



Università degli Studi Mediterranea di Reggio Calabria
Archivio Istituzionale dei prodotti della ricerca

Single and combined abiotic stressors affect maize rhizosphere bacterial microbiota

This is the peer reviewed version of the following article:

Original

Single and combined abiotic stressors affect maize rhizosphere bacterial microbiota / Vescio, R., Malacrino, A., Bennett, A.E., Sorgona', A.. - In: RHIZOSPHERE. - ISSN 2452-2198. - 17:(2021), p. 100318. [10.1016/j.rhisph.2021.100318]

Availability:

This version is available at: <https://hdl.handle.net/20.500.12318/118990> since: 2024-09-02T05:34:54Z

Published

DOI: <http://doi.org/10.1016/j.rhisph.2021.100318>

The final published version is available online at: <https://www.sciencedirect>.

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website

Publisher copyright

This item was downloaded from IRIS Università Mediterranea di Reggio Calabria (<https://iris.unirc.it/>) When citing, please refer to the published version.

(Article begins on next page)

27 July 2026

This is the peer reviewed version of the following article: **Rosa Vescio, Antonino Malacrinò, Alison E. Bennett, Agostino Sorgonà, Single and combined abiotic stressors affect maize rhizosphere bacterial microbiota, Rhizosphere, Volume 17, 2021, 100318, <https://doi.org/10.1016/j.rhisph.2021.100318>**. The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

Single and combined abiotic stressors affect maize rhizosphere bacterial microbiota

Rosa Vescio ^{a,b}, Antonino Malacrinò ^{b,*}, Alison E. Bennett ^b, Agostino Sorgonà ^a

^a*Dipartimento AGRARIA, Università "Mediterranea" di Reggio Calabria, Feo di Vito, 89124, Reggio Calabria, Italy*

^b*Department of Evolution, Ecology and Organismal Biology, The Ohio State University, Columbus, OH, USA*

Keywords: Drought, Heat, Root class, Root zone, 16S rRNA, Metabarcoding

Abstract

Rhizosphere microbiomes are influenced by abiotic stresses, but we know a little about their response to combinations of stresses. In this study we tested: (i) if drought and heat stress influence the maize rhizosphere microbial community; (ii) if the combination of drought and heat has a different outcome compared to a single stress; (iii) if rhizosphere microbiota clusters according to root class and root zone. We setup a microcosm system using maize as model plant. We exposed plants to drought, heat stress and their combination, and used 16S amplicon-sequencing to reconstruct bacterial communities of different root classes (crown and primary) and root zones (apical, sub-apical and basal). We found both drought and heat affect the structure of rhizosphere bacterial communities. The combination of these stressors also influenced the structure of rhizosphere microbial communities, but this effect did not differ compared to the single stresses. Interestingly, we found differences in microbial communities inhabiting the rhizosphere of crown and primary roots in the control treatment, but this difference disappeared once stresses were applied. Stress also lead to an increased abundance of beneficial organisms.

Introduction

Abiotic stressors are a major limiting factor for crop production worldwide (Mantri et al., 2012; Wien 2020). Variation in water availability and increasing of global air temperature are major abiotic stresses posed by climate changes (IPCC 2019). As result, the increase in air temperature and drought events are likely to become more frequent and severe (Spinoni et al., 2018). Recent development in plant microbiome research highlighted the potential of plant-associated microbial communities in alleviating the negative effects of changes in water availability and air temperatures (Hussain et al. 2018; Naylor and

Coleman-Derr 2018; Saikkonen et al. 2020).

Plants play an active role in selecting their own microbiota and can recruit beneficial organisms in response to stresses, especially in the rhizosphere (Berendsen et al. 2012; Turner et al. 2013; Rolfe et al. 2019). Both drought (Naylor and Coleman-Derr 2018) and high air temperatures (van der Voort et al., 2016) can produce a change in the structure of rhizosphere microbiomes. For example, the rhizosphere of plants under drought stress is enriched with plant growth promoting bacteria mainly belonging to the classes Actinobacteria and Firmicutes (Marasco et al., 2012; Edwards et al., 2018; Fitzpatrick et al., 2018; Xu et al., 2018; Simmons et al., 2020). While drought can influence the rhizosphere microbiome directly (via reduction of available water) and indirectly (via the host plant), air temperature only influences the rhizosphere microbiome indirectly (via the host plant). These two different pathways are likely to produce big differences in rhizosphere responses; however, to date, the differential response of rhizosphere microbiomes to heat and water stress has not yet been determined. Furthermore, the combination of heat and water stress commonly occurs in field conditions, and this has an additive detrimental effect on plant growth (Pandey et al., 2017). However, the combination of drought and heat on rhizosphere microbiomes has yet to be examined. The plant root system comprises different root classes, usually classified accordingly to their ontogenesis (i.e. primary, nodal, lateral), each one characterized by distinct developmental, physiological and functional signatures (Waisel and Eshel 2002; Hodge et al., 2009; Tai et al., 2016), and different responses to environmental stresses determining a large within-root phenotypic plasticity (Vescio et al., 2021). For example, different root classes vary in their response to nutrient deficiency (Rubio et al. 2004; Sorgonà et al. 2005; Sorgonà et al., 2007), allelopathy (Abenavoli et al., 2004; Lupini et al., 2016) and drought (Romano et al., 2013; Abenavoli et al., 2016). In addition to this diversification among root classes, roots show differences in functionality in different root zones (Rubio et al., 2004; Sorgonà et al., 2010, 2011). Along the root axis, morphological and functional differences in root architecture reflect the relationship between roots and their environment, with variation in nutrient uptake, water transport, carbon exudation, proton/hydroxyl excretion and respiration (Hodge et al., 2009). Although the physiological and morphological differences between root classes and in different root zones are widely reported, their microbiome remains currently little explored. In this study, we exposed maize plants to drought and heated air, alone and combined, and we characterized the rhizosphere bacterial community in three root zones (apical, subapical and basal) for two different root classes (primary and crown). We hypothesize

that drought and increased air temperatures influence the composition of rhizosphere bacterial microbiome differently due to the difference in direct and in - direct influences on bacterial microbiome composition, and this response will vary with root class and zone. We also hypothesize that the combination of drought and heat will produce a unique rhizosphere microbiome signature.

Material and methods

Study system

Soil was collected from the top 10 cm layer of an uncultivated field located at the Agricultural Experimental Station of the University of Reggio Calabria (38.08N, 15.68E – Tab. S1), and coarsely sieved (4 mm mesh). Pots were filled with a mix of 1 part collected field soil and 2 parts of quartz sand ($\varnothing$ 1–2 mm, Croci Trading Company s.r.l., Italy, autoclaved for 3 h at 121 °C, allowed to cool overnight and then other 3 h at 121 °C). Maize seeds (genotype KXB7554, provided by KWS Italia S.p.A.) were surface sterilized with 20% bleach solution for 20 min and rinsed with deionized water 5 times. Seed germination was synchronized by soaking seeds in deionized water for 24 h and providing air flow through an air pump. We selected maize as model species because: (i) it is an economically important crop and a model species for research; (ii) it has been used as model for abiotic stress research; (iii) it has well-defined root classes and zones.

Experimental design and sample collection

We conducted an experiment testing the effects of two levels of water availability nested in two levels of air temperature on the diversity and composition of the rhizosphere microbiome. Maize plants (*Zea mays* L.) were exposed to two levels of air temperature (25 °C and 32 °C), and two levels of water availability (30% and 80% of soil field capacity, corresponding to a severe drought and no drought). The experiment was split into two blocks to account for the variability introduced by working with two climatic chambers. Each block contained 3 replicates and were temporally distinct: once the first block was harvested the second block was set up, and the air temperature treatments were inverted between the two chambers. The entire experiment yielded a total of 4 treatments (2 water

availability treatments $\times$ 2 air temperature treatments) $\times$ 6 (replicates) = 24 plants. Pairs of plants exposed to the same treatment in the two different blocks were grouped together, and for both we collected two root classes (primary and crown root) that were divided into three zones (apical, sub-apical and basal) yielding a total of 72 samples.

To start each block, after 24 h of soaking, 5 maize seeds were sown in 1 L pots filled with the soil mix. Three replicates of each treatment were then randomly distributed within one of the two climatic chambers (according to the temperature treatment) both initially set at 25 °C, 70% relative humidity and a 14:10 light:dark photoperiod and left to grow for 2 weeks.

During this timeframe each pot was weighted every two days and watered to guarantee a minimum of 80% of soil field capacity. Fifteen days after sowing, plants were exposed to the 30% field capacity and 32 °C air temperature treatments (or both) for 7 days. Heat stress was applied by increasing the air temperature to 32 °C (Hussain et al., 2019) in one climatic chamber. Drought stress was imposed at 30% soil field capacity (Hussain et al., 2019) by reducing water availability from 80% to 30% of the pot capacity (determined by weighing the pots) (Anderson et al. 2018) for plants in both climatic chambers. Control plants were kept at 25 °C and 80% of soil field capacity. With preliminary trials, we determined the amount of water necessary to reach 30% and 80% of field capacity on the same soil used for this experiment. During the experiment, we maintained 30% or 80% of field capacity (according to the treatment) by weighting pots twice a day, calculating the difference in weight compared to our target (either 30% or 80% of field capacity), and compensating this difference with distilled water.

At the end of stress exposure, we measured photosynthetic rate and stomatal conductance to confirm that treated plants were actually stressed (see Supplementary material, Note S1). Plants were then removed from pots and gently shaken to remove bulk soil. The root system was divided into primary and crown roots. Each root class was then divided into three zones: apical (portion from the root tip to the first lateral root), subapical (following the apical portion, same length, but including lateral roots) and basal (same length as the others but excised starting from stem). To extract rhizosphere soil, root sections were put in a 2 ml tube containing 300 μ l of lysis buffer (10 mM Tris, 100 mM NaCl, 10 mM EDTA, 0.5% SDS), and vortexed at maximum speed for 2 min (McPherson et al., 2018). Roots were then discarded, and rhizosphere samples were stored at -80°C before being processed using 16S rRNA metabarcoding procedures (Abdelfattah et al., 2018).

DNA extraction and library preparation

Samples were homogenized with the lysis buffer using two 1 mm $\varnothing$ stainless steel beads per tube, with the aid of a bead mill homogenizer set at 30 Hz for 2 min (TissueLyzer II, Qiagen, USA). Total DNA was extracted using a phenol-chloroform protocol. DNA quality and quantity were checked with a Nanodrop 2000 instrument (Thermo Fisher Scientific Inc., USA). DNA was extracted also from non-template control samples, where experimental samples were replaced by 100 μ l of nuclease-free water, in order to account for contamination of reagents or instruments. PCR amplifications were performed in a reaction mixture containing ~20 ng of template DNA, 1X KAPA HiFi HotStart ReadyMix (KAPA Biosystems, USA), 0.5 μ M of 515 F and 806 R primers (Apprill et al., 2015; Parada et al. 2016), and nuclease free water was added to create a final volume of 12.5 μ L. Amplifications were performed in a Master- cycler Ep Gradient S (Eppendorf, Germany) with an initial denaturation at 94 °C for 3 min, followed by 35 cycles of denaturation at 94 °C for 45 s, 50 °C for 60 s, 72 °C for 90 s and a final extension step at 72 °C for 10 min. All PCR reactions included three non-template control wells, where DNA was replaced with nuclease-free water to check for contamination of PCR reagents. PCR products were inspected for correct amplification on 1% agarose gel. PCR products were then purified with Agencourt AMPure XP kit (Beckman and Coulter, Brea, CA, USA), following the manufacturer's instruction. A short-run PCR was performed on purified samples in order to include the Illumina i7 and i5 indices using the producer's protocol (Nextera XT, Illumina, San Diego, CA, USA). Amplicons were purified again with Agencourt AMPure XP kit as reported above and their concentration was quantified using a Qubit 3.0 fluorometer (Thermo Fisher Scientific Inc., USA). Samples were pooled together at equimolar rations and sequenced with an Illumina MiSeq platform (Illumina, San Diego, CA, USA) using the 300 PE chemistry. No non-template control sample yielded a band after PCR, and the few reads retrieved from sequencing did not pass quality filtering.

Raw reads processing

De-multiplexed forward and reverse reads were merged using the PEAR 0.9.1 algorithm using default parameters (Zhang et al., 2014). Data handling was carried out using QIIME

1.9.1 (Caporaso et al., 2012), and we quality-filtered reads using default parameters, discarded chimeric sequences and binned OTUs with VSEARCH 2.14.2 (Rognes et al., 2016). OTUs coming from amplification of chloroplast DNA were discarded from the downstream analyses. Taxonomy was assigned to each OTU through the BLAST method by querying the SILVA database (v. 132) (Quast et al., 2012).

Data analysis

Data analysis was performed using R statistical software 3.5 (R Core Team, 2020) with the package *phyloseq* (McMurdie and Holmes 2013). Shannon index was calculated using the package *vegan* (Dixon 2003), and comparison among groups was performed by fitting a linear mixed-effect model to account for the nested design, specifying the formula $\sim \text{treatment} * \text{root_class} * \text{root_zone} * (1|\text{root_class}/\text{root_zone}) + (1|\text{temperature}/\text{water})$.

Models were fit using the *lmer* () function under the *lme4* package (Bates et al., 2015) and the package *emmeans* was used to infer pairwise contrasts (corrected using False Discovery Rate, FDR).

Furthermore, we studied the effects of the same factors on the structure of the microbial communities using a multivariate approach. Distances between pairs of samples, in terms of community composition, were calculated using a Bray-Curtis matrix, and then visualized using Canonical Analysis of Principal Coordinates (CAP) procedure. Differences between groups were inferred through PERMANOVA multivariate analysis (999 permutations, stratified using the factors “root_class/ root_zone” and “temperature/water”). Pairwise comparisons were calculated using a custom script, correcting *P* values using the FDR method. We used the R package *DESeq2* (Love et al. 2014) to search for OTUs differentially abundant between each treatment groups and the control. First, we built a model using *treatment*, *root class* and *root zone* as factors, and then we extracted the appropriate contrasts (Stress/Control) for each treatment group. OTUs significantly more abundant in the stressed group were identified by filtering the contrast table by $\log_2\text{FoldChange} > 1$ and $P_{\text{adj}} < 0.05$.

Results

The linear mixed-effects model analysis revealed that microbial rhizosphere Shannon

diversity only varied significantly with the interaction between treatment and root class (Table 1). Specifically, primary roots had a higher diversity than crown roots ($P = 0.003$, Fig. 1B), but this difference disappeared with reduced water availability and increased temperature ($P > 0.05$, Fig. 1B).

We used a multivariate approach (PERMANOVA) to test the influence of treatments, root class, and root zone on the structure of microbial communities, and we found similar results to microbial diversity. We found a significant effect of treatment ($P = 0.005$) and the interaction between treatment and root class ($P < 0.001$) on the structure of rhizosphere microbiome in our experiment (Table 2). In both primary and crown roots, we consistently found that the microbial community of stressed plants was different from control ones and, also, between drought and heat plants (Tab. S2 and S3). However, the rhizosphere microbial community of plants exposed to both single stressors (drought and heat) was not different from the one of plant exposed to their combination (Tab. S2 and S3). Furthermore, for each treatment group, we tested for differences between root classes. We found that the microbiota of primary and crown roots was different in control ($P = 0.001$), but not in any treatment ($P > 0.05$, Tab. S4).

Given the differential response of rhizosphere microbiome to plant stress, we took a closer look to the bacterial taxa that were significantly more abundant in stressed plant compared to the control group (Fig. 2). As general response among the three stresses, we found an increase of three bacterial groups: *Solirubrobacter*, *Massilia*, *Agrobacterium*. While we did not found a specific response to the drought treatment, the treatment with heated air increased the abundance of: *Blastococcus*, *Bosea* (2 OTUs), *Burkholderia*, *Caulobacter* (3 OTUs), *Conexibacter*, *Dactylospor- angium*, *Flavisolibacter*, *Leptothrix*, *Massilia* (5 OTUs), *Mesorhizobium*, *Micromonospora* (3 OTUs), *Niastella*, *Phenoylbacterium*, *Pseudomonas*, *Segetibacterium*, *Solirubrobacter* and 5 unidentified OTUs. On the other hand, when comparing the plant exposed to combined stress to the control, we found a higher abundance of *Rhizobacter*, and 2 unidentified OTUs. Furthermore, 11 OTUs were more abundant in both heat and combined stress treatments: *Acidovorax*, *Bryobacter*, *Massilia*, *Para- burkholderia*, *Pelomonas*, *Rubrobacter*, *Sphingomonas* (2 OTUs), and 3 unidentified OTUs. Two OTUs had a higher abundance in both heat and drought treatments: *Pelomonas* and *Parabulkholderia*.

Discussion

Here, we show that drought and heat stresses induce changes on maize rhizosphere bacterial microbiome. Previous studies, indeed, revealed the effect of various environmental stress on rhizosphere microbiome: drought (Marschner et al. 2005; Cherif et al., 2015; Nuccio et al., 2016; Naylor et al., 2017; Santos-Medellín et al., 2017; Fitzpatrick et al., 2018; Timm et al., 2018), metal-deficiency (Timm et al., 2018), shading (Timm et al., 2018), and nitrogen-deficiency (Allison and Martiny 2008; Roesch et al., 2008; Zhu et al. 2016). Our study supports the effects of plant stressors on rhizosphere microbial assemblages. Furthermore, we tested, for the first time, whether the combination of drought and heat stress produces a different outcome on the rhizosphere microbiome compared to the single stressors. While stress combination was different when compared to the control group, it was not different from the effects of single stressors.

Our results also showed that, in the control group, the rhizosphere of primary and crown roots is inhabited by different bacterial communities. Few previous studies, mostly based on total count and/or trophic strategy and/or culture-dependent techniques only, suggested that different root classes are associated with a different microbiome in the rhizosphere (Gochnauer et al. 1989; De Leij, Whipps and Lynch 1994; Marschner et al. 2005) and, in particular, Sivasithamparam et al. (1979) reported that maize adventitious and seminal roots have similar diversity of bacteria, but the adventitious roots have lower fungal diversity. The only paper focused on root classes soil-based microbiomes, reported nodal roots of *Brachypodium distacum* showing a different structure of bacterial and fungal communities compared to seminal roots (Kawasaki et al., 2016). However, Kawasaki et al. (2016) focused on non-stressed plants while our study included single and combined abiotic stress. Our results supported the differentiation of rhizosphere bacterial microbiome according to root classes. Interestingly, if we extend the same analysis to the stress treatments, the difference between root classes disappears. Currently, we have a very narrow knowledge on this topic, so it is hard to outline an explanatory framework for this effect. We are confident that future studies can expand our results and provide a mechanistic explanation.

Here we also tested the hypothesis that different root zones, within each root class, would be associated to different microbial communities. In our study, we did not find differences in microbial community composition between different zones of the same root class. Following the evidence that different root zones produce different exudates (Walker et al., 2003), we would expect to observe differences in microbial assembly along the root axis. To our knowledge, a single previous study tested the hypothesis that the rhizosphere

microbiome associated with different root zones would respond differently to drought stress (Simmons et al., 2020), and they also found no differences between root zones. Previous studies showed that rhizosphere microorganisms can quickly assimilate root exudates, buffering their influence on rhizosphere microbiomes (Dennis et al. 2010). This mechanism might explain our results, although we should also consider a possible caveat of our study. The lateral roots that stem from sub-apical and basal zones have elongation regions that are morphologically and functionally similar to the apical zone. The presence of these lateral roots might have confounded differences in the microbiome composition between the root zones we targeted.

While each stressor influenced the rhizosphere microbiome in a different way, the analysis of microbial community highlighted that in the control group the microbial community differed between crown and primary roots, but this difference was not found in any of the stressed groups. Previous research found that plants can recruit beneficial microbes in the rhizosphere in an effort to alleviate stress (Lareen et al. 2016), and this can be the mechanism behind our observation. To test this possibility, we focused on the taxa that become significantly more abundant as consequence of plant stress. Indeed, we found that several microbial taxa that were differentially more abundant in our treatment groups are actually associated to plant beneficial organisms: *Massilia* (Ofek et al. 2012), *Solirubrobacter* (Yang et al., 2012; Franke-Whittle et al., 2015), *Burkholderia* (Sua'ez-Moreno et al., 2012), *Caulobacter* (Luo et al., 2019), *Mesorhizobium* (Laranjo et al. 2014), *Micromonospora* (Martínez-Hidalgo et al., 2015), *Rhizobacter* (Lugtenberg and Kamilova 2009), *Paraburkholderia* (Kaur et al. 2017), *Sphingomonas* (Khan et al. 2014, 2017). Furthermore, we found a higher abundance of three genera, *Agrobacterium*, *Pseudomonas* and *Acidovorax*, which are widely known host both pathogenic but also beneficial bacterial species.

Our study brings a novel view to the ecology of plant-associated microorganisms. We showed that root class is an important factor in shaping the rhizosphere bacterial microbiome, and that the presence of plant stressors reduces the differences between root classes. Although more studies on a large set of plant species and genotypes are necessary, our results can contribute in increasing the predictability of plant-microbe relationship, which is an important interaction for securing the productivity of our crops. More generally, our results contribute to the knowledge on the effects of climate changes on crops, showing that two of the major plant stressors caused by climate change influence the plant-microbiome interactions. This has potential impact on the current trend of crafting agricultural practices around a holistic vision of plants-microbe-environment

interactions.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

We are thankful to KWS Italy for providing the maize seeds. We also acknowledge funding from the PhD course “Scienze Agrarie, Alimentari e Forestali” of the Università “Mediterranea” di Reggio Calabria and Dipartimento AGRARIA for supporting RV.

References

- Abdelfattah, A., Malacrino, A., Wisniewski, M., et al., 2018. Metabarcoding: a powerful tool to investigate microbial communities and shape future plant protection strategies. *Biol. Contr.* 120, 1–10.
- Abenavoli, M.R., Leone, M., Sunseri, F., et al., 2016. Root phenotyping for drought tolerance in bean landraces from Calabria (Italy). *J. Agron. Crop Sci.* 202, 1–12.
- Abenavoli, M.R., Sorgonà, A., Albano, S., et al., 2004. Coumarin differentially affects the morphology of different root types of maize seedlings. *J. Chem. Ecol.* 30, 1871–1883.
- Allison, S.D., Martiny, J.B.H., 2008. Resistance, resilience, and redundancy in microbial communities. *Proc. Natl. Acad. Sci. Unit. States Am.* 105, 11512–11519.
- Anderson, S.M., Puertolas, J., Dodd, I.C., 2018. Does irrigation frequency affect stomatal response to drying soil? *Acta Hort.* 1197, 113–138.
<https://doi.org/10.17660/ActaHortic.2018.1197.18>.
- Apprill, A., McNally, S., Parsons, R., et al., 2015. Minor revision to V4 region SSU rRNA 806R gene primer greatly increases detection of SAR11 bacterioplankton. *Aquat. Microb. Ecol.* 75, 129–137.

334 Bates, D., Machler, M., Bolker, B., et al., 2015. Fitting linear mixed-effects models
335 using lme4. *J. Stat. Software* 67, 1–48.

336 Berendsen, R.L., Pieterse, C.M.J., Bakker, P.A.H.M., 2012. The rhizosphere
337 microbiome and plant health. *Trends Plant Sci.* 17, 478–486.

338 Caporaso, J.G., Lauber, C.L., Walters, W.A., et al., 2012. Ultra-high-throughput
339 microbial community analysis on the Illumina HiSeq and MiSeq platforms. *ISME J.* 6,
340 1621–1624.

341 Cherif, H., Marasco, R., Rolli, E., et al., 2015. Oasis desert farming selects
342 environment-specific date palm root endophytic communities and cultivable bacteria
343 that promote resistance to drought. *Environ Microbiol Rep* 7, 668–678.

344 Dennis, P.G., Miller, A.J., Hirsch, P.R., 2010. Are root exudates more important than
345 other sources of rhizodeposits in structuring rhizosphere bacterial communities?
346 *FEMS Microbiol. Ecol.* 72, 313–327.

347 Dixon, P., 2003. VEGAN, a package of R functions for community ecology. *J. Veg. Sci.*
348 14, 927–930.

349 Edwards, J.A., Santos-Medellín, C.M., Liechty, Z.S., et al., 2018. Compositional shifts
350 in root-associated bacterial and archaeal microbiota track the plant life cycle in field-
351 grown rice. In: Gore, J. (Ed.), *PLOS Biol*, vol. 16, e2003862.

352 Fitzpatrick, C.R., Copeland, J., Wang, P.W., et al., 2018. Assembly and ecological
353 function of the root microbiome across angiosperm plant species. *Proc. Natl. Acad.*
354 *Sci. Unit. States Am.* 115, E1157–E1165.

355 Franke-Whittle, I.H., Manici, L.M., Insam, H., et al., 2015. Rhizosphere bacteria and
356 fungi associated with plant growth in soils of three replanted apple orchards. *Plant Soil*
357 395, 317–333.

358 Gochnauer, M.B., McCully, M.E., Labbé, H., 1989. Different populations of bacteria
359 associated with sheathed and bare regions of roots of field-grown maize. *Plant Soil* 114,
360 107–120.

361 Hodge, A., Berta, G., Doussan, C., et al., 2009. Plant root growth, architecture and
362 function. *Plant Soil* 321, 153–187.

363 Hussain, H.A., Men, S., Hussain, S., et al., 2019. Interactive effects of drought and
364 heat stresses on morpho-physiological attributes, yield, nutrient uptake and oxidative

status in maize hybrids. *Sci. Rep.* 9, 3890.

Hussain, S.S., Mehnaz, S., Siddique, K.H.M., 2018. Harnessing the Plant Microbiome for Improved Abiotic Stress Tolerance, pp. 21–43.

IPCC, 2019. Climate Change and Land: an IPCC Special Report on Climate Change, Desertification, Land Degradation, Sustainable Land Management, Food Security, and Greenhouse Gas Fluxes in Terrestrial Ecosystems.

Kaur, C., Selvakumar, G., Ganeshamurthy, A.N., 2017. Burkholderia to Paraburkholderia: the Journey of a Plant-Beneficial-Environmental Bacterium. *Recent Advances in Applied Microbiology*. Springer Singapore, Singapore, pp. 213–228.

Kawasaki, A., Donn, S., Ryan, P.R., et al., 2016. Microbiome and exudates of the root and rhizosphere of *Brachypodium distachyon*, a model for wheat. In: Budak, H. (Ed.), *PLoS One*, vol. 11, e0164533.

Khan, A.L., Waqas, M., Asaf, S., et al., 2017. Plant growth-promoting endophyte *Sphingomonas* sp. LK11 alleviates salinity stress in *Solanum pimpinellifolium*. *Environ. Exp. Bot.* 133, 58–69.

Khan, A.L., Waqas, M., Kang, S.-M., et al., 2014. Bacterial endophyte *Sphingomonas* sp. LK11 produces gibberellins and IAA and promotes tomato plant growth. *J. Microbiol.* 52, 689–695.

Laranjo, M., Alexandre, A., Oliveira, S., 2014. Legume growth-promoting rhizobia: an overview on the *Mesorhizobium* genus. *Microbiol. Res.* 169, 2–17.

Lareen, A., Burton, F., Scha"fer, P., 2016. Plant root-microbe communication in shaping root microbiomes. *Plant Mol. Biol.* 90, 575–587.

De Leij, F.A.A.M., Whipps, J.M., Lynch, J.M., 1994. The use of colony development for the characterization of bacterial communities in soil and on roots. *Microb. Ecol.* 27, 81–97.

Love, M.I., Huber, W., Anders, S., 2014. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol.* 15, 550.

Lugtenberg, B., Kamilova, F., 2009. Plant-growth-promoting rhizobacteria. *Annu. Rev. Microbiol.* 63, 541–556.

Luo, D., Langendries, S., Mendez, S.G., et al., 2019. Plant growth promotion driven by

396 a novel *Caulobacter* strain. *Mol. Plant Microbe Interact.* 32, 1162–1174.

397 Lupini, A., Sorgon`a, A., Princi, M.P., et al., 2016. Morphological and physiological
 398 effects of trans-cinnamic acid and its hydroxylated derivatives on maize root types.
 399 *Plant Growth Regul.* 78, 263–273.

400 Mantri, N., Patade, V., Penna, S., et al., 2012. Abiotic Stress Responses in Plants:
 401 Present and Future. *Abiotic Stress Responses in Plants*. New York, NY. Springer, New
 402 York, pp. 1–19.

403 Marasco, R., Rolli, E., Ettoumi, B., et al., 2012. A drought resistance-promoting
 404 microbiome is selected by root system under desert farming. In: Gilbert, J.A. (Ed.),
 405 *PLoS One*, vol. 7, e48479.

406 Marschner, P., Grierson, P.F., Rengel, Z., 2005. Microbial community composition
 407 and functioning in the rhizosphere of three *Banksia* species in native woodland in
 408 Western Australia. *Appl. Soil Ecol.* 28, 191–201.

409 Martínez-Hidalgo, P., Galindo-Villardó'n, P., Trujillo, M.E., et al., 2015.
 410 *Micromonospora* from nitrogen fixing nodules of alfalfa (*Medicago sativa* L.). A new
 411 promising Plant Probiotic Bacteria. *Sci. Rep.* 4, 6389.

412 McMurdie, P.J., Holmes, S., 2013. phyloseq: an R package for reproducible interactive
 413 analysis and graphics of microbiome census data. In: Watson, M. (Ed.), *PLoS One*, vol.
 414 8, e61217.

415 McPherson, M.R., Wang, P., Marsh, E.L., et al., 2018. Isolation and analysis of
 416 microbial communities in soil, rhizosphere, and roots in perennial grass experiments.
 417 *JoVE*. <https://doi.org/10.3791/57932>.

418 Naylor, D., Coleman-Derr, D., 2018. Drought stress and root-associated bacterial
 419 communities. *Front. Plant Sci.* 8 <https://doi.org/10.3389/fpls.2017.02223>.

420 Naylor, D., DeGraaf, S., Purdom, E., et al., 2017. Drought and host selection influence
 421 bacterial community dynamics in the grass root microbiome. *ISME J.* 11, 2691–2704.

422 Nuccio, E.E., Anderson-Furgeson, J., Estera, K.Y., et al., 2016. Climate and edaphic
 423 controllers influence rhizosphere community assembly for a wild annual grass.
 424 *Ecology* 97, 1307–1318.

425 Ofek, M., Hadar, Y., Minz, D., 2012. Ecology of root colonizing *Massilia*
 426 (*oxalobacteraceae*). In: Vinatzer, B.A. (Ed.), *PLoS One*, vol. 7, e40117.

427 Pandey, P., Irulappan, V., Bagavathiannan, M.V., et al., 2017. Impact of combined

428 abiotic and biotic stresses on plant growth and avenues for crop improvement by
 429 exploiting physio-morphological traits. *Front. Plant Sci.* 8 <https://doi.org/10.3389/fpls.2017.00537>.

431 Parada, A.E., Needham, D.M., Fuhrman, J.A., 2016. Every base matters: assessing
 432 small subunit rRNA primers for marine microbiomes with mock communities, time
 433 series and global field samples. *Environ. Microbiol.* 18, 1403–1414.

434 Quast, C., Pruesse, E., Yilmaz, P., et al., 2012. The SILVA ribosomal RNA gene
 435 database project: improved data processing and web-based tools. *Nucleic Acids Res.*
 436 41, D590–D596.

437 Roesch, L.F.W., Camargo, F.A.O., Bento, F.M., et al., 2008. Biodiversity of
 438 diazotrophic bacteria within the soil, root and stem of field-grown maize. *Plant Soil*
 439 302, 91–104.

440 Rognes, T., Flouri, T., Nichols, B., et al., 2016. VSEARCH: a versatile open source tool
 441 for metagenomics. *PeerJ* 4, e2584.

442 Rolfe, S.A., Griffiths, J., Ton, J., 2019. Crying out for help with root exudates: adaptive
 443 mechanisms by which stressed plants assemble health-promoting soil microbiomes.
 444 *Curr. Opin. Microbiol.* 49, 73–82.

445 Romano, A., Sorgonà, A., Lupini, A., et al., 2013. Morpho-physiological responses of
 446 sugar beet (*Beta vulgaris* L.) genotypes to drought stress. *Acta Physiol. Plant.* 35, 853–
 447 865.

448 Rubio, G., Sorgonà, A., Lynch, J.P., 2004. Spatial mapping of phosphorus influx in
 449 bean root systems using digital autoradiography. *J. Exp. Bot.* 55, 2269–2280.

450 Saikkonen, K., Nissinen, R., Helander, M., 2020. Toward comprehensive plant
 451 microbiome research. *Front Ecol Evol* 8. <https://doi.org/10.3389/fevo.2020.00061>.

452 Santos-Medellín, C., Edwards, J., Liechty, Z., et al., 2017. Drought stress results in a
 453 compartment-specific restructuring of the rice root-associated microbiomes. In:
 454 Ausubel, F.M. (Ed.), *MBio*, vol. 8. <https://doi.org/10.1128/mBio.00764-17>.

455 Simmons, T., Styer, A.B., Pierroz, G., et al., 2020. Drought drives spatial variation in the
 456 millet root microbiome. *Front. Plant Sci.* 11 <https://doi.org/10.3389/fpls.2020.00599>.

458 Sivasithamparam, K., Parker, C.A., Edwards, C.S., 1979. Rhizosphere micro-organisms
 459 of seminal and nodal roots of wheat grown in pots. *Soil Biol. Biochem.* 11, 155–160.

460 Sorgonà, A., Abenavoli, M.R., Gringeri, P.G., et al., 2007. Comparing morphological
 461 plasticity of root orders in slow- and fast-growing citrus rootstocks supplied with
 462 different nitrate levels. *Ann. Bot.* 100, 1287–1296.

463 Sorgonà, A., Abenavoli, M.R., Cacco, G., 2005. A comparative study between two
 464 citrus rootstocks: effect of nitrate on the root morpho-topology and net nitrate uptake.
 465 *Plant Soil* 270, 257–267.

466 Sorgonà, A., Cacco, G., Di Dio, L., et al., 2010. Spatial and temporal patterns of net
 467 nitrate uptake regulation and kinetics along the tap root of *Citrus aurantium*. *Acta*
 468 *Physiol. Plant.* 32, 683–693.

469 Sorgonà, A., Lupini, A., Mercati, F., et al., 2011. Nitrate uptake along the maize primary
 470 root: an integrated physiological and molecular approach. *Plant Cell Environ.* 34,
 471 1127–1140.

472 Spinoni, J., Vogt, J.V., Naumann, G., et al., 2018. Will drought events become more
 473 frequent and severe in Europe? *Int. J. Climatol.* 38, 1718–1736.

474 Suarez-Moreno, Z.R., Caballero-Mellado, J., Coutinho, B.G., et al., 2012. Common
 475 features of environmental and potentially beneficial plant-associated *Burkholderia*.
 476 *Microb. Ecol.* 63, 249–266.

477 Tai, H., Lu, X., Opitz, N., et al., 2016. Transcriptomic and anatomical complexity of
 478 primary, seminal, and crown roots highlight root type-specific functional diversity in
 479 maize (*Zea mays* L.). *J. Exp. Bot.* 67, 1123–1135.

480 Timm, C.M., Carter, K.R., Carrell, A.A., et al., 2018. Abiotic stresses shift belowground
 481 populus-associated bacteria toward a Core stress microbiome. In: Herr, J.R. (Ed.),
 482 *mSystems*, vol. 3. <https://doi.org/10.1128/mSystems.00070-17>.

483 Turner, T.R., James, E.K., Poole, P.S., 2013. The plant microbiome. *Genome Biol.* 14,
 484 209.

485 van der Voort, M., Kempenaar, M., van Driel, M., et al., 2016. Impact of soil heat on
 486 reassembly of bacterial communities in the rhizosphere microbiome and plant disease
 487 suppression. In: Classen, A. (Ed.), *Ecol Lett*, vol. 19, pp. 375–382.

488 Vescio, R., Abenavoli, MR, Sorgona`, A, 2021. Single and Combined Abiotic Stress in
 489 Maize Root Morphology. *Plants* 10 (1), 5. <https://doi.org/10.3390/plants10010005>.

490 Waisel, Y., Eshel, A., 2002. Functional Diversity of Various Constituents of a Single

Root System. Plant Roots. CRC Press, pp. 157–174.

Walker, T.S., Bais, H.P., Grotewold, E., et al., 2003. Root exudation and rhizosphere biology: Fig. 1. Plant Physiol 132, 44–51.

Wien, H.C., 2020. Abiotic Stress Effects on Vegetable Crops. The Physiology of Vegetable Crops. CABI, Wallingford, pp. 71–93.

Xu, L., Naylor, D., Dong, Z., et al., 2018. Drought delays development of the sorghum root microbiome and enriches for monoderm bacteria. Proc. Natl. Acad. Sci. Unit. States Am. 115, E4284–E4293.

Yang, J., Ruegger, P.M., McKenry, M.V., et al., 2012. Correlations between root-associated microorganisms and peach replant disease symptoms in a California soil. In: Yang, C.-H. (Ed.), PLoS One, vol. 7, e46420.

Zhang, J., Kobert, K., Flouri, T., et al., 2014. PEAR: a fast and accurate Illumina Paired-End reAd mergeR. Bioinformatics 30, 614–620.

Zhu, S., Vivanco, J.M., Manter, D.K., 2016. Nitrogen fertilizer rate affects root exudation, the rhizosphere microbiome and nitrogen-use-efficiency of maize. Appl. Soil Ecol. 107, 324–333.

R Core Team (2020). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.

Table 1

Results from the mixed-effect linear model testing the Shannon diversity index against treatment, root class, root zone and their interactions.

Factor	df	F	P
Treatment (T)	3	2.379	0.497
Root class (RC)	1	0.001	0.967
Root zone (RZ)	2	0.028	0.985
T x RC	3	16.171	0.001
T x RZ	6	5.118	0.528
RC x RZ	2	0.994	0.608
T x RC x RZ	6	5.874	0.437

Table 2

Results from PERMANOVA analysis testing the effects of treatment, root class, root zone and their interactions on the structure of maize rhizosphere bacterial communities.

Factor	df	R ²	F	P
Treatment (T)	3	0.065	1.616	0.005
Root type (RC)	1	0.012	0.953	0.491
Root zone (RZ)	2	0.026	0.987	0.451
T x RC	3	0.091	2.267	0.001
T x RZ	6	0.079	0.99	0.501
RC x RZ	2	0.036	1.358	0.065
T x RC x RZ	6	0.095	1.184	0.06




Fig. 1. Comparison of Shannon diversity index between (A) treatments and (B) root classes within treatments. (C) CAP (Canonical Analysis of Principal coordinates) ordination using a Bray-Curtis distance matrix of samples. ** $P = 0.003$.




Fig. 2. Venn diagram representing the number of OTUs (Operational Taxonomic Units) differentially more abundant as response to specific stressors compared to the control group.

Single and combined abiotic stressors affect maize rhizosphere bacterial microbiota

Rosa Vescio, Antonino Malacrino, Alison E. Bennett, Agostino Sorgonà

Rhizosphere (2021), <https://doi.org/10.1016/j.rhisph.2021.100318>

Supplementary material

Note S1. Supplementary results.

The photosynthetic rate and stomatal conductance was measured on intact leaves using the LI-COR LI-6400 system (LI-COR Inc.; Lincoln, NE) with a leaf temperature of 26 °C, a CO₂ concentration of 400 μmol(CO₂) mol(air)⁻¹ (controlled by CO₂ cylinder), an air flow rate of 500 cm³ min⁻¹, and 1200 μmol m⁻²s⁻¹ of photosynthetically active radiation supplied by the LED light source. Each measure was taken between 120 and 200 seconds of waiting time. Between measures, the difference in the CO₂ concentration between the sample and the reference was matched to 50 μmol(CO₂) mol(air)⁻¹. The leaf to-air vapor pressure difference (VPD) was set to 1.5 kPa, and continuously monitored around the leaf during measurements and maintained at a constant level by manipulating the humidity of incoming air as needed. All measurements were performed inside a growth chamber. For each treatment we measured six plants, and for each plant we recorded the mean value of two measures on different leaves. Data was analysed using one-way ANOVA, and contrasts were inferred using Fisher's LSD post hoc test.




Figure S1 – Photosynthetic rate (μmol(CO₂) x m⁻² x s⁻¹) and stomatal conductance (gs, mol H₂O x m⁻² x s⁻¹) of maize plants exposed for seven days to drought (D), heat (H) and their combination (C). The control (O) was obtained in presence of optimal water and temperature (see Materials and Methods). Different letters indicate significant differences between treatments (Fisher's LSD test).

Results show that photosynthetic rate was not influenced by treatments ($F=1.0780$; $df=3$; $P=0.38$ - Fig. S1A), while we observed differences in stomatal conductance between control and all treatments ($F=3.51$; $df=3$; $P=0.034$ - Fig S1B). The stomatal conductance decreased by -20%, -26% and -16% in plants exposed to drought, heat and combined stress, respectively (Fig. S1B). Similar results were obtained in Hussain et al. (2019). This confirms that our treatments were successful in inducing stress to maize plants.

References

Hussain, H.A., Men, S., Hussain, S. et al. (2019) Interactive effects of drought and heat stresses on morpho-physiological attributes, yield, nutrient uptake and oxidative status in maize hybrids. Sci Rep 9, 3890. <https://doi.org/10.1038/s41598-019-40362-7>

Table S1. Soil physical and chemical characteristics

Soil texture	sand 36.0%, silt 32.0%, clay 32.0%
Bulk density	$1.23 \pm 0.04 \text{ kg/dm}^3$
pH_{water}	7.2 ± 0.2
pH_{KCl}	6.4 ± 0.1
Total organic Carbon (C)	$19.3 \pm 0.4 \text{ g/kg dry soil}$
Total Nitrogen (N)	$1.8 \pm 0.2 \text{ g/kg dry soil}$
C:N ratio	10.7
NH₄⁺	$17.1 \pm 1.0 \text{ mg/kg dry soil}$
NO₃⁻	$13.0 \pm 1.0 \text{ mg/kg dry soil}$
Olsen P	$18.3 \pm 2.3 \text{ mg/kg dry soil}$
Total CaCO₃	$8.4 \pm 1.0 \text{ g/kg dry soil}$
Active CaCO₃	$3.9 \pm 0.2 \text{ g/kg dry soil}$
Cation Exchange Capacity (CEC)	$17.1 \pm 1.7 \text{ cmol/kg dry soil}$
Electrical Conductivity (EC) 1:2 at 25°C	$0.165 \pm 0.004 \text{ dS/m}$

Table S2. PERMANOVA pairwise comparison between treatments in primary roots.

Pair	FDR corrected <i>P</i> value
Control – Drought	0.005
Control – Heat	0.001
Control – Combined	0.002
Drought – Heat	0.03
Drought – Combined	0.308
Heat – Combined	0.094

Table S3. PERMANOVA pairwise comparison between treatments in crown roots.

Pair	FDR corrected <i>P</i> value
Control – Drought	0.006
Control – Heat	0.005
Control – Combined	0.004
Drought – Heat	0.03
Drought – Combined	0.253
Heat – Combined	0.073

Table S4. PERMANOVA pairwise comparison between root classes (primary vs crown) within each treatment.

Pair	FDR corrected <i>P</i> value
Control	0.001
Drought	0.200
Heat	0.055
Combined	0.273