



Università degli Studi Mediterranea di Reggio Calabria
Archivio Istituzionale dei prodotti della ricerca

Integrated environmental evaluation of heavy metals and metalloids bioaccumulation in invertebrates and seaweeds from different marine coastal areas of sardinia, mediterranean sea

This is the peer reviewed version of the following article:

Original

Integrated environmental evaluation of heavy metals and metalloids bioaccumulation in invertebrates and seaweeds from different marine coastal areas of sardinia, mediterranean sea / Corrias, F., Atzei, A., Addis, P., Secci, M., Russo, M., Angioni, A.. - In: ENVIRONMENTAL POLLUTION. - ISSN 0269-7491. - 266:Pt 2(2020), p. 115048. [10.1016/j.envpol.2020.115048]

Availability:

This version is available at: <https://hdl.handle.net/20.500.12318/95080> since: 2021-04-11T11:22:46Z

Published

DOI: <http://doi.org/10.1016/j.envpol.2020.115048>

The final published version is available online at: <https://www.sciencedirect>.

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website

Publisher copyright

This item was downloaded from IRIS Università Mediterranea di Reggio Calabria (<https://iris.unirc.it/>) When citing, please refer to the published version.

(Article begins on next page)

Integrated environmental evaluation of heavy metals bioaccumulation in invertebrates and seaweeds from different marine coastal areas of Sardinia, Mediterranean Sea.

Francesco Corrias^a, Alessandro Atzei^a, Piero Addis^b, Marco Secci^b, Mariateresa Russo^c, Alberto Angioni^{a*}

^aDepartment of Life and Environmental Science, Food Toxicology Unit, University of Cagliari, University Campus of Monserrato, SS 554, 09042 Cagliari, Italy,

Francesco Corrias (francesco.corrias@unica.it), Alessandro Atzei (alessandro.atzei@unica.it),

^bDepartment of Life and Environmental Science, Biology Section, University of Cagliari, via Tommaso Fiorelli 2, 09126 Cagliari, Italy, Marco Secci (msecci@unica.it), Piero Addis (addisp@unica.it).

^c Department of Agricultural Science, Mediterranean University of Reggio Calabria, Località Feo di Vito - 89122 Reggio Calabria (RC), Mariateresa Russo (mariateresa.russo@unirc.it)

Running title: Heavy metals accumulation studies in marine coastal areas of Sardinia, Mediterranean Sea.

* Corresponding author:

Alberto Angioni

Food Toxicology Unit, University of Cagliari, University campus of Monserrato, SS 554, 09042 Cagliari, Italy; Phone: +390706758615, Fax: +390706758612; e-mail: aangioni@unica.it

27 **Abstract**

28 In this work, three gastropods *Patella vulgata*, *Osilinus turbinata*, and *Tahis clavigera*, one
29 echinoderm *Paracentrotus lividus*, one coelenterate *Anemonia sulcata*, and two seaweed *Padina*
30 *pavonica*, and *Cystoseira mediterranea* were collected from three different marine areas of Sardinia
31 in the Mediterranean sea and studied for heavy metals and metalloid content and accumulation
32 trends.

33 Inductively coupled plasma optical emission spectrometry (ICP-OES) was used for the
34 determination of Al, AS, B, Ba, Be, Cd, Co, Cr, Cu, Fe, Hg, Mn, Mo, Ni, Pb, Sb, Se, Sn, Sr, Te, Ti,
35 V and Zn in the selected samples. The results showed that *gastropods* were capable of accumulating
36 Al, Ba, Cu, Fe, Sr, and Zn; *seaweeds* can better concentrate Al, Fe, and Zn than all other species. At
37 the same time, *echinoderms* and *coelenterate* had limited ability to store specific metals, showing a
38 much more homogeneous distribution. PCA analysis allowed us to discriminate among the sites and
39 the species. Cala Zafferano was the area with the higher values of accumulation of all metals in all
40 species as expected, considering its proximity to industrial sites. The results of the analysis showed
41 clearly that heavy metal and metalloid accumulation was different for each species studied.
42 Therefore, for a correct environmental assessment of a given area, a comprehensive approach is
43 strongly recommended by exploiting the different properties of both accumulation and
44 concentration of the metals by different aquatic species.

45

46 **Summary**

47 Different species of aquatic living organisms have different heavy metals and metalloid
48 accumulation trends, even if they belong to the same family and the same water column or have
49 similar living habits.

50

51 **Keywords:** Heavy metals; accumulation; gastropods; *Paracentrotus*, *Anemonia*, seaweed, PCA

52

53 **Introduction**

54 Metals are usually found in the environment worldwide, their presence caused by natural events or
55 because of anthropogenic activities, such as industrial sites and agricultural practice
56 (Kamaruzzaman et al., 2011). The increase in environmental metal concentration in these areas can
57 reach 100-1000 times the normal levels of Earth's crust (Carral et al., 1995). The hydrological cycle
58 of water between the land, open water surface and sea can bring to significant pollution, exposing
59 the aquatic ecosystems directly to a great variety of these metals causing severe contamination of
60 marine organisms (Farombi et al., 2007).

61 Metals are mutually essential elements necessary to support biological activities and unneeded
62 chemicals with an unidentified role (Roesijadi and Robinson, 1994). Heavy metals are problematic
63 pollutants because, unlike most organic contaminants, they are non-biodegradable and can
64 accumulate in living tissues and food, causing a significant threat to both human health and
65 environmental safety. The most hazardous heavy metals and metalloids are Be, Cr, Hg, Cd, Pb, and
66 As. These metal ions are low soluble in water, can cause toxicity, and have serious side effects on
67 human health. After discharged in the water, they are usually accumulated by sediments and
68 transferred to invertebrates and biomagnified in fish populations accumulating in the lipid tissue
69 (Fergusson, 1990; Yi et al., 2011; Jakimska, 2011; Jaishankar et al., 2014; Jitar et al. 2015). An
70 accurate assessment of the degree of metal contamination at a specific location requires an adequate
71 estimation of these elements' natural levels in the biological communities and the natural
72 environment. Mussels, as intertidal filter feeders, accumulate pollutants coming from local
73 contamination and distant areas influenced by sea current and thus can lead to inaccurate
74 information on the real environmental situation of restricted territories. Therefore, environmental
75 pollution monitoring studies relying on their ability to concentrate high levels of metal in the soft
76 tissue used mussels, invertebrates, and marine vegetables widely (Philips, 1977; Ingston, 1982;
77 Goldberg et al., 1983; Cossa, 1989; Bryan and Langston, 1992; Abisil et al., 1997; Hummel et al.,
78 1997).

Commentato [IC1]: Ho messo la bibliografia dopo widely.

79 The use of sedentary herbivorous such as gastropods, and echinoderms, seaweeds holdfast, and
80 anemonia has several advantages. These living organisms are available all year, simple to collect,
81 resistant to transportation and conservation, and have enough tissues for the analysis.

82 In the last decades, metals bioaccumulation has been investigated in different biological
83 communities such as invertebrate gastropods (Vinogrijdov et al., 1953; Blackmore, 2000; Perez-
84 Lopez et al., 2003; Amin et al., 2006; Collado et al., 2006; Yüzereroğlu et al., 2009; Ramirez, 2013;
85 Kelepertzis, 2013; Boucetta et al., 2016; Duysak and Azdural, 2017; Aydin-Onen and Ozturk, 2017;
86 Gawad, 2018), anemone (Mitchelmore et al., 2003; Horwitz et al., 2014), or echinoderms (Salvo et
87 al., 2014, Chiarelli et al., 2019). Moreover, some authors investigated seaweed such as *Padina* spp
88 and *Cystoseira* spp (Aydin-Onen and Ozturk, 2017; Rodríguez-Figueroa et al., 2008; Benfares et
89 al., 2015; Ryabushko et al., 2017). However, most articles reported pollution from 4 to 15 metals in
90 one or two species.

91 The study aimed to compare metals accumulation performance of 7 living organisms (invertebrates
92 and vegetables) in three marine areas of Sardinia characterized by the different anthropic impact.
93 Cala Zafferano (CZ) affected by heavy metal contamination, Capo Frasca (CF) and Capo
94 Spartivento (CS), representing potentially clean environments, exposed to NW, SW and SE
95 currents. Three gastropods *Patella vulgata*, *Osilinus turbinata*, and *Tahis clavigera*, one
96 echinoderm *Parancetrotus lividus*, one coelenterate *Anemonia sulcata*, and two seaweed *Padina*
97 *pavonica*, and *Cystoseira mediterranea*, were collected. All samples were subjected to
98 mineralization and analysis by ICP-OES to determine an exhaustive pattern of 23 metals and
99 metalloids and accumulation studies.

100 1. Material and methods

101 2.1 Sample collection and processing

102 Samples were handpicked from three sites of the Sardinia coast, Capo Spartivento (38° 52'41.90 "N;
103 8° 50'28.77 "E), Cala Zafferano Teulada (38° 53'43.26 "N; 8° 39'42.19 "E), and Capo Frasca (39°
104 42'54.98 "N; 8° 26'44.14 "E). Twenty samples of the gastropods *Patella vulgata*, *Osilinus turbinata*,

105 and *Thais haemastoma*, the echinoderm *Parancetritus lividus*, the coelenterate *Anemonia*
106 *sulcata*, and the seaweeds *Padina pavonica*, and *Cystoseira mediterranea*, were collected at a
107 maximum depth of 10 meters in August (Table 1). All species were present at all stations and
108 individually processed.

109 After collection, samples were immediately stored in ice cooler "drain plug open" to let the water
110 run out. This operation avoids deterioration during transport to the lab. Samples brought to the
111 laboratory were weighed, and gastropods soft tissue was separated from the shell. Sea urchin's
112 gonads were collected with a stainless-steel spatula from the shell opened with a stainless-steel
113 cutter for sea urchins. Anemone tentacles and seaweed aerial parts were selected using a stainless-
114 steel scissor. All samples, except gonads, were rinsed with deionized water to remove accidental
115 impurities, and finally stored at -20°C before analysis.

116 *Chemicals*

117 HNO₃ 67-69%, H₂O₂ solution 30%, and standards of Al, AS, B, Ba, Be, Cd, Co, Cr, Cu, Fe, Hg,
118 Mn, Mo, Ni, Pb, Sb, Se, Sn, Sr, Te, Ti, V, Zn were of ICP grade (Carlo Erba Reagents Milan, Italy),
119 HCl 34-37% super pure quality (Romil Spa Cambridge, England). Double-deionized water with a
120 conductivity less than 18.2 MΩ was obtained with a Milli-Q system (Millipore, Bedford, MA,
121 USA). Stock standard solutions were prepared at 1000 mg/L in HNO₃ 10%. Working standard
122 solutions were prepared daily by diluting the stock solution with MilliQ water.

123 *2.2 Moisture and ash*

124 Ten grams of homogenized samples were weighed and dried at 105°C in a thermostatic heater until
125 a constant weight was achieved (~ 24 h). After that, samples were stored in a desiccator to reach
126 ambient temperature and **weighted** for dried mater assessment. Dehydrated samples were then
127 carbonized at 450 C° in a porcelain crucible for five hours for total ash analysis. Moisture and ash
128 determinations were calculated as a percentage of the fresh product.

129

130 *2.3 Heavy metals and metalloids*

Commentato [IC2]: Weighed??

131 About 0.2 g of ashes obtained by carbonization were mineralized in a microwave CEM Mars6
132 system (CEM Corporation, Milano, Italy), by adding 10 mL of a mixture of HNO₃ and HCl (1:3),
133 and 1 mL of H₂O₂. After the mineralization program, the solution was transferred into a 10 mL
134 flask and brought to volume with MilliQ water with a conductivity less than 18.2 MΩ, filtered
135 through 0.45 μm nitrocellulose membrane filter (Whatman, Milan, Italy) in another flask and finally
136 subjected to analysis. Control solvent samples were simultaneously prepared to avoid false positives
137 and contamination during the investigation.

138 Hg, As, and Se sample preparation were carried out using Agilent VGA-77 (Agilent, Milan, Italy)
139 according to Agilent application notes (Agilent AN, 2014; Agilent AN, 2012). Analysis were
140 carried out using a Varian 710ES ICP Optical Emission Spectrometer for the simultaneous analysis
141 of 23 metals and metalloids. Operating conditions were as follows: radiofrequency (RF) generator
142 power 1.2, frequency 40 MHz; argon (99.996% purity) was used both for plasma (15.5 L/min),
143 nebulizer (200 Kpa) and optic supply (1.5 L/min). The sample uptake delay was 30 seconds, and the
144 instrument stabilization delay 15 seconds. The spray chamber was a double-pass, glass cyclonic.
145 Each measurement was made in triplicate. The power and pressure applied were 600 W and 100
146 PSI for 13 minutes. The limit of quantitation (LOQ) was calculated as ten times the standard
147 deviation reading of the blank sample signal (Wenzl et al. 2016). Calibration curves were calculated
148 on five points starting from the LOQ value and were considered acceptable for $R^2 \geq 0.999$.

150 *2.4 Risk assessment*

151 In this study, we used the target hazard quotient (THQ) for risk assessment. THQ represents the
152 ratio of the estimated daily intake (EDI) / reference dose (RfD) for noncarcinogenic risk assessment
153 methods indicating the risk level associated with xenobiotic exposure (Chien et al., 2002; Zhang et
154 al., 2017). RfDs for oral intake were obtained from the Integrated Risk Information System (IRIS-
155 EPA). When THQ is below 1 no adverse effect during a person's lifetime is expected. EDI was
156 calculated as the concentration of heavy metals (mg/kg wet weight) × average daily consumption of

Commentato [IC3]: Mettere 3 al pedice?

Commentato [IC4]: Mettere i 2 al pedice?

Commentato [IC5]: Lo ripetiamo di nuovo dopo i materiali?

Commentato [IC6]: Ho aggiunto questo.

157 invertebrates in the local area (g/day bw) / body weight (Bw). To evaluate the risk of heavy metals
158 and metalloids from invertebrate consumption, we consider an average weight of 60 Kg. Total THQ
159 for each species was calculated as the sum of single metals THQ, while total THQ of each site was
160 calculated assuming the intake from all the selected species in the same day.

161 *2.5 Statistical Analysis*

162 Analysis of variance (ANOVA) was carried out with the software XLSTAT (Addinsolf LTD,
163 Version 19.4), **Mean** comparisons of the effects of treatments were calculated by the Fisher's least
164 significant difference test at $p \leq 0.05$.

165 Analysis data sets were imported into SIMCA 14 (Umetrics AB, Umea, Sweden) for processing
166 using principal component analysis (PCA). Additionally, R2 (the coefficient of determination) and
167 Q2 (the cross-validated correlation coefficient) were used as measures for the robustness of a
168 pattern recognition model (Atherton et al., 2006).

169

170 **3. Results**

171 The selected sites had different anthropic conditions. Cala Zafferano (CZ) is in a military area,
172 downstream of an industrialized coastal area, and subjected to main sea NW currents. Capo
173 Spartivento (CS), located in the Gulf of Cagliari, is protected from the main current NW and
174 exposed to SW and SE, in contrast, Capo Frasca (CF) is located in the south of the Gulf of Oristano
175 but totally exposed to NW current. Both sites are far from industrial and agriculture areas (Figure
176 1). All organisms tested were present in each site and were selected with similar size and weight
177 among the species (Table 1). The analysis of water content and ashes showed similar values in the
178 three sites for gastropods, ranging in average from $74.44 \pm 2.43\%$ (*Osilinus*, CZ), to $79.30 \pm 1.05\%$
179 (*Patella*, CZ). Ash content ranged on average from $3.75 \pm 2.41\%$ (**Thais**, CF), to $8.52 \pm 25.75\%$
180 (*Osilinus*, CZ). However, the higher water levels were detected for *Anemonia*
181 *sulcata*, *Paracentrotus lividus* and *Padina pavonica* accounting in average for $90.85 \pm 1.72\%$, 81.72

Commentato [IC7]: Perché maiuscolo?

Commentato [IC8]: Non deve andare in corsivo?

182 $\pm 2.77\%$, and $82.45 \pm 1.82\%$, respectively, while the lower levels were for *Cystoseira*
 183 *mediterranea* with an average value of $72.52 \pm 7.12\%$ (Table 2).
 184 Seaweeds showed the higher mineral content with ash average values accounting for $15.35 \pm$
 185 17.98% (g/100g $\pm$ RSD, FW) and $12.86 \pm 19.98\%$ (g/100g $\pm$ RSD, FW) in *Padina* CZ
 186 and *Cystoseira* CS, respectively, almost three times higher than the analyzed invertebrates. Instead,
 187 the lowest ash levels were detected in *Anemonia sulcata* CZ and *Paracentrotus lividus* CZ with
 188 average values of $3.50 \pm 7.89\%$ and $2.10 \pm 12.60\%$ (g/100g $\pm$ RSD, FW), respectively.
 189 Gastropods soft tissue, echinoderms gonads, and coelenterate anemone tentacles were used for the
 190 analysis, while the juvenile areal parts were selected for seaweeds.
 191 The ICP-OES analytical method allowed the analysis of 23 heavy metals and metalloids.
 192 Calibration curves showed R^2 values ranging from 0.9967 (Pb) to 0.9997 (Co), and the limits of
 193 quantification of the method were suitable to evaluate overall pollution and risk assessment (Table
 194 3) (Wenzl et al., 2016).
 195 Among the heavy metals and metalloids searched within the methods, eight were missing (Be, Co,
 196 Hg, Mo, Sb, Se, Sn, Te), and only 6 (Al, Cu, Fe, Sr, V, Zn) were found in all samples from the three
 197 sites. Gastropods and seaweeds showed in all sites the presence of 12 heavy metals and
 198 metalloids, *Paracentrotus* 10, and *Anemonia* 9 (Table 4-6).
 199 *3.1 Gastropods*
 200 *Patella vulgata* showed higher values for seven heavy metals and metalloids (Al, Ba, Fe, Mn, Pb,
 201 Sr, Ti) in the samples from CZ, while CS and CF showed even values. B, Cd, Cr, Cu, V, and Zn had
 202 similar values in all three sites. Fe, Sr, and Al from CZ had the higher mean values accounting for
 203 $34.28 \mu\text{g/g}$, $7.06 \mu\text{g/g}$, and $6.44 \mu\text{g/g}$, respectively, about four times higher than those from CF and
 204 CS (Table 3). *Osilinus turbinatus* had the higher mean values in the samples from CZ for Al, Cr,
 205 Cu, Mn, Fe, Ni, Pb, Sr, Ti, and Zn, among these, the highest concentrations were detected for Fe
 206 $40.22 \mu\text{g/g}$, Sr $19.08 \mu\text{g/g}$, Al $8.77 \mu\text{g/g}$, and Ni $7.93 \mu\text{g/g}$. B, Ba, Cd, and V had similar values in
 207 the three sites. Also, *Thais* showed higher values in the samples of CZ. In particular, Al, Ba, Cu, Fe,

Pb, and V were the most concentrated, with Fe 36.89 µg/g, Cu 29.68 µg/g, Zn 21.91 µg/g, and Ba 10.04 µg/g with the higher values (Table 4).

Comparing the three gastropods in the three sites, Al, B, Cu, Fe, and Ti showed similar values in the samples from CZ for *Patella* and *Osilinus*, Mn and Sr for *Patella* and *Thais*, while Cr had similar values in all sites. In CZ, *Thais* showed higher mean values of Ba, Cd, Cu, and Zn, and lower mean values for Al, B, and V.

In CS, B, Fe, and Ti had overlapping values in *Patella* and *Osilinus*, while Mn and Sr were similar in *Osilinus* and *Thais*. Al, Ba, Cd, Pb, and Zn had different values in all sites.

CF showed more considerable variability among all metal's accumulations in the three species. Considering all sites for the single species, the samples of CZ showed higher levels of all metals, except for Zn, which showed overlapping values in the three sites for the three species.

3.2 Seaweeds

Heavy metal and metalloids distribution followed an entirely different and uneven distribution among the sites and between the two seaweed species (Table 5). *Padina* and *Cystoseira* showed both the higher values in the samples of CZ, Fe 225.46 µg/g, Al and Zn both at 104.4 µg/g, and Sr 77.46 µg/g in *Padina*, and Fe 184.03 µg/g, Al 41.85 µg/g, Sr 55.46 µg/g, Zn 41.84 µg/g, Ba 32.08 µg/g, B 12.60 µg/g, and Mn 6.58 µg/g in *Cystoseira*.

Al, Fe, and Zn in *Padina* showed higher concentration in CZ compared to CS and CF, and to *Cystoseira* samples in all sites. *Cystoseira* levels of Al, B, Ba, Cu, Fe, Mn, Sr and Zn were higher in CZ samples, while Cr, Pb, Ti and V were higher in CF.

3.3 Echinoderms and coelenterate

Parancetrotus lividus accumulated only a few metals at moderate concentrations, Al, Sr, and Zn were highly concentrate in CZ. At the same time, the other trace elements showed values in an even range in the three sites. The higher levels were detected for Zn 9.24 mg/g, Al and Sr 5.64 µg/g, and Fe 4.25 µg/g in CZ. The levels of Fe were similar in CZ and CF, and lower in CS (Table 6).

233 *Anemonia sulcata* showed the presence in the tentacles of some minerals at low levels, among the
234 heavy metals detected, Fe 4.07 µg/g, and Al 2.21 mg/g were the most abundant in CZ. Also, Sr and
235 Zn showed values slightly higher in CZ than in CS and CF (Table 5).
236 Accumulation results from all samples showed an overall content of metals higher in Cala
237 Zafferano, especially for Fe, Al, and Zn in seaweeds. These findings confirmed an increase in the
238 environmental pollution in this area which suffered the anthropic impact of the industrial area due
239 to the exposition to NW current (Figure 1).
240 Among the heavy metals and metalloids considered most dangerous, Hg was never detected at a
241 value above the LOQ of the method (0.05 µg/g), which was ten times below the MRL set for fish
242 (CR-EC No 1881/2006). Arsenic was below the LOQ in all samples except in seaweed, and
243 *Cystoseira* showed higher values in all sites. Pb was not present in *Paracentrotus* and *Anemonia*,
244 and the samples of *Osilinus* from CS and CF. In the other samples, its values were all below the
245 MRL (0.3 µg/g), except in the samples of CZ accounting for 0.50 µg/g in *Patella* and 0.46 µg/g
246 in *Thais*. Cd was not detected in *Anemonia*, while showed values below the MRL (0.05 mg/g)
247 in *Paracentrotus* and *Padina* from CS and CF. All other samples showed values above the MRL,
248 with *Patella* showing a value four times higher of the MRL, and *Thais* 20 times higher. No
249 differences were detected among each species between CS and CF sites.
250 **3.4 Multivariate analysis**
251 Multivariate analysis statistics could help in better discriminate among the different species, due to
252 the high number of samples and variables from the analysis. In this paper, we have used the
253 principal component analysis (PCA) approach to describe if the various species studied had
254 different heavy metals accumulation capacity and which were the most discerning heavy metals.
255 PCA biplot of score and loadings for CZ and CS showed a separate group of datasets for each
256 species, allowing to identify 12 clustering groups and to discriminate metals along PC1 and PC2
257 axis (Figure 3).

Commentato [IC9]: Tabella 6

Commentato [IC10]: Ho aggiunto metalloids visto che poi parli di mercurio, ecc...

PCA emphasize the differences in accumulation of the seaweed *Cystoseira* and *Padina* in CZ samples to the other species studied in the same area and to all the other samples collected in CS. Among the gastropod, *Thais* was the most discriminating species. CZ and CS samples were separated from the other gastropod's samples, invertebrate and seaweed, moreover, the samples from CZ were split from CS along the PCA1 axis stressing a more influence of the residues of the most discriminating metals for this axis. The samples of *Patella*, *Osilinus* and *Paracentrotus* fall in a restricted zone of the PCA plot, despite this they were well separated between CZ and CS along the PC1 axis. The metals Pb, Ba, Fe, Mn, Al, Cr, and B were the most discriminating along the PC1 axis, while Cu, Zn, Mo, and Cd were the most discriminating in PC2 axis (Figure 3). Risk assessment showed total THQ for each species below 1, indicating no adverse effects from the consumption of the selected organisms. In all sites, *Thais* showed higher values, while *Anemonia* showed lower values in all sites (Table 7). Moreover, total THQ for the site obtained from the sum of total THQ of the various species were below 1, confirming that no adverse effects were reached also consuming the different species in the same day. The values of total THQ for the site confirmed that CZ had higher harmful metals concentration than CS and CF, and the amounts consumed should be taken into consideration (Table 7).

4. Discussion

Biomonitoring studies have the scope to assess the environmental occurrence of pollutants potentially toxic for human and other living organisms. Different approaches can be used, such as bioconcentration, that studies the direct partition from the environment to target organisms, and biomagnification that is strictly related to the diet leading to increase xenobiotic concentration in the feeder than in the diet. In contrast, bioaccumulation describes the accumulation and enrichment of contaminants in a selected living organism. This process is species-dependent since different organisms have different feeding habits, mechanisms of detoxification and metabolism (Jasminska et al. 2011). Metals and metalloids are commonly present in the environment and can be essential

Commentato [IC11]: Ripetizione. Lo hai già messo a inizio frase.

283 elements supporting biological activities or have adverse effects decreasing aquatic species or being
284 dangerous for human being through the food chain.

285 Gastropods, echinoderm, and anemone species are usually eaten in Sardinia. THQ showed that the
286 intake of single metals from the selected areas through the diet did not create an alarm for the
287 population if food quantities remain in the local typical FDI (food daily intake).

288 Metals do not degrade and along the trophic pyramid tend to increase, and anthropic activity or
289 geochemical conditions can influence their concentration, the natural levels can grow significantly,
290 becoming harmful to human safety. Literature data reported xenobiotic accumulation on suitable
291 species used as indicators. Although most studies have addressed local issues, their impact is
292 important to assess the different bioaccumulation trends of the organisms used.

293 Perez-Lopez et al. (2003) studied the contamination of *Patella vulgata* from Zn, Cd, Pb and Cu with
294 levels ranging in the soft tissue from 91 to 429 µg/g for Zn, 5.6 µg/g for Cd, 1-3 µg/g for Pb, and 2-
295 10 µg/g for Cu.

296 Kelepertzis (2013), studying the heavy metal concentration in *Patella*, postulated that many
297 mineralized areas give rise to higher levels of Pb (96 vs 8 µg/g), Zn (196 vs 48 µg/g), Mn (36 vs 10
298 µg/g), and Cu (28 vs 6 µg/g) bioconcentration. In contrast, Ni (9 µg/g), and Cr (8 µg/g) were even,
299 the levels of metals decreased in the line Zn > Pb > Mn > Cu.

300 Duyssak and Azdural (2017) studied the accumulation from 8 sites in the different tissues
301 of *Patella* of Fe, Zn, Cd, Cu, Co, Ni, Mn, Pb, Cr and Al, with the higher concentration in all sites
302 for Fe, Zn, Cd, Co, Pb, and Al.

303 Collado et al. (2006) found that mean total concentrations of Cd, Cu, Pb, and Zn in *P. rustica* were
304 0.37 ± 0.05 , 1.77 ± 0.09 , 1.27 ± 0.07 and 8.84 ± 0.71 µg/g DW (mean $\pm$ S.E.) respectively; whereas
305 in *P. candei crenata* were 0.71 ± 0.10 , 2.94 ± 0.11 , 0.09 ± 0.01 and 33.74 ± 1.15 µg/g DW (mean $\pm$
306 S.E.).

Commentato [IC12]: Through?

Gadaw (2018) studied the accumulation in the gastropod *Lanistes carinatus* of Mn, Fe, Co, Ni, Cu, Zn, Cd, and Pb. With values ranging from 240 to 567 µg/g dw for Fe, 5.12 – 32.8 µg/g dw for Mn, Zn 5.35-28.15 µg/g dw, 0.69-3.25 µg/g dw for Cu, 4.42-13.87 µg/g dw for Ni, 0-9.24 µg/g dw for Co, 0.03-0.04 µg/g dw for Cd, and 0.24-0.40 µg/g dw for Pb.

Jakimska et al. (2011) reported the levels of Zn and Fe in gastropods. In these molluscs these two metals have essential biochemical functions, Zn is a constituent of haemocyanin, Fe is a constituent of goethite (α -FeOOH) responsible for the proper functioning of the radula. This fact can explain the high levels of Zn and Fe in gastropods. Moreover, the levels of Zn are related to Cu levels through the metabolic processes and homeostasis, metallothionein mediated.

Amin et al. (2006) studied the accumulation of Cu, Zn, Fe, Pb, Cd in the soft tissues of *Thais aculeata* with values of 150.3 ± 58.0 µg/g DW, 158.4 ± 27.9 µg/g DW, 316.7 ± 125.3 µg/g DW, 11.3 ± 5.9 µg/g DW and 0.6 ± 0.3 µg/g DW, respectively.

Ramirez et al. (2013), studied the levels of Cd (0.69-15.81 µg/g), Cu (10.82-55.68 µg/g), Pb (0.21-1.36 µg/g), and Zn (15.78-32.09 µg/g) in *Osilinus*. Later Boucetta et al. (2016) reported the determination in *Osilinus* of Cd, Cr, Cu, Ni, Pb, Zn in two stations, recording different values in the two stations, accounting for 318.10 ± 38.20 µg/g and 6.51 ± 1.47 µg/g for Zn and Pb in the first station. And 3.63 ± 1.14 µg/g for Cd, 32.70 ± 2.90 µg/g for Cu, and 24.80 ± 20 µg/g for Ni in the station 2.

Salvo et al. (2014) studied the accumulation of Hg, As, Cr, Ni, Cu, V, Cd and Pb in *Paracentrotus lividus*, their findings suggested the use of *Paracentrotus* as a bioindicator for pollution in defined areas. Similar findings were presented by Angioni et al. (2012, 2014) analyzing polycyclic aromatic hydrocarbon (PAH) pollution in different Sardinian coastal areas.

Horovits et al. (2014) analyzed symbiotic algae and the tentacles of *Anemonia* for the bioaccumulation of 12 trace elements. *Anemonia viridis* regulates its internal trace element concentrations by compartmentalization and excretion owing a high resilience to environmental contamination from high levels of metals. Pb, Fe, and Cu were mostly associated with symbiotic

algae, while Cd, As, and Zn were associated with the *Anemonia*. 85% of Fe was in the alga fraction, Pb and Cu were evenly distributed, while Zn, As, and Cd were mostly distributed in the *Anemonia*. Mitchelmore et al. (2003) studying the accumulation of heavy metals in anemone pointed out that symbiotic anemones accumulate Cd, Ni e Zn higher than aposymbiotic anemones, and the levels were depleted when recovered in seawater.

Data on the contamination of *Cystoseria* have been reported by Ryabushko et al. (2017) on Pb (35-50 µg/g), Cd (2-3.5 µg/g), Zn (40-65 µg/g), and Cu (13-24 µg/g), and by Benfares et al. (2015) on Cd, Pb (0-5.1 µg/g), Cr (0.1-1.9 µg/g), and Hg.

The same specimen with different size can give rise to different levels of xenobiotics; therefore, to avoid individual variation, we selected specimens from the same species with similar size in each site and among sites.

Literature most studied metals were Cd, Cr, Cu, Pb, and Zn, by using one or two organisms, mainly comparing polluted vs clean areas. The merged data from Sardinia obtained in the present study showed values for all living organisms in the lower range compare to literature data for Cd, Cr and Cu, while Pb and Zinc levels in gastropods were higher than the levels reported in the Canary Islands (Table 8).

Mercury detected in mussels collected along the coastline of Sardinia from two sampling campaigns showed values ranging from 0.035 to 0.11 µg/g and 0.048 to 0.83 µg/g (Ipolyi et al. 2004). Moreover, data reported on carnivorous and herbivorous or omnivorous fish showed that the seconds accumulate till 10 times less the concentration of the former (Levitan et al., 1974; Liu et al., 2014). Therefore, values of Hg below the LOQ in the samples of the present study were not surprising since most of the living organisms studied are mostly herbivorous or seaweed (Table 1).

The results from the studies examined showed that local pollution cannot be used for global discussions but take pictures of a scenario in restricted areas. The Mediterranean basin, which is an enclosed sea, has different contamination characteristics related to the anthropic effects on the coasts and on the sea of the single surrounding countries.

Commentato [IC13]: Forse bisogna mettere Hg?

359 Therefore, the real value of these studies is related to the global information of the accumulation
360 trends of the different living organisms.

361 The present study elucidated that the species of gastropods, *Paracentrotus*, *Anemonia* and seaweed
362 living in the same water column (upper littoral zone) had different uptake capability on metals.

363 These data confirm that accumulation is species and site-dependent, being related to pollution
364 magnitude, feeding habitude and geochemical morphology of the site.

365 MVA analysis emphasizes the different accumulation capacity of the seven species collected at the
366 different upper littoral areas of Sardinia, allowing to distinguish areas with different pollution
367 contaminations in a similar contest.

368

369 **5. Conclusions**

370 The data reported showed different capabilities of the species involved in accumulating different
371 metals. Gastropods are more capable of accumulating Al, Ba, Cu, Fe, Sr, and Zn, but in different
372 amount. *Thais* levels of Cu and Zn are much higher than *Patella* and *Osilinus*, seaweeds can
373 concentrate Al, Fe, and Zn (*Padina* more than *Cystoseira*) in large extent in respect to all other
374 species. At the same time, *Paracentrotus* and *Anemonia* have limited ability to concentrate on
375 specific metals but have a much more homogeneous distribution. THQ showed that *Thais*
376 contributed for almost 50% to total THQ of the site and that among the heavy metals Cd was the
377 most impacting on the overall health concerns for invertebrate eating.

378 All these species have been referred to be model organisms for biomonitoring environmental
379 quality and pollution.

380 Therefore, for a correct environmental assessment of a given area, a diversified approach is strongly
381 recommended by exploiting the different properties of both accumulation and concentration of the
382 metals by different aquatic species.

383

384 **References**

385 Abisil, M., Bemtssen, M., Gerringa, L. 1996. The influence of sediment, food and organic ligands
 386 on the uptake of copper by sediment-dwelling bivalves, *Aquatic Toxicology* 134, 13-29.
 387 Amin, N.M., Noor, J.M., Shazili, A.M. 2006. Analysis of Heavy Metals in Soft Tissues of *Thais*
 388 *Aculeata*, a Gastropod Taken from Chendering Beach, Terengganu as an Attempt to Search for
 389 Indicator of Heavy Metal Pollution in the Aquatic Environment. Proceedings of the 1st International
 390 Conference on Natural Resources Engineering & Technology, Putrajaya, Malaysia, 54-59.
 391 Angioni, A., Porcu, L., Secci, M., Addis, P. 2012. QuEChERS Method for the Determination of
 392 PAH Compounds in Sardinia Sea Urchin (*Paracentrotus lividus*) Roe, Using Gas Chromatography
 393 ITMS-MS Analysis. *Food Anal. Methods*, 5, 1131-1136. DOI 10.1007/s12161-011-9353-7
 394 Angioni, A., Cau, A., Secci, A., Addis, P. 2014. GC–ITMS analysis of PAH contamination levels in
 395 the marine sea urchin *Paracentrotus lividus* in Sardinia. *Marine Pollution Bulletin* 82, 201-207.
 396 Atherton, H.J.; Bailey, N.J.; Zhang, W.; Taylor, J. Major, H.; Shockcor, J. Clarke, K.; Griffin, J.L.
 397 2006. A combined ¹H NMR spectroscopy and mass spectrometry-based metabolomic study of the
 398 PPAR- α null mutant mouse defines profound systemic changes in metabolism linked to the
 399 metabolic syndrome. *Physiological Genomics*. 27, 178-186.
 400 doi:10.1152/physiolgenomics.00060.2006
 401 Aydin-Onen, S., Ozturk, M., 2017. Investigation of heavy metal pollution in eastern Aegean Sea
 402 coastal waters by using *Cystoseira barbata*, *Patella caerulea*, and *Liza aurata* as biological
 403 indicators. *Environmental Science and Pollution Research* 24, 8, 7310-7334. DOI: 10.1007/s11356-
 404 016-8226-4
 405 Benfares, R., Seridi, H., Belkacem, Y., Inal, A. 2015. Heavy Metal Bioaccumulation in Brown
 406 Algae *Cystoseira compressa* in Algerian Coasts, Mediterranean Sea. *Environ. Process.* 2, 429–439.
 407 DOI 10.1007/s40710-015-0075-5
 408 Blackmore, G. 2000. Field Evidence of Metal Transfer from Invertebrate Prey to an Intertidal
 409 Predator, *Thais clavigera* (Gastropoda: Muricidae). *Estuarine, Coastal and Shelf Science*. 51, 2,
 410 127-139.

411 Boucetta, S., Beldi, H., Draredja, B. 2016. Seasonal Variation of Heavy Metals in *Phorcus*
 412 (*Osilinus*) *turbinatus* (Gastropod, Trochidae) in the Eastern Algerian Coast. *Global Veterinaria* 17,
 413 1, 25-41. DOI: 10.5829/idosi.gv.2016.17.01.104129

414 Bryan, G.W., Langston, W.J. 1992. Bioavailability, accumulation and effects of heavy metals in
 415 sediments with special reference to United Kingdom estuaries, a review, *Environ Poll* 76, 89-102.

416 Carral, E., Puente, X., Villares, R., Carballeira, A. 1995. Background heavy metal levels in
 417 estuarine sediments and organisms in Galicia (northwest Spain) as determined by modal analysis.
 418 *Sci. Total. Environ.* 172, 175–188.

419 Chiarelli, R., Martino, C., Roccheri, M.C. 2019. Cadmium stress effects indicating marine pollution
 420 in different species of sea urchin employed as environmental bioindicators. *Cell Stress and*
 421 *Chaperones* 24, 675–687. <https://doi.org/10.1007/s12192-019-01010-1>

422 Collado, C., Ramirez, R., Bergasa, O., Hernandez-Brito, J.J., Gelado-Caballero, M.D., Haroun, R.J.
 423 2006. Heavy metals (Cd, Cu, Pb, and Zn) in two species of limpets (*Patella rustica*, *Patella candei*
 424 *crenata*) in the canary island. *WIT Transactions on Ecology and the Environment* DOI:
 425 10.2495/WP060051

426 COMMISSION REGULATION (EC) No 1881/2006 of December 19 2006, setting maximum
 427 levels for certain contaminants in foodstuff. *Official Journal of the European Union*. 20.12.2006, L
 428 364/5.

429 Cossa, D. 1989. A review of the use of *Mytilus* spp as quantitative indicator of cadmium and
 430 mercury) contamination in coastal waters, *Oceanologica Acta* 12, 417-432.

431 Duysak, Ö., Azdural, K. 2017. Evaluation of Heavy Metal and Aluminium Accumulation in a
 432 Gastropod, *Patella caerulea* L., 1758 in Iskenderun Bay, Turkey. *Pakistan J. Zool.*, 49, 2, 629-637.
 433 DOI: <http://dx.doi.org/10.17582/journal.pjz/2017.49.2.629.637>

434 Farombi, E.O., Adelowo, O.A., Ajimoko, Y.R. 2007. Biomarkers of oxidative stress and heavy
 435 metal levels as indicators of environmental pollution in African catfish (*Clarias gariepinus*) from
 436 Nigeria Ogun River. *Int. J. Environ. Res. Public Health*, 4, 158-165.

437 Fergusson, J. 1990. The heavy elements: chemistry, environmental impact and health effects,
 438 Pergamon Press, Oxford.

439 Gawad, S.S.A. 2018. Concentrations of heavy metals in water, sediment and mollusc gastropod,
 440 *Lanistes carinatus* from Lake Manzala, Egypt. *Egyptian Journal of Aquatic Research* 44, 77–82.
 441 <https://doi.org/10.1016/j.ejar.2018.05.001>

442 Goldberg, E., Bowen, V., Hodge, Y., Flegal, A., Martin, J. 1983. U.S. mussel watch: 1977-1978.
 443 Results of trace metals and radionuclides, *Estuar. Coast Shelf Sci.* 16, 69-93.

444 Horwitz, R., Borell, E.M., Fine, M., Shaked, Y. 2014. Trace element profiles of the sea anemone
 445 *Anemonia viridis* living nearby a natural CO₂ vent. *PeerJ* 2:e538. DOI 10.7717/peerj.538

446 Hummel, H., Moderman, R., Amirad-Triquet, C., Rainglet, F., Duijn, Y., Herssevoost, M., de Jong,
 447 J., Bogaards, R., Bachelet, G., Desprez, M., Marehand, J., Sylvand, B., Amirad, J.C., Rybarczyk,
 448 H., de Wolf, L. 1997. A comparative study on the relation between copper and condition in marine
 449 bivalves and the relation with copper in the sediment, *Aquatic Toxicology* 38, 165 – 181.

450 Ingston, W.J. 1982. The distribution of mercury in British estuarine sediments and its availability to
 451 deposit feeding bivalves, *J Mar Biol Assoc UK* 62, 667-674.

452 Jaishankar, M., Tseten, T., Anbalagan, N., Mathew, B. B., Beeregowda, K.N. 2014. Toxicity,
 453 mechanism and health effects of some heavy metals. *Interdiscip Toxicol.* 7(2), 60–72. doi:
 454 10.2478/intox-2014-0009

455 Jakimska, A., Konieczka, P., Skóra, K., Namieśnik, J. 2011. Bioaccumulation of Metals in Tissues
 456 of Marine Animals, Part I: The Role and Impact of Heavy Metals on Organisms. *Pol. J. Environ.*
 457 *Stud.* 20, 5, 1117-1125.

458 Jitar O., Teodosiu, C., Oros, A., Plavan, G., Nicoara, M. 2015. Bioaccumulation of heavy metals in
 459 marine organisms from the Romanian sector of the Black Sea. *New Biotechnology.* 32, 3, 369-378.

460 Kamaruzzaman, B.Y., Rina, John, B.A., Jalal, K.C.A. 2011. Heavy Metal Accumulation in
 461 Commercially Important Fishes of South West Malaysian Coast. *Research Journal of*
 462 *Environmental Sciences* 5, 6, 595-602.

463 Kelepertzis, E. 2013. Heavy Metals baseline concentrations in soft tissues of *Patella* sp. From the
 464 Stratoni coastal environment, NE Greece. *Ecol Chem Eng S.* 20, 1, 141-149.
 465 Mitchelmore, C.L., Verde, E.A., Ringwood, A.H., Weis, V.M. 2003. Differential accumulation of
 466 heavy metals in the sea anemone *Anthopleura elegantissima* as a function of symbiotic state.
 467 *Aquatic Toxicology* 64, 317-329.
 468 Perez-Lopez, M., Alonso, J., Novoa-Valinas, M. C., Melgar, M. J. 2003. Assessment of Heavy
 469 Metal Contamination of Seawater and Marine Limpet, *Patella vulgata* L., from Northwest Spain.
 470 *Journal of Environmental science and Health, Part A—Toxic/Hazardous Substances &*
 471 *Environmental Engineering* 38, 12, 2845–2856.
 472 Philips, D. 1977. The use of biological indicator organisms to monitor trace metal pollution in
 473 marine and estuarine environments - a review, *Environ Pollut* 13, 281-317.
 474 Ramirez, R. 2013. The gastropod *Osilinus atrata* as a bioindicator of Cd, Cu, Pb and Zn
 475 contamination in the coastal waters of the Canary Islands. *Chemistry and Ecology.* 29, 3, 208-220.
 476 <https://doi.org/10.1080/02757540.2012.735659>
 477 Rodríguez-Figueroa, G. M., Shumilin, E., Sánchez-Rodríguez, I. 2008. Heavy metal pollution
 478 monitoring using the brown seaweed *Padina durvillaei* in the coastal zone of the Santa Rosalía
 479 mining region, Baja California Peninsula, Mexico. *J Appl Phycol.* 21:19–26. DOI 10.1007/s10811-
 480 008-9346-0
 481 Roesijadi, G., Robinson, W.E. 1994. Metal regulation in aquatic animals: mechanisms of uptake,
 482 accumulation, and release. In *Aquatic Toxicology: Molecular, Biochemical, and Cellular*
 483 *Perspectives*; Malins, D.C., Ostrander, G.K., Eds.; Lewis Publishers: Boca Raton, 387–420.
 484 Ryabushko, V. I., Prazukin, A. V., Gureeva, E. V., Bobko, N. I., Kovrigina, N. G., Nekhoroshev,
 485 M. V. 2017. Морской биологический журнал (*Marine Biological Journal*). 2, 2, 70–79. doi:
 486 10.21072/mbj.2017.02.2.07
 487 Salvo, A., Giorgi, A.G., Cicero, N., Bruno, M., Lo Turco, V., Di Bella, G., Dugo, G. 2014.
 488 Statistical characterization of heavy metal contents in *Paracentrotus lividus* from Mediterranean

489 Sea. Natural Product research formerly Natural Products letters. 1-9. DOI:
490 10.1080/14786419.2013.878937

491 Vinogradov, A., Efron, J., Setlow, J. 1953. Elementary Composition of Coelenterata. In ODUM V.
492 (Ed.), Memoir II: The Elementary Chemical Composition of Marine Organisms (pp. 194-219).
493 NEW HAVEN: Yale University Press. Retrieved March 25, 2020, from
494 www.jstor.org/stable/j.ctvbc0gk.11

495 Wenzl, T., Haedrich, J., Schaechtele, A., Robouch, P., Stroka, J. 2016. Guidance Document on the
496 Estimation of LOD and LOQ for Measurements in the Field of Contaminants in Feed and Food.
497 EUR 28099, Publications Office of the European Union, Luxembourg, ISBN 978-92-79-61768-3;
498 doi:10.2787/8931

499 , Y.J., Yang, Z., Zhang, S. 2011. Ecological risk assessment of heavy metals in sediment and human
500 health risk assessment of heavy metals in fishes in the middle and lower reaches of the Yangtze
501 River basin. Environmental Pollution, Volume 159, Issue 10, October 2011, Pages 2575-2585

502 Yüzereroğlu, T.A., Firidin, G.G., Coğun, H.Y., First, O., Aslanyavrsu, S., Maruldali, O., Kargin, F.
503 2009. Heavy metals in *Patella caerulea* (Mollusca, Gastropoda) in polluted and non-polluted areas
504 from the Iskenderun Gulf (Mediterranean Turkey). Environmental Monitoring and Assessment 167,
505 1-4, 257-64. DOI: 10.1007/s10661-009-1047-x

506

507

508

509

510

511

512

513

514

515

516

517 **Table 1.** Humidity and ash content of the edible part of gastropods, Paracentrotus,
518 anemonia, and of the seaweed sampled.
519

	Humidity			Ash		
	g/100g ± RSD FW			g/100g ± RSD FW		
	C. Zafferano (CZ)	C. Spartivento (CS)	C. Frasca (CF)	C. Zafferano (CZ)	C. Spartivento (CS)	C. Frasca (CF)
<i>Patella vulgata</i>	79.46 a(a)	78.91 a(a)	78.01 a(a)	4.61 a(a)	4.29 a(a)	5.04 ab(a)
<i>Osilinus turbinatus</i>	74.44 a(a)	77.54 a(a)	75.62 a(a)	8.91 b(a)	5.10 a(b)	6.59 a(ab)
<i>Thais haemastoma</i>	76.53 a(a)	74.90 a(a)	76.20 a(a)	5.18 a(a)	3.75 ab(b)	4.24 b(ab)
<i>Paracentrotus lividus</i>	84.06 b(a)	79.54 a(a)	81.56 b(a)	2.10 c(a)	2.75 b(a)	3.30 b(a)
<i>Anemonia sulcata</i>	89.06 b(a)	91.44 b(a)	91.96 c(a)	3.50 c(a)	3.85 ab(a)	3.71 b(a)
<i>Padina pavonica</i>	81.67 b(a)	81.50 c(a)	84.18 b(a)	15.35 d(a)	13.13 c(a)	6.30 a(b)
<i>Cystoseira mediterranea</i>	70.21 c(a)	68.91 d(a)	78.44 b(b)	10.85 e(a)	12.86 c(a)	10.48 c(a)

520 * Values among species (without brackets) or among sites (in brackets) followed by unlike letters differ significantly by Fisher's least
521 significant difference (LSD) procedure, P ≤ 0.05. Letters without brackets relate to comparisons of the influence of species within each site.
522 Letters in brackets relate to comparisons of the influence of the sites among each species for each parameter.
523

524

525

526

527

Table 2. Heavy metals method parameters.

Element	λ	MRL µg/g	LOQ	Linear regression equation	R ²
Al	237.3		0.10	y=670.89x-45.48	0.9978
AS	188.98		0.10	y=79.38x-10.14	0.9991
B	249.77		0.50	y=5151.33x+789.8	0.9995
Ba	493.40		0.05	y=8035x-289.9	0.9993
Be	313.04		0.005	y=259128x-6848	0.9985
Cd	226.50	0.05	0.005	y=4804x-3.55	0.9995
Co	228.61		0.025	y=6622.7x+58.6	0.9997
Cr	267.71		0.01	y=9908.7x+97.7	0.9994
Cu	324.75		0.01	y=17117.2x-95.93	0.9996
Fe	259.94		0.10	y=3059.9x+80.8	0.9992
Hg	194.16	0.5	0.05	y=392.2x+11	0.9994
Mn	257.61		0.01	y=70780.6x-8029	0.9990
Mo	204.59		0.05	y=604.4x-5.5	0.9988
Ni	216.55		0.05	y=1642.2x-22.3	0.9986
Pb	220.35	0.3	0.05	y=297.3x-19.4	0.9967
Sb	217.58		0.10	y=49.7x-5.8	0.9981
Se	196.02		0.025	y=157.7x-11.5	0.9992
Sn	189.92		0.05	y=74.3x-4.9	0.9986
Sr	407.77		0.10	y=829945.3x+17943	0.9984
Te	214.28		0.10	y=106.6x-10.1	0.9972

528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562

Ti	336.12		0.01	y=14275x+521.8	0.9976
V	292.40		0.005	y=13753.2x+231.2	0.9996
Zn	213.85		0.05	y=4785.9x-316	0.9986

Table 3. Heavy metals (µg/g ± RSD%) in gastropods from the three sites investigated.

metal	<i>Patella vulgata</i>			<i>Osilinus turbinatus</i>			<i>Thais</i>		
	C. Zafferano (CZ)	C. Spartivento (CS)	C. Frasca (CF)	C. Zafferano (CZ)	C. Spartivento (CS)	C. Frasca (CF)	C. Zafferano (CZ)	C. Spartivento (CS)	C. Frasca (CF)
Al	6.44a(a)	1.41b(b)	1.35b(b)	8.77 a(a)	2.94 b(c)	2.80 b(c)	0.37B a(d)	0.18 b(e)	0.15 b(e)
AS	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ
B	1.63a(a)	1.44a(a)	1.37a(a)	1.54 a(a)	1.35 a(a)	1.40 a(a)	1.14 a(b)	0.90 a(b)	0.92 a(b)
Ba	1.21a(a)	0.52b(b)	0.48b(b)	1.98 a(c)	1.63 a(c)	1.57 a(c)	10.04 a(d)	7.07 b(e)	6.58 b(e)
Be	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ
Cd	0.19a(a)	0.23a(a)	0.20a(a)	0.08 a(b)	0.10 a(b)	0.09 a(b)	0.90 a(c)	1.14 a(c)	1.01 a(c)
Co	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ
Cr	0.03a(a)	0.03a(a)	0.03a(a)	0.11 a(a)	0.03 b(a)	0.03 b(a)	0.04 a(a)	0.03 a(a)	0.03 a(a)
Cu	0.28a(a)	0.26a(a)	0.27a(a)	2.29 a(a)	1.64 b(a)	1.58 b(a)	29.68 a(a)	8.71 b(a)	7.58 b(a)
Fe	34.28a(a)	7.37b(b)	6.58b(b)	40.22 a(a)	15.36 b(c)	14.91 b(c)	36.89 a(a)	10.73 b(d)	11.51 b(d)
Hg	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ
Mn	0.33a(a)	0.07b(b)	0.05b(b)	0.69 a(c)	0.23 b(a)	0.21 b(a)	0.30 a(a)	0.36 a(a)	0.32 a(a)

Mo	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ
Ni	< LOQ	0,13a(a)	0,11a(a)	7,93 a(b)	0,26 b(c)	0,25 b(c)	< LOQ	0,08 a(a)	0,05 a(a)
Pb	0,50a(a)	0,07b(b)	0,08b(b)	0,18 a(c)	< LOQ	< LOQ	0,46 a(a)	0,16 b(c)	0,14 b(c)
Sb	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ
Se	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ
Sn	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ
Sr	7,06a(a)	1,57b(b)	1,65b(b)	19,08 a(c)	8,04 b(a)	7,58 b(a)	6,01 a(a)	6,10 a(a)	6,05 a(a)
Te	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ
Ti	0,13a(a)	0,07b(b)	0,06b(b)	0,18 a(a)	0,09 b(b)	0,10 b(b)	< LOQ	< LOQ	< LOQ
V	0,28a(a)	0,30a(a)	0,31a(a)	0,60 a(b)	0,59 a(b)	0,54 a(b)	0,06 a(c)	0,51 b(b)	0,48 b(b)
Zn	1,69a(a)	1,88a(a)	1,75a(a)	0,62 a(b)	2,16 b(c)	1,98 b(c)	21,91 a(d)	20,31 a(d)	21,03 a(d)

^a Values among species (without brackets) or among sites (in brackets) followed by unlike letters differ significantly by Fisher's least significant difference (LSD) procedure, $P \leq 0.05$. Letters without brackets relate to comparisons of the influence of species within each site. Letters in brackets relate to comparisons of the influence of the sites among each metal.

Table 4. Heavy metals seaweeds.

Element	<i>Padina pavonica</i>			<i>Cystoseira mediterranea</i>		
	C. Zafferano (CZ)	C. Spartivento (CS)	C. Frasca (CF)	C. Zafferano (CZ)	C. Spartivento (CS)	C. Frasca (CF)
Al	104,40a(a)	9,13b(b)	7,30b(b)	41,85a(c)	3,85b(d)	15,09c(e)
AS	0,73a(a)	0,25b(b)	0,26b(b)	1,06a(c)	2,56a(c)	1,05a(c)
B	< LOQ	0,74a(a)	< LOQ	12,60a(b)	5,10b(c)	< LOQ
Ba	9,96a(a)	3,52b(b)	0,68c(c)	32,08a(d)	17,77b(e)	4,46c(b)
Be	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ
Cd	0,07a(a)	0,01b(b)	0,02b(b)	0,08a(a)	0,02b(b)	0,08a(a)
Co	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ
Cr	0,18a(a)	0,02b(b)	0,06b(b)	0,15a(a)	0,03b(b)	0,24c(ac)
Cu	0,42a(a)	0,10b(b)	0,15b(b)	0,79a(c)	0,18b(b)	0,15b(b)
Fe	225,46a(a)	11,41b(b)	15,79b(b)	184,03a(a)	3,94b(c)	3,56b(c)

Hg	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ
Mn	2,88a(a)	0,84b(b)	< LOQ	6,58a(c)	1,40b(a)	0,73c(b)
Mo	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ
Ni	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ
Pb	0,71a(a)	0,11b(b)	0,25c(c)	0,58a(a)	0,13b(b)	0,92c(d)
Sb	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ
Se	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ
Sn	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ
Sr	77,46a(a)	5,39b(a)	5,65b (b)	55,46a(a)	1,53b(c)	1,15b(c)
Te	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ
Ti	2,60a(a)	0,20b(b)	0,92c(c)	1,18a(c)	0,03b(d)	4,41c(e)
V	0,72a(a)	0,21b(b)	0,25b(b)	1,10a(a)	0,52b(c)	1,50c(d)
Zn	104,40a(a)	0,23b(b)	11,42c(c)	41,85a(d)	1,50b(e)	2,97c(f)

^a Values among species (without brackets) or among sites (in brackets) followed by unlike letters differ significantly by Fisher's least significant difference (LSD) procedure, $P \leq 0.05$. Letters without brackets relate to comparisons of the influence of species within each site. Letters in brackets relate to comparisons of the influence of the sites among each metal.

Table 5. Heavy metals in echinoderms, and coelenterate.

Element	<i>Paracentrotus lividus</i>			<i>Anemonia sulcata</i>		
	C. Zafferano (CZ)	C. Spartivento (CS)	C. Frasca (CF)	C. Zafferano (CZ)	C. Spartivento (CS)	C. Frasca (CF)
Al	5.64a(a)	0.27a(a)	1.74a(a)	2.21a(a)	1.95a(a)	2.05a(a)
AS	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ
B	1.25a(a)	0.53a(a)	1.27a(a)	0.63a(a)	0.59a(a)	0.60a(a)
Ba	0.98a(a)	0.25a(a)	0.40a(a)	0.13a(a)	0.12a(a)	0.10a(a)
Be	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ
Cd	0.06a(a)	0.04a(a)	0.03a(a)	< LOQ	< LOQ	< LOQ
Co	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ
Cr	0.02a(a)	0.04a(a)	0.02a(a)	< LOQ	< LOQ	< LOQ
Cu	0.27a(a)	0.14a(a)	0.17a(a)	0.20a(a)	0.12a(a)	0.16a(a)
Fe	4.25a(a)	1.85a(a)	4.38a(a)	4.07a(a)	3.81a(a)	2.95a(a)
Hg	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ
Mn	< LOQ	< LOQ	< LOQ	0.22a(a)	0.35a(a)	0.27a(a)
Mo	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ
Ni	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ
Pb	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ
Sb	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ
Se	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ
Sn	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ
Sr	5.64a(a)	0.79a(a)	2.89a(a)	0.64a(a)	0.29a(a)	0.31a(a)
Te	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ	< LOQ
Ti	0.09a(a)	0.02a(a)	0.08a(a)	< LOQ	< LOQ	< LOQ
V	0.12a(a)	0.08a(a)	0.10a(a)	0.02a(a)	0.03a(a)	0.02a(a)
Zn	9.24a(a)	4.87a(a)	2.47a(a)	0.62a(a)	0.22a(a)	0.13a(a)

* Values among species (without brackets) or among sites (in brackets) followed by unlike letters differ significantly by Fisher's least significant difference (LSD) procedure, $P \leq 0.05$. Letters without brackets relate to comparisons of the influence of species within each site. Letters in brackets relate to comparisons of the influence of the sites among each metal.



680

681 **Figure 1.** Localization of the three sites in Sardinia, main current  NW, and  industrial area.

682




Figure 2. Histograms of the distribution of the metals among the species in the three locations.




Figure 3. PCA biplot of the score and loading in CZ and CS sites.