

# Università degli Studi Mediterranea di Reggio Calabria

Archivio Istituzionale dei prodotti della ricerca

Benzofuran-2-acetic esters as a new class of natural-like herbicides

This is the peer reviewd version of the followng article:

Original

Benzofuran-2-acetic esters as a new class of natural-like herbicides / Araniti, F; Mancuso, R; Lupini, A; Sunseri, F; Abenavoli, Mr; Gabriele, B. - In: PEST MANAGEMENT SCIENCE. - ISSN 1526-498X. - 76:(2020), pp. 395-404. [10.1002/ps.5528]

Availability: This version is available at: https://hdl.handle.net/20.500.12318/3139 since: 2024-11-22T15:43:06Z

Published DOI: http://doi.org/10.1002/ps.5528 The final published version is available online at:https://onlinelibrary.wiley.com/doi/10.1002/ps.5528

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website

Publisher copyright

This item was downloaded from IRIS Università Mediterranea di Reggio Calabria (https://iris.unirc.it/) When citing, please refer to the published version.

(Article begins on next page)

This is the peer reviewed version of the following article: Araniti et al (2020) Benzofuran-2-acetic esters as a new class of natural-like herbicides. Pest Management Science. 10.1002/ps.5528.

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website. Received: 3 May 2019

Revised: 17 June 2019

(wileyonlinelibrary.com) DOI 10.1002/ps.5528

# Benzofuran-2-acetic esters as a new class of natural-like herbicides

# Fabrizio Araniti,<sup>a</sup> Raffaella Mancuso,<sup>b</sup> Antonio Lupini,<sup>a</sup> Francesco Sunseri,<sup>a</sup> Maria Rosa Abenavoli<sup>a\*</sup> and Bartolo Gabriele<sup>b\*</sup>®

## Abstract

BACKGROUND: In recent decades, the use of synthetic herbicides has been increasing, mainly in emerging countries. However, their intensive and indiscriminate application is a major cause of environmental pollution and human health injury. Therefore, there is an increasing need to develop new herbicides with safer toxicological and environmental profiles. A promising strategy is to synthesize new molecules containing the core of natural products as a template for the production of 'bio-inspired' or 'natural-like' herbicides.

RESULTS: The potential herbicidal activity of some benzofuran-2-acetic esters was assessed *in vitro* on *Arabidopsis thaliana*, a model species. All five molecules (M1-M5) showed significant phytotoxic activity, reducing both shoot and root system at low concentrations. In particular, methyl 2-(5-methoxybenzofuran-2-yl)hexanoate (M3) exhibited the highest phytotoxicity displayed against two crops and weeds, monocots (*Zea mays* L. and *E. crus-galli*) and dicots (*Lactuca sativa* L. and *Amaranthus retroflexus* L.). The M3 activity was also compared with glyphosate, a common herbicide, showing a lower but similar activity. Moreover, the results evidenced that M3 was more effective in post-emergency.

CONCLUSION: Readily synthesizable benzofuran-2-acetic esters possessing the benzofuran ring as 'bio-inspired' core, show significant herbicidal activity making them very efficient even at low concentrations. They can be sprayed in liquid form, and the addition of adjuvants can improve penetration through the leaf cuticle. These results confirm the importance of these molecules as models for the development of new natural-like herbicides. © 2019 Society of Chemical Industry

Keywords: benzofurans; crop protection; herbicides; natural-like herbicides

## **1 INTRODUCTION**

Weeds have the largest negative impact on crop productivity<sup>1,2</sup> since they compete for soil, water and nutrients, causing significant economic losses.<sup>3,4</sup> Cultural, mechanical, chemical and biological methods are the most prominent approaches used for weed management. Among these, synthetic herbicides, whose use is increasing in recent decades, mainly in emerging countries, remain to date the most effective method for their ease of application and greater accessibility for farmers.<sup>5</sup> However, the intensive and indiscriminate application of these chemicals, very persistent in agricultural end-products and not easily biodegradable, is a major cause of environmental pollution and human health injury. In addition, their poor selectivity along with the rapid evolution of weed resistance determine huge damage to crops and negative consequences in agriculture.<sup>6</sup> Therefore, there is an increasing search to develop new herbicides with safer toxicological and environmental profiles and new mechanisms of action (MOA). A very promising strategy is to synthesize new molecules containing the core of natural products as a template for the production of 'bio-inspired' or 'natural-like' herbicides. They could be characterized by high biological activity, novel target sites and alternative MOA, low or no toxicity, high biodegradability compared to the commercial herbicides, and consequently an increased environmental and health safety.<sup>7</sup> Until now, natural products have not played an important role in herbicide development compared to

other pesticides,<sup>8,9</sup> although some natural-like compounds with herbicidal activity have been developed.<sup>10–17</sup>

In this paper, benzofuran-2-acetic esters are presented as a new class of 'bio-inspired' potential herbicides.<sup>19</sup> The benzofuran motif constitutes the core of many pharmacologically active substances, with biological activities that include anti-inflammatory, anticancer, cytotoxic, antimicrobial, antioxidant, insecticidal and antiplasmodial effects.<sup>20–25</sup> However, the herbicidal activity of benzofuran-2-acetic esters has not been reported so far.<sup>19</sup> In this work, the phytotoxic activity of these compounds, readily synthesizable by a catalytic carbonylation process, was evaluated on the model plant *Arabidopsis*, crops and weeds, and was also compared to glyphosate, a common commercial herbicide.

- \* Correspondence to: MR Abenavoli, Dipartimento AGRARIA, Università Mediterranea di Reggio Calabria, 89124 Reggio Calabria, Italy. E-mail: mrabenavoli@unirc.it; B Gabriele, Dipartimento di Chimica e Tecnologie Chimiche, Università della Calabria, via P. Bucci 12/C, 87036 Arcavacata di Rende, CS, Italy. E-mail: bartolo.gabriele@unical.it
- a Department of AGRARIA, University "Mediterranea" of Reggio Calabria, Reggio Calabria, Italy
- b Laboratory of Industrial and Synthetic Organic Chemistry (LISOC), Department of Chemistry and Chemical Technologies, Arcavacata di Rende, Cosenza, Italy



**Figure 2.** Dose – response curve of shoot fresh weight (SFW) of *A. thaliana* seedlings treated with benzofurans **M1-M5** for 14 days. Data are expressed as percentage compared to the control. Different letters along the curves indicate significant differences at  $P \le 0.05$  (Tukey's test, n = 4).

## 2 MATERIALS AND METHODS

#### 2.1 Synthesis of benzofuran-2-acetic esters

Benzofuran-2-acetic esters M1-M5 were prepared by palladiumcatalyzed carbonylation of (2-allyloxyphenyl)-2-yn-1-ols, as we already reported.<sup>26</sup> Briefly, a 250 mL stainless steel autoclave was charged with Pdl<sub>2</sub> (7.2 mg, 2.0·10<sup>-2</sup> mmol), KI (332.0 mg, 2.0 mmol), PPh<sub>3</sub> (21.0 mg,  $8.0 \cdot 10^{-2}$  mmol) and a solution of the substrate (2.0 mmol) in anhydrous MeOH (9.1 mL). Water (72 µL, 4.0 mmol) was then added, and the autoclave was sealed, purged at room temperature several times with CO with stirring (10 atm) and eventually pressurized to 30 atm. After stirring at 100 °C for 15 h, the autoclave was cooled and degassed. The solvent was eliminated by rotary evaporation, and products were purified by column chromatography on silica gel using 1:1 hexane-CH<sub>2</sub>Cl<sub>2</sub> (M1), 8:2 hexane-acetone (M2), 8:2 hexane-AcOEt (M3, M5), and 95:5 hexane-AcOEt (M4). The yields obtained are shown in Figure 1. Characterization data for all products can be found in the above-mentioned paper.26

#### 2.2 In vitro bioassays on Arabidopsis thaliana

The *in-vitro* bioassays were carried out as previously reported by Araniti *et al.* with some modifications.<sup>27</sup> *Arabidopsis thaliana* (L.) Heynh. seeds ecotype Columbia (Col-0) were surface sterilized in 50% EtOH (3 min), 0.5% NaOCl with 0.01% Triton X-100 (3 min), and then rinsed three times in distilled water.

Seeds were maintained in 0.1% agar at 4 °C for 72 h to promote the synchronization of germination. Then, 25 seeds were sown in Petri dishes  $(100 \times 150 \text{ mm})$  containing agar medium (0,8%)w/v) enriched with micro- and macronutrients (Murashige-Skoog, Sigma-Aldrich) and sucrose (1%). Plates were placed vertically in the growth chamber (22  $\pm$  2 °C temperature, 55% HR, and 90 mol $\cdot$ m<sup>-2</sup> $\cdot$ s<sup>-1</sup> light intensity) to promote geotropic root growth. Four days after sowing, six seedlings (7 days old) were transferred to a single plate and grown in the same medium containing 0, 50, 100, 200, 400, 800 µM of synthetic benzofuran derivatives (M1-M5), for 14 days (see section 2.1 for benzofuran derivatives preparation). All the molecules were previously dissolved in 0.1% 







**Figure 3.** Pigment content [Chlorophyll a (Chla), chlorophyll b (Chlb) and carotenoids (Ct)] in young *Arabidopsis* plants treated with **M1-M5** for 14 days. Data are expressed as percentage compared to the control. Different letters along the bars indicate significant differences at  $P \le 0.05$  (Tukey's test, n = 4).

EtOH (v/v), and then autoclaved. The same amount of EtOH was added to the control plates. After treatment, seedlings were collected and separated into shoot and root. Shoot Fresh Weight (SFW), Leaf Number (LN), Total (TRL) and Lateral Root Length (LRL) were evaluated. The root system was imaged by scanning (STD 1600, Régent Instruments Inc., Quebec, QC, Canada) and analyzed by WinRhizo Pro System v. 2002a (Instruments Règent Inc., Quebec, Canada). Root hair density (RHD) was evaluated with an Olympus microscope (SZX9).

#### 2.3 Pigments content

6 Chlorophyll a (Chla), chlorophyll b (Chlb), and carotenoid (Ct) 7 content was quantified on *Arabidopsis* seedlings treated with each 8 molecule (**M1-M5**) for 14 days. One hundred milligrams of leaf 9 tissue, for each replicate and treatment, were ground in liquid 9 nitrogen and extracted with 1.5 mL methanol. The extract was 9 then centrifuged at 170 g for 5 min, and afterwards, 500  $\mu$ L of 9 supernatant was mixed with 500  $\mu$ L methanol. The absorbance of the extracts was determined at 470, 653, 666 and 750 nm. The pigment content (mg  $g^{-1}$  of SFW) was evaluated according to the following equations proposed by Wellburn<sup>28</sup>:

Chla (
$$\mu$$
g) = [15.65 (DO<sub>666</sub> - DO<sub>750</sub>) - 7.34 (DO<sub>653</sub> - DO<sub>750</sub>)] × V

Chlb (
$$\mu$$
g) = [27.05 (DO<sub>653</sub>-DO<sub>750</sub>)-11.21 (DO<sub>666</sub>-DO<sub>750</sub>)] × V

Carotenoids (X + C) (
$$\mu$$
g) = [1000 (DO<sub>470</sub>-DO<sub>750</sub>)  
-2.86 Chla-129.2 Chlb)/221] × V

where DO is the optical density, V is the volume of methanol used (mL) and (X + C) the sum of carotenoids and xanthophylls.

#### 2.4 Benzofuran M3 vs glyphosate

The potential herbicidal activity of the most bioactive benzofuran **M3** was compared to glyphosate, a common commercial herbicide. The preparation of the commercial herbicide (Roundup<sup>®</sup>)

85

86

87

88

89

90

91

92

93

94

95

96

97

98

99

100

101

102

103

104

105

106

107 108

109

115

116

117

118

119 120 121

122 123

124





**Figure 4.** Arabidopsis thaliana seedlings grown in agarized medium and treated with benzofuran **M1** for 15 days. (A) Control; (B) 50  $\mu$ M; (C) 100  $\mu$ M; (D) 200  $\mu$ M.

was carried out following package instructions to simulate an open field application. **M3** was solubilized in 96% ethanol (1:5 w/w), and then diluted with deionized water to reach 0 (control), 3.6 and 7.2 mM concentrations. Further, the **M3** solubility was increased by adding in the solution few drops of 0.001 mM NaOH, as emulsifying agent together with a surfactant and a wetting product [Etravon Pro (1 L/200 L/ha), Syngenta, Italy] that increased its penetration and efficacy during foliar treatment. Foliar-applied distribution was carried out by spraying, calculating an equal amount of active principle to be diluted in 200–300 L, the amount of solution distributed in a hectare.

# 2.5 In vivo bioassays on crops and weeds and pigment content

The phytotoxic activity of benzofuran **M3** and glyphosate were assayed *in vivo* on two crops and their respective weeds: *Zea mays* L. (monocot) and *Echinochloa crus-galli* (monocot), *Lactuca sativa* L. (dicot) and *Amaranthus retroflexus* (dicot). These species

are commonly used in phytotoxic experiments, but, above all, 63 this choice allowed us to evaluate the potential selectivity of 64 M3 against monocot and dicot.<sup>29,30</sup> Seeds of each species were 65 surface sterilized in 15% NaClO solution (v/v) for 15 min and 66 then rinsed three times with deionized water. Then, seeds (a 67 seed/hole) were placed in a tray (0.2 m<sup>2</sup>) with hole (5  $\times$  5 cm) 68 corresponding to a total volume of about 250 cm<sup>3</sup>, containing 69 potting soil. After sowing, trays were placed in a growth chamber 70 (26 °C, 65% humidity, 12 h photoperiod with a light intensity of 71 350 mol  $m^{-2} s^{-1}$ ) and watered with Hoagland nutrient solution 72 (Hoagland and Arnon, 1950) every 2 days. Twenty-one days after 73 sowing, young plantlets of each species were sprayed with a 74 solution containing 0 (control), 3.6 and 7.2 mM M3 or glyphosate 75 concentrations. After 7 days of treatment, shoots were collected 76 for measuring fresh weight (g), stem length (cm) and pigment 77 content (%). Stem length was measured by the WinRHIZO system 78 pro STD 1600 (Instruments Régent Inc. Canada) after scanner 79 acquisition (Epson Expression 800). Shoot dry weight (g) was 80 determined after keeping seedlings at 72° C until constant weight 81 was reached. The determination of pigments content was carried 82 out as seen in Section 2.3. 83

#### 2.6 Statistical analysis

A completely random design with four replications was adopted to evaluate the effects of the benzofuran derivatives in the in-vitro Arabidopsis bioassay. Data were evaluated for normality (Kolmogorov-Smirnov test) and tested for the homogeneity of variances (Levene's test). Shoot fresh weight (SFW), TRL, LRL of seedlings in response to different doses of benzofuran derivatives were evaluated by a nonlinear regression model using a log-logistic function that allowed to estimate the ED<sub>50</sub> parameter, the dose required to reduce 50% of measured parameter compared to control. The phytotoxicity comparison among molecules was performed by one-way ANOVA using the ED<sub>50</sub> as a variable and the molecule as main factor. The ED<sub>50</sub> data were first checked for deviations from normality (Kolmogorov-Smirnov test) and tested for homogeneity (Levene's test). The statistical significance of differences among means was estimated by analysis of variance and Tukey test ( $P \le 0.05$ ). The statistical significance of pigments



www.soci.org



indicate statistical differences with  $P \le 0.05$ . N = 4.

content was estimated by analysis of variance (one-way ANOVA) and the data was processed as previously described.

A completely random design with 10 replications was adopted to evaluate the effects of **M3** and glyphosate in the *in vivo*-bioassays with crops and weeds. All the data were evaluated for normality (Kolmogorov–Smirnov test) and tested for homogeneity of variances (Levene's test). The statistical significance of differences among means was estimated by analysis of variance (two-way ANOVA) where the experimental factors were molecule/herbicide and concentration followed by Tukey's test ( $P \le 0.05$ ). All the analyses were performed by using SPSS ver. 6.1 software (Insightful Corporation, Seattle, WA, USA).

#### **3 RESULTS**

#### 3.1 In vitro bioassays on Arabidopsis thaliana

All the benzofuran derivatives **M1-M5** caused a strong inhibitory effect on *Arabidopsis* seedlings. Except for **M5**, all molecules caused a significant SFW reduction already at the lowest concentration (50 μM), with inhibition values ranging from 44% (**M2**) to 83% (**M3**). As the concentration increased (≥ 100 μM), all molecules showed a SFW inhibition higher than 76% (Fig. 2). On the contrary, **M5** induced a significant SFW reduction only at concentrations higher than  $100 \,\mu$ M, with inhibition values ranging from 70% ( $200 \,\mu$ M) to 97% ( $800 \,\mu$ M) (Fig. 2). These effects were sometimes accompanied by a failure of leaf development, as shown for **M1** and **M3** (Figs. 4 and 5, respectively), whose treated seedlings failed to develop the first true leaves. In **M1** treated plants, the cotyledons appeared to be depigmented (Figs. 3 and 4(d)).

All molecules significantly affected pigment content at a different extent (Fig. 3). Among them, **M2**, **M3** and **M4** were the most effective, causing a high reduction of all the pigments already at the lowest concentration. In particular, they induced a chlorophyll a reduction about 32% at the lowest concentration (50  $\mu$ M), reaching inhibition values higher than 87% at 800  $\mu$ M (Fig. 3). A similar trend was observed in chlorophyll b content, which was reduced by 20% at the lowest concentration, reaching an inhibition value higher than 75% at the highest concentration (800  $\mu$ M) (Fig. 3). Finally, the carotenoid content was the most affected parameter in seedlings treated with 50  $\mu$ M, pointing out inhibition values ranging from 38% (**M3**) to 57% (**M4**). At the highest concentration (800  $\mu$ M), an inhibition higher than 79% was observed (Fig. 3).

Moreover, root system showed a significant reduction in TRL, NLR and root hair density (data not shown). To compare the potential phytotoxicity of the benzofuran derivatives on root system, the  $ED_{50}$ , which defines the dose required to reduce 50% of the

<sup>12</sup>4 տ





M3

Glyphosate



**Figure 7.** Dry weight (g) of *Z. mays, L. sativa, E. cruss-galli* and *A. retroflexus* exposed to 0, 3.6 or 7.2 mM **M3** or glyphosate for 7 days. Different letters within species indicate significant differences at *P* < 0.05 (Tukey's test, *n* = 10).

M3

Glyphosate



**Figure 9.** Chlorophyll a content ( $\mu$ g g<sup>-1</sup> shoot fresh) of *Z. mays, L. sativa, E. cruss-galli* and *A. retroflexus* exposed to 0, 3.6 or 7.2 mM **M3** or glyphosate for 7 days. Different letters within species indicate significant differences at *P* < 0.05 (Tukey's test, *n* = 10).

60

61

62

1

1





**Figure 10.** Chlorophyll b content ( $\mu$ g g<sup>-1</sup> shoot fresh) of *Z. mays, L. sativa, E. cruss-galli* and *A. retroflexus* exposed to 0, 3.6 or 7.2 mM **M3** or glyphosate for 7 days. Different letters within species indicate significant differences at *P* < 0.05 (Tukey's test, *n* = 10).

total response, was calculated. The results showed that all these molecules exhibited a high phytotoxicity at <466  $\mu$ M concentrations. In particular, TRL was strongly reduced at concentrations ranged from 12 to 466  $\mu$ M, while lower concentrations, ranging from 10 to 145  $\mu$ M, were able to cause the SFW and NRL reduction (Table 1). Among the benzofuran derivatives, **M3** appeared the most effective, showing ED<sub>50</sub> values for SFW, TRL and NLR at <20  $\mu$ M concentration (Fig. 5 and Table 1). By contrast, **M5** resulted the less effective, while the other molecules showed different effects based on the parameter taken into account (Table 1).

3.2 In vivo bioassays on crops and weeds: M3 vs glyphosate

Due to its higher phytotoxicity, M3 was chosen for the in vivo assays on two crops (maize and lettuce, monocot and dicot, respectively) and two weeds (E. crus-galli and A. retroflexus, monocot and dicot, respectively). The M3 effectiveness was compared with that of glyphosate based on biomass production (shoot fresh and dry weights, SFW and SDW) and photosynthetic pigments (chlorophyll a and b, carotenoids content). All showed a significant reduction after treatment with both the molecules in both crops and weeds. In particular, a significant SFW reduction (52%) was observed in maize at the highest M3 concentration (7.2 mM), even though lower compared to that caused by commercial herbicide (85%). This inhibitory effect on fresh weight was also evident, at lower extent, in the other species, resulting in a reduction of 20, 24, and 30% compared to the control in A. retroflexus, L. sativa, and E. cruss-galli, respectively. By comparison, the glyphosate was able to reduce significantly SFW also at the lowest concentration (3.6 mM) (Fig. 6). Shoot dry weight (g) was less affected by M3 at the lowest concentration, while glyphosate confirmed greater efficacy being able to reduce SDW already at 3.6 µM in all the species except for A.

*retroflexus*, which did not appear to be affected by both molecules (Fig. 7). At the highest concentration, **M3** was able to reduce SDW in maize and *E. crus-galli* (41 and 21% compared to the control, respectively) but not lettuce (Fig. 7).

After 7 days of treatment, a significant decrease in stem length was observed (Fig. 8). In particular, **M3** caused a reduction of 45, 28, 22, and 21% in maize, lettuce, *E. crus-galli* and *A. retroflexus*, respectively, at the highest concentration with no significant effects observed at the lowest one (Fig. 8). By contrast, the commercial herbicide showed a high phytotoxicity on maize and *E. crus-galli* at the lowest concentration (3.2 mM), causing plant death in maize at the highest concentration (Fig. 8). Finally, in all the treated plants, both **M3** and glyphosate treatment significantly reduced all the pigments content (chlorophyll a and b, carotenoids) at all the concentrations (Figs. 9–11).

## **DISCUSSION**

Benzofuran derivatives constitute a major group of biologically active heterocyclic compounds among natural and synthetic products.<sup>31</sup> The benzofuran core is present in many biologically active compounds, characterized by antifungal, insecticidal, antioxidant, antiviral and anticancer activities.<sup>32,33</sup> For this reason, benzofuran has also been introduced as the core to develop new herbicides as reported in recent literature.<sup>19,34,35</sup> Recently, the investigation of methods for developing novel benzofuran derivatives with herbicidal activity has become an important goal to develop new natural-inspired herbicides. 

In this paper, we have assessed the potential herbicidal activity of some synthetic benzofuran-2-acetic esters **M1-M5**, synthesized according to Figure 1. All five benzofuran derivatives showed 



**Figure 11.** Carotenoid content ( $\mu$ g g<sup>-1</sup> shoot fresh) of *Z. mays, L. sativa, E. cruss-galli* and *A. retroflexus* exposed to 0, 3.6 or 7.2 mM **M3** or glyphosate for 7 days. Different letters within species indicate significant differences at *P* < 0.05 (Tukey's test, *n* = 10).

a high phytotoxic potential, inhibiting biomass production (SFW) and reducing shoot pigment content in Arabidopsis. These effects suggested that these compounds could limit photosynthesis causing subsequent plant growth inhibition. Interestingly, pigments have often been used as biomarkers for new molecules with herbicidal activity in crops and model plants.<sup>36,37</sup> Fai et al. demonstrated that some herbicides caused chlorophyll inhibition blocking its synthesis.<sup>38</sup> However, this may not be the only cause because pigment content reduction could be determined by a chemically-induced degradation or an oxidative stress induced by benzofuran derivatives. The effects induced by benzofurans were dose- and molecule-dependent for all the pigments. Carotenoids appeared to be more sensitive than both Chla and Chlb. As known, carotenoids play an important role to overcome oxidative stress,<sup>39</sup> quenching reactive oxygen species (ROS) and stabilizing photosynthetic complexes.<sup>40–42</sup> Furthermore, carotenoids act as 'energy reducing agents' for capturing excess energy during photosynthesis, which in turn destroys membranes of cells and organelles such as chloroplasts. Therefore, carotenoid reduction and ROS production could be the cause of chlorophyll biosynthesis inhibition and, consequently, of chlorotic and less developed leaves as observed in Arabidopsis seedlings treated with all benzofurans M1-M5. Similar effects on plant biomass and pigment content were previously observed with other compound classes.<sup>17,18</sup>

To understand the different effectiveness of these chemicals, the  $ED_{50}$  of SFW, TRL and NLR were compared. These values confirmed the strong inhibitory activity of all the molecules underlying also their significant different efficacy. Among them, **M3** was the most effective, showing the lowest  $ED_{50}$  values for each parameter taken into account. It is well known that the presence of substituents in the aromatic ring may affect phytotoxicity. For example, the hydroxylation of *trans*-cinnamic acid decreased ion uptake ability,<sup>39</sup> and the position and/or the isomerization of the substitution group made *cis*-cinnamic acid 10 times more active than *trans*-cinnamic acid in *Arabidopsis* root growth inhibition.<sup>40</sup> Furthermore, lactonization increased the effect of coumarin making it a stronger inhibitor than *trans*-cinnamic acid in the germination process.<sup>41</sup> Comparing the chemical structure of the five benzofurans, **M3** shows a methoxy substituent at the C-5 position, which might be responsible for its higher phytotoxicity. For this reason, **M3** was chosen to be compared with glyphosate, the most common herbicide used in agriculture on several crop and weed species.

90

91

92

93

95

96

97

98

99

100

101

102

103

104

105

106

107

108

109

110

111

112

113

114

115

116

117

118

119

120

0

The *in vivo* bioassay results indicate that **M3** possesses a moderate/good herbicidal efficacy, highlighting also a certain selectivity against monocots, as reported for other compounds containing the benzofuran moiety.<sup>35</sup> Despite the higher effectiveness of glyphosate, **M3** caused a significant plant growth inhibition mainly at the highest concentration (7.2 mM). Moreover, the results show a strong effect on the photosynthetic pigments as already observed in *Arabidopsis*, suggesting that the **M3** target could be the shoots, and consequently, it was more effective in post-emergency (spray treatment on sprout).

#### 5 CONCLUSIONS

In conclusion, we have found that readily available benzofuran-2-acetic esters possess a high herbicidal activity even at low doses. In particular, all the molecules assayed *in vitro* strongly affected the growth and development of the model species *Arabidopsis thaliana* affecting both shoot and root growth and development. The results obtained suggest that these natural-like molecules represent a promising pharmacophore for the development of new agrochemicals.

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

0

63

64

65

66

67

68

69

70

71

72

For future applications, in order to improve the efficacy of these molecules, benzofuran-2-acetic esters could be formulated using adjuvants to reduce the surface tension between the spray droplet and the leaf surface and to improve their penetration through leaf cuticle.

#### REFERENCES

- Tilman D, Cassman KG, Matson PA, Naylor R and Polasky S, Agricultural sustainability and intensive production practices. *Nature* 418:671–677 (2002).
- 2 Pimentel D, Zuniga R and Morrison D, Update on the environmental and economic costs associated with alien-invasive species in the United States. *Ecol Econ* **52**:273–288 (2005).
- 3 Charudattan R, Biological control of weeds by means of plant pathogens: significance for integrated weed management in modern agro-ecology. *BioControl* 46:229–260 (2001).
- 4 Rajcan I and Swanton CJ, Understanding maize-weed competition: resource competition, light quality and the whole plant. *Field Crops Res* 71:139–150 (2001).
- 5 Monaco TJ, Weller SC and Ashton FM, *Weed Science: Principles and Practices*, 4th edn. John Wiley & Sons, New York (2002).
- 6 Bennett R, Phipps R, Strange A and Grey P, Environmental and human health impacts of growing genetically modified herbicide-tolerant sugar beet: a life-cycle assessment. *Plant Biotechnol J* 2:273–278 (2004).
- 7 Dayan FE and Duke SO, Natural compounds as next-generation herbicides. *Plant Physiol* **166**:1090–1105 (2014).
- 8 Copping LG and Duke SO, Natural products that have been used commercially as crop protection agents. *Pest Manag Sci* **63**:524–554 (2007).
- 9 OF H, Use of natural products in the crop protection industry. *Phy-tochem Rev* 10:185-194 (2011).
- 10 Duke SO, Dayan FE, Rimando AM, Schrader KK, Aliotta G, Oliva A et al., Chemicals from nature for weed management. Weed Sci 50:138–151 (2002).
- 11 Cornes D, Callisto: a very successful maize herbicide inspired by allelochemistry, in Proceedings of Fourth World Congress on Allelopathy, Wagga Wagga, NSW, Australia. (2005) Available: http://www. regional.org.au/au/allelopathy/2005/2/7/2636\_cornesd.htm.
- 12 Mitchell G, Bartlett DW, Fraser TEM, Hawkes TR, Holt DC, Townson JK et al., Mesotrione: a new selective herbicide for use in maize. Pest Manag Sci 57:120–128 (2001).
- 13 Grossmann K and Ehrhardt T, On the mechanism of action and selectivity of the corn herbicide topramezone: a new inhibitor of 4-hydroxyphenylpyruvate dioxygenase. *Pest Manag Sci* 63:429–439 (2007).
- 14 Romagni JG, Allen SN and Dayan FE, Allelopathic effects of volatile cineoles on two weedy plant species. J Chem Ecol 26:303–313 (2000).
- 15 Grayson BT, Williams KS, Freehauf PA, Pease RR, Ziesel WT, Sereno RL et al., The physical and chemical properties of the herbicide cinmethylin (SD 95481). *Pestic Sci* 21:143–153 (1987).
- 16 Barton AF, Dell B and Knight AR, Herbicidal activity of cineole derivatives. J Agric Food Chem 58:10147–10155 (2010).
- 17 Araniti F, Mancuso R, Ziccarelli I, Sunseri F, Abenavoli MR and Gabriele B, 3-(methoxycarbonylmethylene) isobenzofuran-1imines as a new class of potential herbicides. *Molecules* 19:8261–8275 (2014).
- 18 Araniti F, Mancuso R, Lupini A, Giofré SV, Sunseri F, Gabriele B et al., Phytotoxic potential and biological activity of three synthetic coumarin derivatives as new natural-like herbicides. *Molecules* 20:17883–17902 (2015).
- 19 An Italian patent has been filed on July 13, 2017 (# 102015902347682), titled: "Benzofurani ad attività erbicida". Inventors: Maria Rosa ABENAVOLI, Bartolo GABRIELE, Antonio LUPINI, Raffaella MAN-CUSO, Francesco SUNSERI, Fabrizio ARANITI; Applicants: University "Mediterranea" of Reggio Calabria and University of Calabria, Italy. The Italian Patent Office issued the patent on August 31, 2017, with # 0001429724.
- 20 Kao CL and Chern JW, A convenient synthesis of naturally occurring benzofuran ailanthoidol. *Tetrahedron Lett* 42:1111–1113 (2001).
- 21 Spaniol M, Bracher R, Ha HR, Follath F and Krähenbühl S, Toxicity of amiodarone and amiodarone analogues on isolated rat liver mitochondria. *J Hepatol* **35**:628–636 (2001).

- 22 Carlsson B, Singh BN, Temciuc M, Nilsson S, Li YL, Mellin C *et al.*, Synthesis and preliminary characterization of a novel antiarrhythmic compound (KB130015) with an improved toxicity profile compared with amiodarone. *J Med Chem* **45**:623–630 (2002).
- 23 Narimatsu S, Takemi C, Kuramoto S, Tsuzuki D, Hichiya H, Tamagake K et al., Stereoselectivity in the oxidation of bufuralol, a chiral substrate, by human cytochrome P450s. *Chirality* 15:333–339 (2003).
- 24 De Campos MP, Cechinel Filho V, Da Silva RZ, Yunes RA, Zacchino S, Juarez S et al., Evaluation of antifungal activity of Piper solmsianum C. DC. var. solmsianum (Piperaceae). Biol Pharm Bull 28:1527–1530 (2005).
- 25 Aslam SN, Stevenson PC, Phythian SJ, Veitch NC and Hall DR, Synthesis of cicerfuran, an antifungal benzofuran, and some related analogues. *Tetrahedron* 62:4214–4226 (2006).
- 26 Gabriele B, Mancuso R, Salerno G and Costa M, Cascade reactions: sequential homobimetallic catalysis leading to benzofurans and β, γ-unsaturated esters. Adv Synth Catal 348:1101–1109 (2006).
- 27 Araniti F, Costas-Gil A, Cabeiras-Freijanes L, Lupini A, Sunseri F, Reigosa MJ et al., Rosmarinic acid induces programmed cell death in Arabidopsis seedlings through reactive oxygen species and mitochondrial dysfunction. PLoS One 13:e0208802 (2018).
- 28 Wellburn AR, The spectral determination of chlorophylls a and b, as well as total carotenoids, using various solvents with spectrophotometers of different resolution. *J Plant Physiol* **144**:307–313 (1994).
- 29 Díaz-Tielas C, Sotelo T, Graña E, Reigosa MJ and Sánchez-Moreiras AM, Phytotoxic potential of trans-chalcone on crop plants and model species. J Plant Growth Regul 33:181–194 (2014).
- 30 Araniti F, Marrelli M, Lupini A, Mercati F, Statti GA and Abenavoli MR, Phytotoxic activity of *Cachrys pungens* Jan, a mediterranean species: separation, identification and quantification of potential allelochemicals. *Acta Physiol Plant* **36**:1071–1083 (2014).
- 31 Khanam H, Bioactive benzofuran derivatives: a review. *Eur J Med Chem* 97:483–504 (2005).
- 32 Naik R, Harmalkar DS, Xu X, Jang K and Lee K, Bioactive benzofuran derivatives: moracins A–Z in medicinal chemistry. *Eur J Med Chem* **90**:379–393 (2015).
- 33 Khodarahmi G, Asadi P, Hassanzadeh F and Khodarahmi E, Benzofuran as a promising scaffold for the synthesis of antimicrobial and antibreast cancer agents: a review. *J Res Med Sci* **20**:1094–1104 (2015).
- 34 Yang ZH, Li W, Yang ZH, Shao DF and Hu AX, Preparation and application of benzofuranol derivatives containing triazole group. *Chin Pat CN* **104**:236 A (2015).
- 35 Yan Z, Yang Z, Deng X, Lin D, Wu M, Li J et al., Novel aryloxyphenoxypropionate derivates containing benzofuran moiety: design, synthesis, herbicidal activity, docking study and theoretical calculation. Pest Biochem Physiol 154:78–87 (2019).
- 36 Yancheva SD, Georgieva LD, Kostova M, Halkoglu P, Dimitrova M and Naimov S, Plant pigments content as a marker for herbicide abiotic stress in corn (*Zea mays L.*). *Emirates J Food Agric* 28:332–336 (2016).
- 37 Sandmann G, Schneider C and Böger P, A new non-radioactive assay of phytoene desaturase to evaluate bleaching herbicides. Z Naturforsch C 51:534–538 (1996).
- 38 Fai PB, Grant A and Reid B, Chlorophyll a fluorescence as a biomarker for rapid toxicity assessment. *Environ Toxicol Chem* 26:1520–1531 (2007).
- 39 Smirnoff N, The role of active oxygen in the response of plants to water deficit and desiccation. *New Phytol* **125**:27–58 (1993).
- 40 Demmig-Adams B, Gilmore AM and Adams WW, Carotenoids 3: in vivo function of carotenoids in higher plants. *FASEB J* **10**:403 – 412 (1996).
- 41 Bassi R and Caffarri S, Lhc proteins and the regulation of photosynthetic light harvesting function by xanthophylls. *Photosynth Res* **64**:243–256 (2000).
- 42 Munné-Bosch S and Penuelas J, Photo-and antioxidative protection during summer leaf senescence in *Pistacia lentiscus* L. grown under mediterranean field conditions. *Ann Bot* **92**:385–391 (2003).
- 43 Yu JQ and Matsui Y, Effects of root exudates of cucumber (*Cucumis sativus*) and allelochemicals on ion uptake by cucumber seedlings. J Chem Ecol 23:817–827 (1997).
- 44 Wong WS, Guo D, Wang XL, Yin ZQ, Xia B and Li N, Study of cis-cinnamic acid in Arabidopsis thaliana. Plant Physiol Biochem 43:929–937 (2005).
- 45 Reynold T, Comparative effects of heterocyclic compounds on inhibition of lettuce fruit germination. *J Exp Bot* **40**:391–404 (1989).

# **Contents**

Published online in Wiley Online Library:

**Research Article** 

There is an increasing need to develop new 'bio-inspired' herbicides. This article shows how readily available benzofuran-2-acetic esters constitute a new class of synthetic compounds with significant herbicidal activity.



Benzofuran-2-acetic esters as a new class of natural-like herbicides

Fabrizio Araniti, Raffaella Mancuso, Antonio Lupini, Francesco Sunseri, Maria Rosa Abenavoli and Bartolo Gabriele

Pest Manag Sci 2019; **0**: 0