

Università degli Studi Mediterranea di Reggio Calabria

Archivio Istituzionale dei prodotti della ricerca

Effects of rice husk biochar on rill detachment capacity in deforested hillslopes

This is the peer reviewd version of the followng article:

Original

Effects of rice husk biochar on rill detachment capacity in deforested hillslopes / Parhizkar, M; Shabanpour, M; Lucas-Borja, Me; Zema, Da. - In: ECOLOGICAL ENGINEERING. - ISSN 0925-8574. -191:106964(2023), pp. 1-7. [10.1016/j.ecoleng.2023.106964]

Availability: This version is available at: https://hdl.handle.net/20.500.12318/141556 since: 2024-11-22T10:56:37Z

Published DOI: http://doi.org/10.1016/j.ecoleng.2023.106964 The final published version is available online at:https://www.sciencedirect.

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website

Publisher copyright

This item was downloaded from IRIS Università Mediterranea di Reggio Calabria (https://iris.unirc.it/) When citing, please refer to the published version.

1	This is the peer reviewed version of the following article:
2	
3	Parhizkar, M., Shabanpour, M., Lucas-Borja, M.E., Zema D.A. 2023. <i>Effects of rice husk</i>
4	biochar on rill detachment capacity in deforested hillslopes. Ecological Engineering (Elsevier),
5	161:106964,
6	
7	which has been published in final doi
8	
9	10.1016/j.ecoleng.2023.106964
10	
11	(https://www.sciencedirect.com/science/article/pii/S0925857423000733)
12	
13	The terms and conditions for the reuse of this version of the manuscript are specified in the
14	publishing policy. For all terms of use and more information see the publisher's website

15	Effects of rice husk biochar on rill detachment capacity in deforested hillslopes		
16			
17	Misagh Parhizkar ^{1,*} , Mahmood Shabanpour ¹ , Manuel Esteban Lucas-Borja ² , Demetrio Antonio		
18	Zema ³		
19			
20	¹ Department of Soil Science, Faculty of Agricultural Sciences, University of Guilan, Rasht, Iran		
21	² Escuela Técnica Superior Ingenieros Agrónomos y Montes, Universidad de Castilla-La Mancha.		
22	Campus Universitario, E-02071, Albacete, Spain.		
23	³ Department AGRARIA, Mediterranean University of Reggio Calabria, Loc. Feo di Vito, I-89122,		
24	Reggio Calabria, Italy		
25			
26	* Correspondence: <u>misagh.parhizkar@gmail.com</u>		
27			
28	Abstract		
29			
30	Biochar addition to soil has been found to be a useful practice to control		
31	erosion. However, few studies have explored the effects of blochar		
32	produced from husk of rice on rill erosion, due to overland flow in		
33	deforested areas. This study has evaluated the rill detachment capacity (D) is the little (U_i) is the li		
34	(D_c) and erodibility (K_r) in soil samples treated with rice husk blochar in		
35	comparison to untreated sites (control), collected in deforested hillslopes		
36	of Northern Iran. D_c was measured in a lab-scale hydraulic flume at		
37	Four-bed slopes (9.7, 14.3, 19.5, and 24.2%) and five water discharges (0.24, 0.26, 0.47, 0.50, 10.60 L $-\frac{-1}{2}$ N		
38	(0.24, 0.36, 0.47, 0.59, and 0.69 L m s). Moreover, organic carbon,		
39	aggregate stability and bulk density of soils were measured. D_c was		
40	lower (on average -32%) in the treated soil compared to the control. D_c		
41	was not correlated to the other physico-chemical properties of soil. K _r ,		
42	estimated by linear regressions between D_c and shear stress, was		
43	These results avidence that rise buck biocher is effective at controlling		
44	and mitigating soil datashment, and at improving arganic matter containts		
45	and initigating son detachment, and at improving organic matter content		
40	rill gradibility and shoar stress of deforested hillslopes (trested or		
4/	untreated) are helpful to model soil resistance to rill program in process		
48	based erosion models		
49			

50

51 **Keywords:** rill erosion; soil conservation; organic matter content; soil aggregate stability; rill 52 erodibility; shear stress.

53

55

56

57

58

59

60

61

62

63

64

65

66

67

68

69

70

71

72

73

74

75

76

77

78

79

80

81

82

83

84

54 1. Introduction

Soil detachment, which is the initial stage of erosion, is the removal of particles from the soil matrix due to raindrop impact or overland flow (Jiang et al. 2020). In the case of clear water, the maximum value of soil detachment in rills formed by erosion is indicated as "rill detachment capacity" (Nearing et al., 1992). This is a key parameter for estimating the rate of soil removed by overland flow (Govers et al., 2007) and describing the overall erosion process (Nearing et al., 1991). Moreover, this parameter is widely used in process-based erosion models, such as the Water Erosion Prediction Project (WEPP) (Wang et al., 2016).

Rill detachment capacity is widely variable, and this variability is due to many factors, such as the soil and plant root characteristics as well as the land use and management (Parhizkar et al., 2021b). The latter factor is essential to reduce the overall erosion rates, since effective soil conservation practices decrease the sediment source for subsequent soil transport and deposition (Zhang et al., 2009). These practices are important in all environments, and essential in deforested areas with high susceptibility to soil erosion (Parhizkar et al., 2020b). For example, some hillslopes that were deforested in recent years in Northern Iran (e.g., Guilan province) have undergone intolerable erosion rates (Parhizkar et al., 2021d), while the previous plant cover prevented noticeable soil loss.

> Rill detachment, as other erosion forms, can be noticeably reduced by soil amendment, but with various levels of effectiveness (Li et al., 2021). For instance, supplying soil with biochar, which is produced through pyrolysis of carbon-rich residues in the lack of oxygen (Peng et al., 2011), is a feasible conservation practice that has been successfully used in several environments to mitigate erosion. Theoretically, the biochar has a great capability to improve soil structure, as an organic conditioner (Burrell et al., 2016), since it increases soil carbon thanks to its content

in porous organic matter with high stability (Gurwick et al., 2013). Therefore, application of biochar to soil as conservation practice improves soil quality with beneficial effects on its physical, chemical, and microbial properties (Obia et al., 2016; Domene et al., 2014).

85

86

87

88

89

90

91

92

93

94

95

96

97

98

99

100

101

102

103

104

105

106

107

108

109

110

111

112

113

114

115

116

117

118

Rice, together with corn and wheat, is a common staple food, especially in Asia, where much of the world production is concentrated. Therefore, the vegetal residues of rice yield (such as straw and husk) are abundant at low or zero cost in several Asian countries. Various studies have demonstrated that rice husk is a good substrate to produce biochar, and therefore rice residues can effectively improve soil aggregate stability with consequent reductions in runoff and erosion rates (Jien and Wang, 2013; Gamage et al., 2016). Rice husk biochar can improve the hydrological and other physico-chemical properties of soil, since the cellulosic substances of husk can be incorporated and then degraded (Parhizkar et al., 2020b). As such, rice husk biochar can be practically used for soil conservation in those deforested lands that are prone to soil erosion and degradation, due to the removal of plant cover. It is essential, however, to evaluate how much rice hush biochar is effective and thus beneficial at reducing runoff and erosion rates in deforested lands.

The literature about the impacts of biochar on soil erosion is abundant. For instance, Prakongkep et al. (2021) showed that biochar increases the average size of aggregates and wet aggregate stability by 6 to 25% in comparison to untreated soil in farm fields of Thailand. Ghorbani et al. (2019) reported that the application of biochar significantly increases mean weight diameter (MWD), geometric mean diameter (GMD) and water stable aggregates (WSA) of soil compared to the untreated sites. According to Hseu et al. (2014), during a rainfall event simulated at an intensity of 80 mm h⁻¹ on a slope of 20°, erosion significantly decreased by 35% to 90% in soils treated with biochar in comparison to the untreated soil left bare. In Iran, Ahmadi et al. (2020) showed that, in soil with a loamy texture, the application of biochar reduced the runoff coefficient and erosion in comparison to the soil without biochar. These authors also pointed out the need of further studies to obtain a detailed knowledge of the impact of biochar on soil erosion. However, in spite of generally positive experiences, some studies report contrasting effectiveness of biochar at reducing erosion and, more in general, at improving soil quality. For instance, while Nicosia et al. (2021) stated that biochar supplied as soil conditioner increases the storage capacity and prevents soil particle detachment, Zhang et al. (2019) reported that biochar could increase soil erosion, particularly in croplands. Moreover, the investigations focusing the effectiveness of biochar produced from rice husk at reducing the rill erosion are quite scarce, especially on steep slopes in sensitive ecosystems, such as forests or deforested lands (Lucas-Borja et al., 2019a; Parhizkar et al., 2020a; 2021e). Since aggregate stability is a key physical property driving particle detachment, studies investigating its impacts on rill detachment capacity and erodibility in treated vs. untreated soils are needed.

This study: (1) explores the changes in rill detachment capacity of soils treated with rice husk biochar in comparison to untreated sites; (2) analyzes the soil properties that mostly influence rill detachment capacity; and (3) develops regression models to predict rill erodibility from detachment capacity for both treated and untreated soils. The investigation has been carried out using a hydraulic flume that simulated rill detachment at different longitudinal slope and water discharges on soil samples collected in deforested areas of Northern Iran. The results of the study could give land managers insight about the usefulness of treatments with rice husk biochar in deforested lands, where soil conservation is essential, and proposes reliable regression models to predict the magnitude of the rill detachment process, when more precise estimations are not possible or very difficult.

2. Materials and methods

2.1. Study area

A forestland (Saravan Forestland Park) in Guilan province (Northern Iran, 37°08′20″ N; 49°39′42″ E, elevation between 50 and 250 m above

the mean sea level) was chosen as case study (Figure 1). The climate of the area, typically Mediterranean, is of Csa type according to the classification by Kottek et al. (2006). On average, the annual temperature is 16.3 °C and the rainfall 1360 mm (IRIMO, 2016). The soil has a silty clay loam texture (sand 12.9%, silt 47.8% and clay 39.3%) (SDSD, 2017).



Figure 1 - Geographical location of the study area (a); aerial view (b) of deforested hillslopes
(Saravan Forestland Park, Guilan province, Northern Iran) (Source: Google[®] Map[®]).

In recent years, Sarawak Forestland Park was deforested and subjected to unsuitable management practices, such as the vegetation removal on hillslopes to install high-voltage power towers (Parhizkar et al., 2020a; 2021b). This resulted in intense soil erosion, particularly in its rill form (Parhizkar et al., 2020c).

2.2. Soil sampling

172Between July and August 2021, 20 pairs of soil samples were collected173from the uppermost 20 cm in randomly-chosen points of the deforested174hillslopes, according to the procedure by Parhizkar et al. (2021b). Of175each pair of samples, one was used for both rill detachment simulations176in the flume experiments (see section 2.3), and another was subjected to

177physico-chemical analysis (see section 2.6). Prior to soil sampling, litter178and weeds were removed from soil surface. The samples were179transported to the Soil Testing Laboratory of the College of Agriculture180of Guilan University (Iran).181

2.3. Plot preparation

182

183

202

203

204

One of each pair of soil samples was placed in an experimental plot with 184 the same size and characteristics of those described by Parhizkar et al. 185 (2020b; 2021b). Two plots were prepared: a first plot received an 186 untreated soil (hereafter indicated as "control" soil); in the other plot, a 187 soil-biochar mixture at 3% (w/w) of biochar was applied ("treated" soil). 188 Biochar was produced from rice husk by an electrical muffle furnace at a 189 peak temperature of 500° C and a duration of 30-45 min for the pyrolysis 190 process, according to the procedure by Ghorbani et al. (2019). 191 Immediately after production, biochar was applied on the plots with 192 treated soils and left there for three months before the experiments by 193 the hydraulic flume (see section 2.5). 194 Both plots were subjected to wetting and drying cycles in the open air 195 for 24 hours, in order to set the water content of soil at the field capacity. 196 Then, the plots were placed in a room at 25° C throughout a period of 10 197 months. Finally, small soil samples were extracted from the treated and 198 control plots, using a steel ring (0.1 m in diameter and 0.05 m in height) 199 (Khanal and Fox, 2017; Parhizkar et al., 2021b) for the subsequent flume 200 experiments. 201

2.4. Flume experiments

The samples extracted from the two plots were subjected to experiments simulating rill detachment in a slope-adjustable hydraulic flume (length \times width was 3.5 m \times 0.2 m) made of steel. This allowed the measurement of the rill detachment capacity of both treated and control soils at variable flow conditions and longitudinal slopes.

In more detail, each soil sample was inserted in a hole of the flume bed, close to the downstream outlet. Their surface was sprayed with water for 24 h before the experiment, in order to fully saturate the flume bed with water. Then, the flow characteristics were measured according to the procedure setup by Parhizkar et al. (2020a). To summarize, a graduated cylinder was used to measure the water discharge at the downstream outlet of the flume in six replications. The surface flow velocity of the water stream was determined in nine replications using the fluorescent dye technique. The mean velocity was calculated after correction using the coefficients propose by Abrahams et al. (1985). A level probe and a self-made digital instrument with accuracy of 1 mm (Parhizkar et al., 2021c) were used to measure the mean water depth, averaging six measurements in two cross-sections and three points per section (Parhizkar et al., 2020a). The experiment ended when the depth of the eroded soil in the steel ring was approximately 0.015 m or after five minutes from the experiment start. The weight of scoured sediments in each experiment was determined after soil drying in oven for 24 h at 105 °C. Further details about the flume characteristics and the experiments are reported in Parhizkar et al. (2020a; 2021a).

223

224 2.5. Experimental design

225

Twenty combinations of four slopes of the flume bed (9.7%, 14.3%, 19.5%, and 24.2%) and five water discharges (0.24, 0.36, 0.47, 0.59, and 0.69 L m⁻¹ s⁻¹) were setup for each soil sample (Table 1). The values of bed slope were selected according to the longitudinal gradient of the natural hillslopes in the study area, while the water discharges were chosen from field measurements of overland flows (Parhizkar et al., 2020a; 2021b). Each experiment was conducted in five replicates. Therefore, the experimental designed consisted of 200 flume experiments: two soil conditions (treated vs. control) × four bed slopes × five water discharges × five replications.

233

234

2.6. Calculation of rill detachment capacity, erodibility and shear stress

235

The mean rill detachment capacity (D_c , kg s⁻¹ m⁻²), under each combination of bed slope and water discharge, was calculated by equation (1):

238

$$D_c = \frac{\Delta M}{A \cdot \Delta t \cdot \cos \alpha} \tag{1}$$

240

239

where ΔM is the dry weight of detached soil (kg), A is the area of the soil sample (m²), Δt is the experiment duration (s) and α the slope angle of the flume bed (°).

Rill erodibility (K_r , s m⁻¹) and critical shear stress (τ_c , Pa) are meaningful indicators of the soil resistance to rill erosion (Wang et al., 2014), commonly used in hydrological models, such as the WEPP model (Nearing et al., 1989). These parameters were estimated as the slope and intercept of the following equation, which linearly interpolates D_c and shear stress (τ):

$$248 D_C = K_r(\tau - \tau_c) (2)$$

The shear stress, according to Nearing et al. (1997), was estimated using the following equation (3): $\tau = \rho gRS$ (3)

where ρ is the water density (kg m⁻³), g is the acceleration of gravity (m s⁻²), R is the hydraulic radius (m) and S is the bed slope (m m⁻¹).

Table 1 - Values of the hydraulic parameters in the flume experiments for measuring the rill detachment capacity on samples collected in deforested hillslopes of Guilan province (Northern Iran) and treated with rice husk biochar or left undisturbed (control).

Sland	Water	Water	Hydraulic	Shear
Slope	discharge	depth	radius	stress
(8, %)	$(q, L m^{-1} s^{-1})$	(h, cm)	(R, m)	(τ, Pa)
	0.24	0.560	0.005	5.041
	0.36	0.680	0.006	6.053
9.7	0.47	0.790	0.007	6.960
	0.59	0.880	0.008	7.689
	0.69	0.956	0.009	8.295
	0.24	0.466	0.004	6.201
	0.36	0.577	0.005	7.645
14.3	0.47	0.694	0.006	9.095
	0.59	0.781	0.007	10.152
	0.69	0.881	0.008	11.347
	0.24	0.342	0.003	6.319
	0.36	0.427	0.004	7.826
19.5	0.47	0.571	0.005	10.322
17.5	0.59	0.681	0.006	12.184
	0.69	0.772	0.007	13.696
	0.24	0.291	0.003	6.706
	0.36	0.364	0.004	8.329
24.2	0.47	0.485	0.005	10.970
~ r.~	0.59	0.583	0.006	13.065
	0.69	0.676	0.006	15.017

2.7. Soil analysis

Each soil sample extracted for physico-chemical analysis was air-dried and sieved through a 2-mm mesh. The following physico-chemical properties were measured: (1) organic carbon (OC, using the Walkley -Black technique, Allison, 1975); (2) medium weight diameter (MWD) and geometric mean diameter (GMD) of soil aggregates as well as water stable aggregate (Kemper and Rosenau, 1986; Kandeler, 1996), by the wet sieving method; and (3) bulk density (BD), using the clod method (Radcliffe and Simunek, 2010).

2.8. Statistical analysis

A paired sample t-test was used to identify statistically significant differences in rill detachment capacity between control and treated soils at *p*-level < 0.01. QQ-plots were used to check the normality of sample distribution.

A Principal Component Analysis (PCA) was applied to identify possible associations among rill detachment capacity and physico-chemical properties of the treated and control soils.

Finally, a linear regression analysis was carried out to study the relationships between the rill detachment capacity and the shear stress.

The statistical analyses were performed using XLSTAT 9.0 software (Addinsoft, Paris, France).

3. Results

3.1. Variations of rill detachment capacity and soil physico-chemical properties between treated and control soils

The rill detachment capacity (D_c) in the control soil was significantly (p < 0.01) higher compared to the treated plot. Soil treatment with rice husk biochar decreased D_c by 32.5% (0.046 ± 0.024 kg m⁻² s⁻¹ against a value 0.031± 0.014 kg m⁻² s⁻¹ measured for the control soil). The latter soil

condition showed a lower variability in D_c, shown by the lower standard deviation compared to the control (Figure 2).



304

300

301 302 303

Figure 2 - Box plots of rill detachment capacity measured using flume experiments on samples collected in deforested hillslopes of Guilan province (Northern Iran) and treated with rice husk biochar or left undisturbed (control).

308 309

All soil properties were significantly different between the treated and control plots (p < 0.01). In comparison to the control, the treated soil showed an average increase in OC by 21.2%, while MWD, WSA and GMD increased on average by 47.8%, 20.9% and 23.7%, respectively. In contrast, the BD of the treated soil was lower by 10% compared to the control (Table 2).

316

Table 2 - Values of the main soil properties (mean \pm std. dev.) measured in samples (n = 20) collected in deforested hillslopes of Guilan province (Northern Iran) and treated with rice husk biochar or left undisturbed (control).

Soil	Soil condition		
properties	Treated	Control	
OC (%)	2.15 ± 0.28 a	$1.77\pm0.08~b$	
MWD (mm)	0.98 ± 0.15 a	$0.66\pm0.08~\text{b}$	
WSA (%)	45.38 ± 4.44 a	37.55 ± 3.89 b	
GMD (mm)	0.64 ± 0.11 a	0.52 ± 0.06 b	
$BD (kg m^{-3})$	1411 ± 45.27 a	1568 ± 23.83 b	

Notes: OC = organic carbon; MWD = medium weight diameter of soil aggregates; WSA = water stable aggregate; GMD = geometric mean diameter of soil aggregates; BD = bulk density. Different letters in each line indicate significant differences (p < 0.01, Tukey test) between control and treated soils.

The PCA identified two principal components (PC1 and PC2), which explained together 70.4% of the total variance of the original variables (rill detachment capacity and physico-chemical properties of soils). More specifically, PC1 and PC2 depicted 55.19% and 15.17% of this variability, respectively. All the physico-chemical properties of soil influenced the first PC (loadings > 0.483), while the rill detachment capacity heavily weighted in the PC2 (loading of 0.767). In other words, the PC1 increased with OC, BD, MWD, WSA and GMD, while the second PC increased only with D_c (Table 3 and Figure 3a). This means that the analyzed physico-chemical properties did not influence D_c, since the two PCs are uncorrelated.

A clear gradient along the first PC was found between treated and control soils, when the scores of the soil samples were plotted on the first two PCs. Two well-discriminated clusters, one for each soil condition, were evident, which discriminated soils treated with rice husk biochar from untreated soils (Figure 3b).

342

321

322 323

324 325

326

327

328

329

330

331

332

333

334

335

336

337

Table 3 - Loadings of the original variables (rill detachment capacity and soil physico-chemical properties) on the first Principal Components (PC1 and PC2) measured on samples collected in deforested hillslopes of Guilan province (Northern Iran) and treated with rice husk biochar or left undisturbed (control).

Original variables	PC ₁	PC ₂
D_c	0.168	0.767
OC	0.576	0.072
BD	0.738	0.018
MWD	0.658	0.025
WSA	0.690	0.001
GMD	0.483	0.027

Notes: D_c = rill detachment capacity; OC = organic carbon; BD = bulk density; MWD = medium weight diameter of soil aggregates; WSA = water stable aggregate; GMD = Geometric mean diameter of soil aggregates. Significant parameters at p < 0.01 are reported in bold.

0.54



(a)



Legend: $D_C = rill$ detachment capacity; OC = organic carbon; BD = bulk density; MWD = medium weight diameter of soil aggregates; WSA = water stable aggregate; GMD = Geometric mean diameter of soil aggregates.

 Figure 3 - Loadings of rill detachment capacity and soil physicochemical properties (a), and scores (b) of the original variables (rill detachment capacity and soil properties) on the first Principal Components (PC1 and PC2) measured on samples collected in deforested hillslopes of Guilan province (Northern Iran) and treated with rice husk biochar (A) or left undisturbed (control, C).

3.2. Estimation of rill erodibility using linear regression equations

The linear regression models that interpolate D_c and τ by equation (2) showed high coefficients of determination (0.92 and 0.88 for control and treated soil, respectively, p < 0.01) (Figure 4), and this highlights the strong influence of the shear stress on the rill detachment process. The slope (K_r) and intercept (τ_c) were different between the two soil conditions, as expected. In more detail, both K_r and τ_c were lower in the treated soils (0.0047 s m⁻¹ and 2.51 Pa, respectively) compared to the control (0.0081 s m⁻¹ and 3.46 Pa) (Figure 4).



378

382

383

384

385

386

387

388

389

390

391

392

393

394

395

396

Figure 4 – Linear regression equations between the rill detachment capacity ($y = D_c$) and shear stress ($x = \tau$) estimated on soil samples collected in deforested hillslopes of Guilan province (Northern Iran) and treated with rice husk biochar or left undisturbed (control).

4. Discussions

4.1. Influence of biochar on soil properties and rill detachment capacity

The soils of the study area (treated with rice husk biochar and untreated), of similar type and texture, were subjected to rill detachment simulations under the same experimental conditions. Hence, the impacts of the treatment on the rill detachment capacity and erodibility can be easily identified.

The analysis of the main physico-chemical properties of soils carried out in this investigation confirms the higher quality of treated soils as well as more resistance to rill detachment compared to the control plots. In contrast, the latter soils (simulating the deforested and untreated sites) showed higher degradation and erodibility.

The significant increase in organic carbon content of the treated soil compared to the untreated soil is consistent with the findings of other authors (e.g., Masulili et al., 2010; Ghorbani et al., 2019). The increased organic carbon content indicates beneficial conditions for cultivation of the investigated soils (Gamage et al., 2016), since a higher organic matter content (linked to the organic carbon) improves soil fertility and thus enhances crop growth and yield. Biochar plays a significant effect on soil organic carbon, since this substrate contains fresh organic matter (Dume et al., 2016). Moreover, some investigations report that biochar, due to its stable structure, is resistant to surface oxidation of organic carbon, and therefore this soil conditioner is effective at improving the agronomic conditions of the soil (Nguyen et al., 2008).

400

401

402

403

404

405

406

407

408

409

410

411

412

413

414

415

416

417

418

419

420

421

422

423

424

425

426

427

428

429

The significant differences in the indices of soil aggregate stability observed in this study for the treated soil may be due to the organic matter supply, as observed by Zeraatpisheh et al. (2021). These authors showed that soil organic matter has a very high influence on aggregate stability indices. This increase is beneficial for soil aggregation, since the organic compounds act as cement for soil particles. In general, soil disturbance due to anthropogenic factors (e.g., land abandonment, deforestation, cultivation) result in decreases in soil organic matter and thus aggregate stability (Lucas-Borja et al., 2019b; Shabanpour et al., 2020). According to Li et al. (1992), high aggregate stability results in high resistance to rill erosion of soil due to the overland flow.

The lower bulk density measured in the treated soil may be attributed to the increase in the volume of the soil-biochar mixture over time (Ghorbani et al., 2019). In other words, the rearrangement of soil and biochar particles, which is due to the release of applied pressure by soilorganic complex, could have reduced the bulk density in soils treated with biochar (Pandian et al., 2016). This result also agrees with the findings of other authors (e.g., Herath et al., 2013; Jien and Wang, 2013), who reported that the total porosity and macroporosity of soils significantly increase when organic soil conditioners, such as biochar, are added.

430Rill detachment variability was in the same direction as the changes in431soil physico-chemical properties. The decrease in soil erodibility thanks432to treatment with rice hush biochar can be considered as a beneficial433effect against soil degradation in forestland. This decrease is consistent

with the statements of Jien and Wang (2013). These authors report that biochar is able to improve the physical and chemical properties of degraded soil, and reduce erosion, because its application increases macro-aggregates and help in mitigating erosion. In contrast, biochar may not be an appropriate material for all conditions, such it could enhance the loss of humus in forestland (Wardle et al., 2008) and increase the displacement of the biochar particles on the soil surface (Verheijen et al. 2009).

434

435

436

437

438

439

440

441

442

443

444

445

446

447

448

449

450

451

452

453

454

455

456

457

458

459

460

461

462

463

464

465

466

467

As highlighted by the principal component analysis, the two soil conditions (soils treated with rice husk biochar and untreated sites) are clearly different in terms of rill detachment capacity, organic carbon, and aggregate stability. However, the PCA did not associate the variability in rill detachment capacity with organic carbon, aggregate stability, and bulk density. This contrasts the findings of Parhizkar et al. (2020a), who report that an increase in organic carbon and aggregate stability implies a decrease in rill detachment capacity, and the opposite happens for bulk density. That investigation was carried out in undisturbed woodlands and forestlands. In our study, the management operations carried out in recent years in the experimental hillslopes may have altered the soil surface, which may explains the inconsistency between the results of this study and those by Parhizkar et al. (2020a). Other studies report negative correlations between the indices related to the soil aggregate stability, and the detachment capacity (e.g., De Baets et al., 2006; Wang et al., 2018). Moreover, the soil samples collected under the two conditions (treated and untreated plots) are clustered in two well-discriminated groups, and this may be a result of the differences in the analyzed soil properties (organic matter, aggregate stability, bulk density and rill detachment). This means that the changes in these properties due to biochar addition to soil may alter the soil erodibility. It is noteworthy that the distinction between untreated soils and sites supplied with substrates that are rich in organic matter (such the rice husk biochar) is consistent with the differences in soil erodibility and quality found between bare and hydromulched soils in a previous study carried out in the same environment (Parhizkar et al., 2021b).

4.2. Influence of rice husk biochar on rill erodibility

468

469

470

471

472

473

474

475

476

477

478

479

480

481

482

483

484

485

486

487

488

489

490

491

492

493

494

495

496

497

498

499

500

501

This study has demonstrated that the soil treatment with biochar is able to reduce rill erodibility. For the untreated soil, this parameter was about two-fold the value measured in the treated soil. This result is in accordance with Seitz et al. (2020), who found that the soil erodibility decreases in sandy and silty soils with biochar addition. Moreover, Li et al. (2021) showed that biochar successfully decreases flow turbulence on rills formed by erosion and thus reduces the sediment transport rate on hillslopes. Overall, the soil conservation practices based on treatment with biochar can increase the water storage capacity of soil and prevent soil particle detachment (Nicosia et al., 2021). In contrast, biochar addition could increase soil erosion, especially in cultivated lands (Zhang et al., 2019).

In this investigation, the linear regression model based on the interpolation between rill detachment capacity and shear stress is very accurate, as shown by the high coefficients of determination calculated for the two soil conditions. This result agrees with the results of the previous works of Parhizkar et al. (2020a; 2020b; 2020c; 2021b) in the same study area, and Zhang et al. (2008). These studies also report very high coefficients of determination for equations established between the rill detachment capacity and shear stress under different experimental conditions.

It is important to highlight that the soil erodibility estimated for the soil treated with rice husk biochar is higher compared to the values measured for hydromulched soils in the previous work by Parhizkar et al. (2021b). This can be due to the beneficial effects of hydromulch roots on soil properties, which exert a noticeable reduction in rill erodibility. These beneficial effects are in accordance with the outcomes of Mamo and Bubenzer (2001), who report that the presence of living plant roots supports the reduction in rill detachment capacity and erodibility of soils. However, the hydrological effects of soil treatment with biochar develop more quickly compared to grass growth in the hydromulched sites, and this characteristic may be appreciated in emergency actions for soil conservation.

Overall, the results of this study show that the reduced values of rill erodibility in sites treated with rice husk biochar in comparison to the untreated soils may be of interest for land planners, in order to implement suitable soil conservation programs to reduce the soil erosion risks. Actually, the rill erodibility, as being the slope of the equation interpolating rill detachment capacity and shear stress, is a measure of the soil resistance to rill erosion (Nearing et al., 1992). Therefore, the precise estimation of rill erodibility is necessary to implement effective soil conservation actions.

5. Conclusions

In the case study of deforested hillslopes of Northern Iran, rill detachment capacity in soil treated with rice husk biochar was significantly lower compared to the untreated sites. In contrast, the organic carbon and aggregate stability of soils were significantly higher and the bulk density was lower. Rill detachment capacity was not correlated to the other physico-chemical properties of soil. Rill erodibility of deforested hillslopes, calculated by linear regressions between rill detachment capacity and shear stress, was markedly lower in the treated soil compared to the untreated sites. These findings highlight the effectiveness of the rice husk biochar to control and mitigate soil detachment and improve organic matter content and aggregate stability in deforested hillslopes. Moreover, the proposed values of rill erodibility and shear stress for deforested hillslopes (treated or not) are helpful for both hydrologists (who must reproduce soil resistance to rill erosion in process-based erosion models) and land planners (who must setup the most effective soil conservation actions) working in delicate ecosystems, such as deforested lands that are exposed to intense erosion.

534

502

503

504

505

506

507

508

509

510

511

512

513

514

515

516

517

518

519

520

521

522

523

524

525

526

527

528

529

530

531

532

533

535 Funding

536

537 Faculty of Agricultural Sciences, University of Guilan.

528	
556	

539 Acknowledgments

540

The authors thank the Faculty of Agricultural Sciences, University of Guilan for their support and experimental assistance.

- 543
- 544

References

- 545
- Abrahams, A.D., Parsons, A.J., Luk, S.H., 1985. Field measurement of the velocity of overland flow using dye tracing. Earth Surface Processees and Landforms 11, 653–657.
- 548 Ahmadi, S.H., Ghasemi, H., Sepaskhah, A.R., 2020. Rice husk biochar influences runoff features,
- soil loss, and hydrological behavior of a loamy soil in a series of successive simulated rainfall
 events. Catena 192, 104587.
- Allison, L.E., 1975. Organic Carbon. In: CA, Black (Ed.), Methods of Soil Analysis, Part 2.
 American Society of Agronomy, Madison, WI, pp. 1367–1378.

Burrell, L.D., Zehetner, F., Rampazzo, N., Wimmer, B., Soja, G., 2016. Long-term effects of
biochar on soil physical properties. Geoderma 282, 96–102.

⁵⁵⁵ De Baets, S., Poesen, J., Gyssels, G., Knapen, A., 2006. Effects of grass roots on the erodibility of ⁵⁵⁶ topsoils during concentrated flow. Geomorphology 76, 54–67.

557 Domene, X., Mattana, S., Hanley, K., Enders, A., Lehmann, J., 2014. Medium-term effects of corn

- ⁵⁵⁸ biochar addition on soil biota activities and functions in a temperate soil cropped to corn. Soil
 ⁵⁵⁹ Biology And Biochemistry 72, 152–162.
- Dume, B., Mosissa, T., Nebiyu, A., 2016. Effect of biochar on soil properties and lead (BD) availability in a military camp in South West Ethiopia. African Journal of Environmental Science and Technology 10 (3), 77-85.
- Gamage, D.N.V., Mapa, R.B., Dharmakeerthi, R.S., Biswas, A., 2016. Effect of rice-husk biochar
 on selected soil properties in tropical Alfisols. Soil Research 54, 302-310.
- 565 Ghorbani, M., Asadi, H., Abrishamkesh, S., 2019. Effects of rice husk biochar on selected soil
- 566 properties and nitrate leaching in loamy sand and clay soil. International Soil and Water
- 567 Conservation Research 7, 258–265.

- ⁵⁶⁸ Gurwick, N.P., Moore, L.A., Kelly, C., Elias, P., 2013. A systematic review of biochar research, ⁵⁶⁹ with a focus on its stability in situ and its promise as a climate mitigation strategy. PLoS ONE 8(9),
- 570 e75932.
- ⁵⁷¹ Hseu, Z-Y., Jien, S-H., Chien, W-H., Liou, R-C., 2014. Impacts of Biochar on Physical Properties
- and Erosion Potential of a Mudstone Slopeland Soil. The Scientific World Journal 2014 (6), 602197.
- Herath, H., Camps-Arbestain, M., Hedley, M., 2013. Effect of biochar on soil physical properties in
 two contrasting soils: An alfisol and an andisol. Geoderma 209, 188–197.
- IRIMO (Islamic Republic of Iran Meteorological Organization), 2016. Annual Rainfall Report.
 Available online: www.irimo.ir (accessed on 20 September 2021).
- Jiang, F., He, K., Huang, M., Zhang, L., Lin, G., Zhan, Z., Li, H., Lin, I., Håkansson, J., Ge, H.,
- Huang, Y., 2020. Impacts of near soil surface factors on soil detachment process in benggang
 alluvial fans. Journal of Hydrology 590, 125274.
- Jien, S-H., Wang, C-S., 2013. Effects of biochar on soil properties and erosion potential in a highly weathered soil. Catena 110, 225-233.
- Govers, G., Giménez, R., Van Oost, K. 2007. Rill erosion: exploring the relationship between
 experiments, modelling and field observations. Earth-Science Reviews 84(3-4), 87-102.
- 585 Kemper, W.D., Rosenau, R.C., 1986. Aggregate stability and size distribution. In: Method of Soil
- Analysis. part 1. Physical and mineralogical methods. Agronomy Monographs 9, Klute, A. (Eds.),
- 587 American Society of Agronomy, Madison, pp. 425-442.
- Khanal, A., Fox, G.A., 2017. Detachment characteristics of root-permeated soils from laboratory jet
 erosion tests. Ecological Engineering 100, 335-343.
- Kottek, M., Grieser, J., Beck, C., Rudolf, B., Rubel, F., 2006. World Map of the Köppen-Geiger
 climate classification updated. Meteorolgische Zeitschrift 15, 259–263.
- ⁵⁹² Li, T., Yu, P., Liu, D., Fu, Q., Hou, R., Zhao, H., Xu, S., Zuo, Y., Xue, P., 2021. Effects of Biochar
- on Sediment Transport and Rill Erosion after Two Consecutive Years of Seasonal Freezing and
- Thawing. Sustainability 13, 6984.
- 595 Li, Y., Xu, X.Q., Zhu, X.M., 1992. Preliminary-study on mechanism of plant-roots to increase soil antiscouribility on the
- Loess Plateau. Science in China Series B-Chemistry 35, 1085–1092. Liu, J., Zhou, Z., Zhang, X.J., 2019. Impacts
- of sediment load and size on rill detachment under low flow discharges. Journal of Hydrology 570,
 719–725.
- ⁵⁹⁹ Lucas-Borja, M.E., Delgado-Baquerizo, M., 2019a. Plant diversity and soil stoichiometry regulates
- the changes in multifunctionality during pine temperate forest secondary succession. Science of the
- 601 Total Environment 697, 134204.

- Lucas-Borja, M.E., Zema, D.A., Plaza-Álvarez, P.A., Zupanc, V., Baartman, J., Sagra, J., de las
 Heras, J., 2019b. Effects of different land uses (abandoned farmland, intensive agriculture and
 forestland) on soil hydrological properties in Southern Spain. Water 11, 503.
- Mamo, M., Bubenzer, G.D., 2001. Detachment rate, soil erodibility, and soil strength as influenced
 by living plant roots part I: Laboratory study. Transactions of the ASABE 44(5), 1167-1174.
- Masulili, A., Utomo, W.H., Syechfani, M., 2010. Rice husk biochar for rice based cropping system in acid soil 1. The characteristics of rice husk biochar and its influence on the properties of acid sulfate soils and rice growth in West Kalimantan Indonesia. Journal of Agricultural Science 2(1),
- 610 **39**.
- Nearing, M.A., Bradford, J.M., Parker, S.C., 1991. Soil detachment by shallow flow at low slopes.
- 612 Soil Science Society America Journal 55, 339–344.
- Nearing, M.A., Bradford, J.M., Parker, S.C., 1992. Soil detachment by shallow flow at low slopes.
- 614 Soil Science Society America Journal 56, 1654.
- Nearing, M.A., Foster, G.R., Lane, L.J., Finkner, S.C., 1989. A process based soil erosion model
- for USDA Water Erosion Prediction Project technology. Transactions of the ASAE 32, 1587–
 1593.
- Nearing, M.A., Norton, L.D., Bulgakov, D.A., Larionov, G.A., West, L.T., Dontsova, K.M., 1997.
- Hydraulics and erosion in eroding rills. Water Resources Research 33, 865–876.
- Nguyen, B., Lenmann, J., Kinyangi, J., Smernik, R., Riha, S., Engelhard, M., 2008. Long-term
 black carbon dynamics in cultivated soil. Biogeochemistry 89, 295–308.
- Nicosia, A., Pampalone, V., Ferro, V., 2021. Effects of Biochar Addition on Rill Flow Resistance.
 Water 13, 3036.
- Obia, A., Mulder, J., Martinsen, V., Cornelissen, G., Børresen, T., 2016. In situ effects of biochar on
 aggregation, water retention and porosity in light-textured tropical soils. Soil Tillage Research 155,
 35–44.
- Odhiambo, B.O., Mihara, M., 2018. Effects of Gypsum and Rice Husk Biochar on Surface
 Discharge and Nutrient Loss from Farmlands in Budalangi, Kenya. International Journal of
 Environmental and Rural Development 9, 94-100.
- Parhizkar, M., Shabanpour, M., Khaledian, M., Asadi, H., 2021a. The evaluation of soil detachment
- capacity induced by vegetal species based on the comparison between natural and planted forests.
- ⁶³² Journal of Hydrology 595, 126041.
- Parhizkar, M., Shabanpour, M., Khaledian, M., Cerdà, A., Rose, C.W., Asadi, H., Lucas-Borja,
- 634 M.E., Zema, D.A., 2020a. Assessing and Modeling Soil Detachment Capacity by Overland Flow in
- ⁶³⁵ Forest and Woodland of Northern Iran. Forests 11(1), 65.

- Parhizkar, M., Shabanpour, M., Lucas-Borja, M.E., Zema, D.A. 2021b. Hydromulch roots reduce
 rill detachment capacity by overland flow in deforested hillslopes. Journal of Hydrology 598,
 126272.
- Parhizkar, M., Shabanpour, M., Lucas-Borja, M.E., Zema, D.A. 2021c. Variability of rill
 detachment capacity with sediment size, water depth and soil slope in forest soils: A flume
 experiment. Journal of Hydrology 601, 126625.
- Parhizkar, M., Shabanpour, M., Lucas-Borja, M.E., Zema, D.A., Li, S., Tanaka, N., Cerdà, A.,
- 2020b. Effects of length and application rate of rice straw mulch on surface runoff and soil loss
 under laboratory simulated rainfall. International Journal of Sediment Research 36, 468-478.
- Parhizkar, M., Shabanpour, M., Miralles, I., Cerdà, A., Tanaka, N., Asadi, H., Lucas-Borja, M.E.,
- Zema, D.A., 2021d. Evaluating the effects of forest tree species on rill detachment capacity in a
 semi-arid environment. Ecological Engineering 161, 106158.
- Parhizkar, M., Shabanpour, M., Miralles, I., Zema, D.A., Lucas-Borja, M.E., 2021e. Effects of plant
 species on soil quality in natural and planted areas of a forest park in northern Iran. Science of the
 Total Environment 778, 146310.
- Parhizkar, M., Shabanpour, M., Zema, D.A., Lucas-Borja, M.E., 2020c. Rill Erosion and Soil
 Quality in Forest and Deforested Ecosystems with Different Morphological Characteristics.
 Resources 9 (11), 129.
- Pandian, K., Subramaniayan, P., Gnasekaran, P., Chitraputhirapillai, S., 2016. Effect of biochar
 amendment on soil physical, chemical and biological properties and groundnut yield in rainfed
 Alfisol of semi-arid tropics. Archives of Agronomy and Soil Science 62(9), 1293-1310.
- Peng, X., Ye, L.L., Wang, C.H., Zhou, H., Sun, B., 2011. Temperature and duration-dependent rice
 straw-derived biochar: Characteristics and its effects on soil properties of an Ultisol in southern
- ⁶⁵⁹ China. Soil and Tillage Research 112, 159–166.
- Prakongkep, N., Gilkes, R., Wisawapipat, W., Leksungnoen, P., Inboonchuay, T., Delbos, E.,
- 661 Strachan, L-J., Ariyasakul, P., Ketdan, C., Hammecker, C., 2021. Effects of Biochar on Properties
- of Tropical Sandy Soils under Organic Agriculture. Journal of Agricultural Science 13, 1–17.
- Qu, J., Li, B., Wei, T., Li, C., Liu, C., 2014. Effects of rice-husk ash on soil consistency and
 compactibility. Catena 122, 54–60.
- Radcliffe, D.E., Simunek, J., 2010. Soil physics with HYDRUS: Modeling and applications. CRC
 press, <u>Boca Raton, Florida, USA</u>.
- 667 SDSD (Soil Science Division Staff), 2017. Soil Survey Manual, Ditzler, C., Scheffe, K., Monger,
- 668 H.C., Eds, USDA Handbook 18 Government Printing Office: Washington, DC, USA.

- 669 Seitz, S., Teuber, S., Geißler, C., Goebes, P., Scholten, T., 2020. How Do Newly-Amended Biochar
- Particles Affect Erodibility and Soil Water Movement?—A Small-Scale Experimental Approach.
- 671 Soil Systems 4 (4), 60.
- 672 Shabanpour, M., Daneshyar, M., Parhizkar, M., Lucas-Borja, M.E., Zema, D.A., 2020. Influence of
- crops on soil properties in agricultural lands of northern Iran. Science of the Total Environment134694.
- Ulyett, J., Sakrabani, R., Kibblewhite, M., Hann, M., 2014. Impact of biochar addition on water
- retention, nitrification and carbon dioxide evolution from two sandy loam soils. European Journal of
- 677 Soil Science 65, 96–104.
- Van Liew, M.W., Arnold, J.G., Garbrecht, J.D., 2003. Hydrologic simulation on agricultural
 watersheds: Choosing between two models. Transactions of the ASAE 46 (6), 1539-1551.
- Verheijen, F.G.A., Jeffery, S., Bastos, A.C., van der Velde, M., Diafas, I., 2009. Biochar
 Application to Soils A Critical Scientific Review of Effects on Soil Properties, Processes and
 Functions. EUR 24099 EN, Office for the Official Publications of the European Communities,
- 683 Luxembourg, pp. 149.
- Wang, B., Zhang, G.H., Shi, Y.Y., Zhang, X.C., 2014. Soil detachment by overland flow under
 different vegetation restoration models in the Loess Plateau of China. Catena 116, 51–59.
- Wang, B., Zhang, G.H., Yang, Y.F., Li, P.P., Liu, J.X., 2018. The effects of varied soil properties
 induced by natural grassland succession on the process of soil detachment. Catena 166, 192–199.
- Wang, D.D., Wang, Z.L., Shen, N., Chen, H., 2016. Modeling soil detachment capacity by rill flow
 using hydraulic parameters. Journal of Hydrology 535, 473–479.
- Wick, A.F., Huzurbazar, S.V., Stahl, P.D., 2009. Use of Bayesian methods to model soil
 aggregation in undisturbed semiarid grasslands. Soil Science Society of America Journal 73, 1707–
 1714.
- Wardle, D.A., Nilsson, M.C. Zackrisson, O., 2008. Fire-derived charcoal causes loss of forest
 humus. Science 320(5876), 629.
- ⁶⁹⁵ Zeraatpisheh, M., Ayoubi, S., Mirbagheri, Z., Mosaddeghi, M.R., Xu, M., 2021. Spatial prediction
- of soil aggregate stability and soil organic carbon in aggregate fractions using machine learning
- algorithms and environmental variables. Geoderma Regional 27, 400-440.
- ⁶⁹⁸ Zhang, F., Huang, C., Yang, M., Zhang, J., Shi, W., 2019. Rainfall simulation experiments indicate
- that biochar addition enhances erosion of loess-derived soils. Land Degradation and Development
 30, 2272–2286.
- Zhang, G.H., Liu, G.B., Tang, K.M., Zhang, X.C., 2008. Flow detachment of soils under different
- ⁷⁰² land uses in the Loess Plateau of China. Transaction of the ASABE 51, 883–890.

Zhang, G.H., Tang, K.M., Zhang, X.C., 2009. Temporal variation in soil detachment under different
land uses in the Loess Plateau of China. Earth Surface Processes and Landforms 34, 1302–1309.