



Università degli Studi Mediterranea di Reggio Calabria
Archivio Istituzionale dei prodotti della ricerca

An Integrated Pest Management Program Outperforms Conventional Practices for Tomato (*Solanum lycopersicum*) in Cambodia

This is the peer reviewed version of the following article:

Original

An Integrated Pest Management Program Outperforms Conventional Practices for Tomato (*Solanum lycopersicum*) in Cambodia / Porras, Mf; Malacrinò, A; An, C; Seng, Kh; Socheath, O; Norton, G; Miller, S; Rosa, C; Rajotte, Eg; O'Rourke, Me. - In: PLANT HEALTH PROGRESS. - ISSN 1535-1025. - 23:2(2022), pp. 206-211. [10.1094/PHP-09-21-0123-RS]

Availability:

This version is available at: <https://hdl.handle.net/20.500.12318/130247> since: 2022-10-18T13:06:53Z

Published

DOI: <http://doi.org/10.1094/PHP-09-21-0123-RS>

The final published version is available online at: <https://apsjournals.apsnet.org/doi/10.1094/PHP-09-21-0123-RS>

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website

Publisher copyright

This item was downloaded from IRIS Università Mediterranea di Reggio Calabria (<https://iris.unirc.it/>) When citing, please refer to the published version.

(Article begins on next page)

An Integrated Pest Management Program Outperforms Conventional Practices for Tomato (*Solanum lycopersicum*) in Cambodia

Mitzy F. Porras, Antonino Malacrinò, Chanratha An, Kim Hian Seng, Ong Socheath, George Norton, Sally Miller, Cristina Rosa, Edwin G. Rajotte, Megan E. O'Rourke

Journal: Plant Health Progress

DOI: 10.1094/PHP-09-21-0123-RS

URL: <https://apsjournals.apsnet.org/doi/10.1094/PHP-09-21-0123-RS>

Publisher: The American Phytopathological Society (APS)

Please, cite this article as:

Mitzy F. Porras, Antonino Malacrinò, Chanratha An, Kim Hian Seng, Ong Socheath, George Norton, Sally Miller, Cristina Rosa, Edwin G. Rajotte, and Megan E. O'Rourke (2022). An Integrated Pest Management Program Outperforms Conventional Practices for Tomato (*Solanum lycopersicum*) in Cambodia. *Plant Health Progress* 23:2, 206-211

Mitzy F. Porras

Plant Health Progress

An integrated pest management program outperforms conventional practices for tomato (*Solanum lycopersicum* L.) in Cambodia

Mitzy F. Porras ^{a*}, Antonino Malacrino ^b, Chanratha An ^{c,d}, Kim Hian Seng ^d, Ong Socheath ^c, George Norton ^c, Sally Miller ^f, Cristina Rosa ^g, Edwin G. Rajotte ^a, Megan E. O'Rourke ^{h,i}

^a*Department of Entomology, Pennsylvania State University, 501 ASI Bldg. University Park, Pennsylvania 16802 USA.

^b Department of Evolution, Ecology, and Organismal Biology, The Ohio State University, Columbus, Ohio 43210 USA.

^c Royal University of Agriculture, Phnom Penh, Cambodia

^d IDE Cambodia, Phnom Penh, Cambodia

^e Department of Agriculture and Applied Economics, Virginia Tech, Blacksburg, VA 24061, USA.

^f Department of Plant Pathology, The Ohio State University, Wooster, OH 44626 USA.

^g Department of Plant Pathology, Pennsylvania State University, 321 Buckhout Lab. University Park, Pennsylvania 16802 USA.

^h School of Plant and Environmental Sciences, Virginia Tech, Blacksburg, VA, USA. ⁱPresent address: Institute of Bioenergy, Climate, & Environment, USDA-NIFA, MO 6414, USA.

*Corresponding author: M.F. Porras; E-mail::mfp145@psu.edu

Abstract

For several years, pest management in tomato production in Cambodia has generally focused on the use of synthetic pesticides. We compared conventional pest management (farmers' traditional practices) with an IPM program on 12 farms in the northwestern part of Cambodia. The IPM program

combined cultural, biological, and chemical practices. We found that IPM practices reduced aphid damage by 46% and diseases such as Fusarium wilt and damping-off were substantially reduced. Our results indicate that the IPM package increased tomato yield and income by an average of 23% and 34%, respectively, compared to conventional practices during both dry and rainy seasons.

Keywords: Arthropods, disease, biological control, cultural control, chemical control

Funding: the United States Agency for International Development under Cooperative Agreement No. AID-OAA-L-15-00001

Introduction

Tomato (*Solanum lycopersicum* L.) (Solanaceae) is one of the most important vegetables in Cambodia. The total area cultivated is 1,065 ha and represents a source of employment for more than 1,400 families in rural and suburban areas (World Bank Group 2015). Few Cambodian provinces supply vegetable markets, it appears that only six provinces produce 30% of consumed vegetables. The other 70% are imported from Vietnam (Eliste and Zorya 2015). Two main varieties of tomato, Neang Pich and Neang Tamm, are cultivated in Cambodia (Mund 2011), but the yield is severely affected by insect pests and pathogens, causing losses of 40% (Schreinemachers et al. 2015). Despite the importance of tomato and its severe pest problems, the need to establish integrated pest management (IPM) programs has received little attention in Cambodia. This lack of attention has resulted in the overuse of pesticides, increased pesticide resistance, reduction of natural enemies, and growing threats to the environment and human health (Mund 2011). Adopting an IPM approach that integrates cultural, biological, and chemical practices may increase tomato yield while reducing dependence on harmful pesticides (Mund 2011).

Numerous pests can cause economic losses in tomato crops; however, little is known about tomato pests in Cambodia. Tomato crops in Cambodia can be affected by pest species reported in other countries in Southeast Asia. For example, many studies report that several arthropods cause damage to the foliage and fruits, including spider mites (*Tetranychus urticae* Koch), and insects such as armyworms

(*Spodoptera exigua* (Hübner) and *S. litura* (Fabricius), leafminer (*Liriomyza* sp.), tomato budworm (*Helicoverpa armigera* (Hübner)), whiteflies (e.g., *Bemisia tabaci* (Gennadius)), aphids (*Myzus persicae* Sulzer and *Machrosiphum euphorbiae* Thomas), and thrips (e.g., *Thrips palmi* Karny) (McDougall et al. 2013) (Sen 2016). Bacteria, fungi, and viruses can also cause significant damage in tomato, for example, bacterial wilt (*Ralstonia pseudosolanacearum*) (Klass et al. 2020) are important diseases, Fusarium wilt (*Fusarium oxysporum* f. sp. *lycopersici*), powdery mildew (*Sphaerotheca fuliginea*), and late blight (*Phytophthora infestans*). The most abundant viruses in tomato crops are *Tomato yellow leaf curl virus* (TYLCV) (Tolin and Fayad 2016, Yule et al. 2019). Currently, insect pests are managed using synthetic pesticides such as imidacloprid, pyrethroids, and organophosphates and effective fungicides are lacking except for late blight management. These pesticides are often misused, resulting in added cost to the farmer, inadequate pest control, reduction of ecosystem services such as pollination, and water pollution (Kevan and Phillips 2001).

IPM is a strategic approach to managing pests that uses multiple tactics to manage all of the pest threats observed in a crop system (Barzman et al. 2015). The first step in an IPM approach requires proper identification of pest and pathogen species as well as beneficial species. Once these species are known, an IPM Package can be designed (Dinakaran et al. 2013). An IPM package consists of tactics or practices that address pest problems in a crop in a given location over a whole growing season (Barzman et al. 2015, Grasswitz 2019).

Confirmation of the pest in a given field can be done using systematic observations. Some pest species can be anticipated because they have appeared in previous years. Preventative measures may be incorporated into the package based on this anticipation. These measures may include selection of resistant crop varieties, manipulation of planting dates, soil amendments, sanitizing seed treatments, encouragement of natural enemies by establishing plant species that are food sources for natural enemies and refraining from using pesticides that are harmful to natural enemies, mechanical barriers such as using a row cover on the seedbed, and so on (Grasswitz 2019). For IPM decisions, a regular monitoring regime (scouting) is established to ascertain when pest species appear during a season, how much of a

threat they are to the crop, and whether natural enemies are present and offer pest suppression. Monitoring can be aided by sex pheromone traps and sticky cards, as well as human observations (Barzman et al. 2015). Within-season control practices can include rogueing of diseased plants and judicious pesticide use when necessary based on crop monitoring. Each practice is selected based on its efficacy, economic benefits, and risks to human health and the environment (Barzman et al. 2015). A challenge for developing and implementing an IPM program is to successfully balance yield, quality, and profits while reducing pesticide use. Several IPM programs have been introduced for tomato production in Asia. Pretty and Bharucha (2015) analyzed IPM implementation programs in several countries in East Asia and Africa and found yield increases from 3.3 to 17.5 t/ha, demonstrating the critical role of IPM for the food production. In Bangladesh, where the pesticide applications were significantly reduced, profit increased by 39% (Rahman et al. 2018).

The development of appropriate management strategies is essential to increase yield and profit for Cambodian tomato farmers. Here, we explored the impacts of an IPM package on pest infestation, yield, income, cost, and profit on 12 tomato farms during both dry and wet seasons in 2019. We hypothesized that the IPM package would be at least as effective as conventional management at controlling tomato pests and that profitability would not be harmed by adopting an IPM management package. The research was done on farmers' fields using a participatory research approach (Rajotte et al. 2005).

2. Materials and methods

Field trials were conducted during the dry season (October 2018-March 2019) and wet season (May 2019-October 2019) in 12 farms located in four districts, including Chi Kraeng, Angkor Thom, Batheay Srei and Pouk of Siem Reap, Cambodia. Dry season temperatures ranged from 21.5 to 36°C, with a total rainfall of 370 mm, with 18–71% relative humidity (Weather Stats 2021). The rainy season had total rainfall of 1050 mm (Weather Stats 2021), with temperatures ranging from 22.5 to 35.5°C and 33–100% relative humidity. We followed an on-farm research approach that underscored participatory experimental trials and surveys (Norton et al. 1999, Darnhofer et al. 2012) essentially as

described in Malacrinò et al. (2020). Briefly, we compared IPM and conventional management treatments and a control. The IPM treatment protocol was standardized across farms (details below) (Dinakaran et al. 2013) whereas the conventional treatment comprised a mix of farm practices; farmers managed their conventional plots according to their usual practices.

2.2. Agronomy

Soil was plowed and fertilized with compost/manure (5 t ha^{-1}) and lime (320 kg ha^{-1}) about two weeks before transplanting. Next, raised beds (80 cm wide x 30 cm high) with a 70 cm between rows and a drip irrigation system lengthwise in the middle of each bed and covered with silver plastic mulch a week before transplanting. Tomato plants (3-week old) were transplanted at a density of 2 plants m^{-2} and vertically grown on bamboo and wooden poles (1 m) and plastic string. Tomato plants were irrigated daily or every two days according to crop needs. Plots were fertilized with 50 kg ha^{-1} N, 60 kg ha^{-1} P, and 250 kg ha^{-1} K, and 18 kg ha^{-1} Ca and 0.3 kg ha^{-1} Mg during both dry and rainy seasons. Fertilization was divided between basal fertilizer and fertigation applications with 40% of total N and 30% of total P, and just 5% of total K; the basal application was performed during the second plow and weekly fertigation applications.

2.3. Pest management

The IPM treatment targeted aphids, fruit borer, leaf miner, thrips, spider mite, whitefly, worms (*Helicoverpa armigera* and *Spodoptera* spp.), bacterial wilt, damping off, Fusarium wilt, and powdery mildew (Table 1). IPM treatment comprised seedlings planted in seed trays in coco peat (Jiffy DecoGro, Jiffy International AS, Norway) mixed with *Trichoderma viride* (Royal University of Agriculture, Cambodia), a biofungicide to manage diseases caused by fungal pathogens, at 1 g L^{-1} of coco peat, protected by insect exclusion nets (50 mesh inch $^{-1}$). Then, seedlings were transplanted into the plot and watered in with a *T. viride* and cow manure slurry solution after three weeks.

Next, plants were sprayed weekly with a mix of three biopesticides (*Bacillus thuringiensis* (*Bt*) subsp. *aizawa* (TAB 1, Ladda Co., Ltd., Thailand), *Bacillus subtilis* (Tab Innovation Co., Ltd., Thailand), and *Beauveria bassiana* (Tab Innovation Co., Ltd., Thailand) (Table S1) at 0.4 kg, 0.25 kg ha⁻¹, and 0.4 kg ha⁻¹ respectively. In addition, we periodically sprayed sodium tetraborohydrate decahydrate in orange essential oil (Prev-Am, ORO Agri Inc, USA) when needed to manage insect damage (aphids, whiteflies, or thrips) at 0.4% per 100 gal. Furthermore, conventional pesticides were used in the trial including imidacloprid (Confidor 200SL, Bayer, Germany) was sprayed once by each farmer at a rate of 667 ml ha⁻¹ during the dry season to control thrips, and metalaxyl (Galaxy 35WP, Papaya Trade Co., Cambodia) for late blight sprayed one or two times as needed at a rate of 590 g ha⁻¹, once or twice a week during the rainy season (Table S1). We used pheromone lures and yellow sticky traps (35 x 40 cm) for monitoring insect populations. Disease incidence and severity were monitored from plant symptoms in the field (10 plants/plot). The conventional treatment is summarized in Table S2. Seedlings were grown in seed trays using field soil; in total, fifteen different pesticides were sprayed (11 insecticides and four fungicides) during both dry and rainy seasons. Tomato plants of control plots were grown in coco peat trays without insecticide or fungicide applications.

IPM package components were selected using our experience in similar systems in other countries such as Bangladesh, Nepal, and India. The work in Cambodia is considered adaptive research, so the successful packages developed in other countries would be validated in Cambodia (Rajotte et al. 2005, Dinakaran et al. 2013, Malacrino et al. 2019)

Table 1. Arthropod pest and pathogens scored in tomato plants during both dry and rainy seasons in small-farm scales

Pest category	Common name	Scientific name
Arthropods	Aphid	<i>Macrosiphum euphorbiae</i>

Fruit borer	<i>Leucinodes orbonalis</i> Guenée	
Leaf miner	<i>Liriomyza sativae</i>	
Thrip	<i>Frankliniella schultzei</i>	
Spider mite	<i>Tetranychus urticae</i>	
White fly	<i>Bemisia tabaci</i> (Genadius.)	
Worms	<i>Helicoverpa</i>	
	<i>Spodoptera</i> spp.	
Pathogens	Bacterial wilt	<i>Ralstonia</i> spp.
	Damping off	<i>Phytophthora</i> spp.
	Fusarium wilt	<i>Fusarium</i> spp.
	Powdery mildew	<i>Oidium</i> spp.

2.4. Data collection and analysis

We scouted all plots weekly and selected five random plants per treatment and measured damage by arthropod pests and pathogens (Table 1). We followed a protocol essentially as described in Malacrino et al. (2020), in which pest and disease levels on each plant were scored on a scale from 0 to 3, which was granularly calibrated for replicability between farms. Damage or severity was classified as follows: 0 = none, 1 = little, 2 = moderate, and 3 = severe (Table S3). We also measured yield, input costs (excluding labor), income from sales, and profit. Input costs included expenses from fertilizers to trellis system, values were adjusted for depreciation. Farmers documented yield, including marketable weight (kg) and price of tomatoes per kg (U.S. dollars [USD]) for each treatment. Then, we calculated costs and benefits for each treatment by subtracting total expenses from income produced by marketable tomatoes.

For statistical analyses, we used a Generalized Linear Mixed-Effects Model (GLMM) with treatments (IPM, conventional, and control) as a fixed effect, and pest, farm, and sampling week as random variables. Analyses of damage scores for each species pest were similarly analyzed, without speciespest as a random variable. Models were fitted with a Poisson family error distribution because data were counts and log link function using the *glmer* function from the *lme4* package of R (Bates et al. 2014). Then, we assessed a treatment's effect using Type II Wald χ^2 from the package *car* and *Anova* function. Post-hoc analyses such as pairwise contrasts using Tukey's multiple comparisons were conducted using the package *emmeans*. Pests for which at least one treatment did not have an average damage score of at least 0.02 (i.e., at least 1 of 50 plants was scored as 1 or higher per week) were not statistically analyzed due to lack of data and model convergence issues.

We used a Linear Mixed-Effects Model (LME) to analyze yields, costs, incomes, and profits data using the *lmer* function with treatment as a fixed effect and farm as a random variable, then pairwise contrasts were inferred using Tukey's multiple comparisons, as described above. Since growing

conditions were different between dry and rainy seasons, statistical analyses were separately performed for each season using R 4.0.2 (Team 2013).

3. Results

3.1. Damage

The global analysis (GLMM) showed no significant differences in disease incidence among the three treatments. However, during the dry season, Fusarium wilt and damping-off was decreased by 45% and 55%, respectively, in IPM plots compared to conventional management and control plots (Fig. 1, Table 2). In contrast, bacterial wilt and powdery mildew were not significantly affected by management practices. In 2019 (wet season), disease incidence followed a similar trend as in 2018. (Table 2).

GLMM showed no significant differences in arthropod pests among the three types of plots. However, the incidence of worms (*Helicoverpa* and/or *Spodoptera* spp.) was reduced by 25% (Fig. 1, Table 3). During the wet season, the arthropod pest populations decreased in all experimental plots; however, the lowest pest incidences were observed in plots where the IPM practices were implemented (Table 3).

3.2. Productivity

During the dry season, tomato yield increased 6% in plots that used IPM practices compared to plots using conventional or no management. Income increased 9% and 14 %in IPM plots relative to conventional and control plots, respectively. The costs associated with using the IPM package in the dry season were 5% higher than in plots using conventional management and 29% higher than control plots (Table 4). During the rainy season, tomato yield in IPM plots was 8% higher than in plots using conventional management and 32% higher than the untreated control. On average, the income during the rainy season in IPM plots increased by 63%, and the costs were only 4% higher than plots using conventional management and 11% higher than control. The cost associated with IPM package was 3.5% higher than in plots using conventional management and 26% higher than control plots (Table 4). The profit in IPM plots was 15% higher than in conventional management and 40% higher than in control plots (Table 4).

Discussion

IPM practices have long been recognized as essential for sustainable tomato production (Strand 1998, Pretty and Bharucha 2015). Few studies, however, have examined the effects of IPM on tomato production in Cambodia (Mund 2011), and fewer still have evaluated the differences between IPM and conventional practices. The results of our field experiment demonstrate that using an IPM package reduces the incidence of arthropod pests and disease incidence in tomato crops during both dry and wet seasons in Cambodia.

Plots in which the IPM package was deployed had significantly higher yield and income than plots using conventional management. The IPM package uses cultural, physical, biological, and synthetic control tools, including several that increased the cost on average by 30% compared to conventional management. However, the successful control of tomato pests resulted in an average of both seasons of 45% more profit in plots using the IPM approach in both seasons than in those using conventional management.

The costs of IPM practices were higher than the conventional management and no-spray trials. However, adopting this IPM program led to substantial profit increases during the rainy season and a similar profit compared to the no management program. The IPM strategy used on these small farms was demonstrated to be a success when multiple tactics such as physical control and biological and synthetic pesticides were used. Thus, sustainable tomato production on small farms in Cambodia could benefit from adopting IPM packages that align with tomato plant development and the biology and ecology of pests and diseases.

Incidence of the soilborne diseases Fusarium wilt and damping-off was significantly reduced during both dry and wet seasons in tomatoes in the IPM program treated with *Trichoderma viride* as seedling transplants and subsequently with *Bacillus subtilis* and *Pseudomonas fluorescens* applications. These microbes have been shown to be successful biocontrol agents of pathogens such as root rot fungi (Elad 2000). Biocontrol agents degrade the cell walls of the plant pathogen (de la Cruz et al. 1995). Bacterial wilt, Fusarium wilt, powdery mildew, damping off, and virus symptoms were all detected but at low infection levels in IPM plots. This suggests the efficacy of the rational use of biopesticides within an IPM program (Dara 2016).

The numerical differences in arthropod pest incidence in IPM plots resulted from the combined effect of physical, biological, and synthetic control practices. Physical control, such as the use of insect exclusion nets during the seedling stage has been shown to reduce early insect infestation, resulting in a significant reduction in tomato damage (Caroline et al. 2017, Singh et al. 2018). Spraying *Bt* and *Beauveria bassiana* significantly reduced damage caused by insects during both dry and rainy seasons, suggesting a synergistic impact on a wide range of tomato pests. Several studies have shown that *Bt* bacterium controls many insect pests, including *Helicoverpa zea*, *Manduca sexta*, and *Spodoptera frugiperda* (Arthurs and Dara 2019). In addition, studies have shown that *Bt* reduces leafminer populations (Badran et al. 2016, Jamshidnia et al. 2018) and sucking insects such as whiteflies and aphids (Badran et al. 2016). Aphid populations were absent during the rainy season, possibly due to the environmental constraints and colonies of *B. bassiana* in the farms. The effects of the fungus *B. bassiana*

on the control of insect pest populations have been well documented (Wari et al. 2020). Other biological control agents such as *Bt* directly impact caterpillars attacking foliage and fruits (i.e., *S. frugiperda*, *M. sexta*). Alternating biopesticides and synthetic insecticides seem to be a good alternative to reduce arthropod pest impact as well as insecticide use in tomato crops of Chi Kraeng, Angkor Thom, Batheay Srei and Pouk, Siem Reap provinces.

Small-scale farms are a fundamental component of agricultural production in Cambodia. They have an important role in promoting social benefits of sustainable agriculture, including the conservation of natural resources and biodiversity, as well as the enhancement of local food security (Grasswitz 2019). Low tomato production in Cambodia may result from the combined effect of poor soil fertility, unfavorable weather, poor-quality seeds, and high cost of inputs and/or poor postharvest knowledge of the growers (Bhattarai et al. 2011). Growing tomato can provide high returns to small-scale farmers, thus reducing the importation from neighboring countries such as Vietnam and Thailand (Bhattarai et al. 2011).

Challenges associated with the cost-effectiveness of IPM may be addressed by prioritizing phytosanitary problems per location; thus, different IPM packages may be designed in response to a broader range of farmers' needs. An IPM approach may reduce the risk associated with the development of resistance in tomato pests. The success of tomato production in Cambodia relies on a better understanding of the pest population dynamics combined with cultural, physical, and rational applications of biopesticides and synthetic pesticides.

In summary, our results indicate that our IPM package approach can significantly reduce tomato pest severity and disease incidence and that rational use and timing of biopesticide applications can result in sustainable insect and disease control over time. Moreover, this study demonstrates the cost-effectiveness of implementing IPM tactics on small farms.

Acknowledgments

We thank the United States Agency for International Development under Cooperative Agreement No. AID-OAA-L-15-00001 to Virginia Tech for the Feed the Future Innovation Lab for Collaborative Research on Integrated Pest Management (IPM IL). The opinions expressed herein are those of the authors and do not necessarily reflect the views of the U.S. Agency for International Development. We also thank anonymous reviewers for their comments and suggestions to this work.

Competing interests

The authors declare no competing interests.

Data availability

The data that support the findings of this study are available on request from the corresponding author MFP and MEO

References

- Arthurs, S., and S. K. Dara.** 2019. Microbial biopesticides for invertebrate pests and their markets in the United States. *Journal of invertebrate pathology* 165: 13-21.
- Badran, A. H., V. M. Guzov, Q. Huai, M. M. Kemp, P. Vishwanath, W. Kain, A. M. Nance, A. Evdokimov, F. Moshiri, and K. H. Turner.** 2016. Continuous evolution of *Bacillus thuringiensis* toxins overcomes insect resistance. *Nature* 533: 58-63.
- Bank, W.** 2015. Cambodian agriculture in transition: opportunities and risks. World Bank Group, Agriculture Washington, DC.
- Barzman, M., P. Bärberi, A. N. E. Birch, P. Boonekamp, S. Dachbrodt-Saaydeh, B. Graf, B. Hommel, J. E. Jensen, J. Kiss, and P. Kudsk.** 2015. Eight principles of integrated pest management. *Agronomy for sustainable development* 35: 1199-1215.
- Bates, D., M. Mächler, B. Bolker, and S. Walker.** 2014. Fitting linear mixed-effects models using lme4. arXiv. org. preprint]. doi 10.
- Bhattarai, S. P., M. C. Palada, D. J. Midmore, D. Wu, and R. Salas.** 2011. ON-FARM EVALUATION OF FERTILIZER BRIQUETTES AND LOW-COST DRIP IRRIGATION FOR SMALLHOLDER VEGETABLE PRODUCTION IN CAMBODIA. *Irrigation and Drainage* 60: 318-329.
- Caroline, L. J., M. Saidi, and A. Opiyo.** 2017. Effect of Coloured Agro-Net Covers on Insect Pest Control and Yield of Tomato (*Solanum lycopersicum* Mill). *Journal of Agricultural Science* 9.
- Darnhofer, I., D. Gibbon, and B. Dedieu.** 2012. Farming systems research: an approach to inquiry, pp. 3-31, Farming systems research into the 21st century: The new dynamic. Springer.
- Dinakaran, D., G. Gajendran, S. Mohankumar, G. Karthikeyan, S. Thiruvudainambi, E. Jonathan, R. Samiyappan, D. Pfeiffer, E. G. Rajotte, and G. Norton.** 2013. Evaluation of Integrated Pest and Disease Management Module for Shallots in Tamil Nadu, India: a Farmer's Participatory Approach. *Journal of Integrated Pest Management* 4: B1-B9.
- Eliste, P., and S. Zorya.** 2015. Cambodian agriculture in transition: Opportunities and risks. The World Bank.

- Grasswitz, T. R. 2019.** Integrated Pest Management (IPM) for small-scale farms in developed economies: Challenges and opportunities. *Insects* 10: 179.
- Jamshidnia, A., S. Abdoli, S. Farrokhi, and R. Sadeghi. 2018.** Efficiency of spinosad, *Bacillus thuringiensis* and *Trichogramma brassicae* against the tomato leafminer in greenhouse. *BioControl* 63: 619-627.
- Kevan, P. G., and T. P. Phillips. 2001.** The economic impacts of pollinator declines: an approach to assessing the consequences. *Conservation ecology* 5.
- Klass, T. L., M. M. Hayes, K. H. Seng, C. An, F. Rotondo, W. Shoaf, S. Ong, K. Tho, C. Allen, and S. Miller. 2020.** First Report of Bacterial Wilt of Tomato Caused by *Ralstonia pseudosolanacearum* (*Ralstonia solanacearum* phylotype I) in Cambodia. *Plant Disease* 104: 969-969.
- Malacrinò, A., K. H. Seng, C. An, S. Ong, and M. E. O'Rourke. 2019.** Integrated pest management for yard-long bean (*Vigna unguiculata* subsp. *Sesquipedalis*) in Cambodia. *Crop Protection*: 104811.
- Malacrinò, A., K. H. Seng, C. An, S. Ong, and M. E. O'Rourke. 2020.** Integrated pest management for yard-long bean (*Vigna unguiculata* subsp. *Sesquipedalis*) in Cambodia. *Crop Protection* 135: 104811.
- McDougall, S., A. Watson, B. Stodart, G. Kelly, D. Troldahl, and L. Tesoriero. 2013.** Tomato, capsicum, chilli and eggplant: a field guide for the identification of insect pests, beneficials, diseases and disorders in Australia and Cambodia, Australian Centre for International Agricultural Research (ACIAR).
- Mund, J.-P. 2011.** The agricultural sector in Cambodia: Trends, processes and disparities. *Pacific News* 35: 10-14.
- Norton, G. W., E. G. Rajotte, and V. Gapud. 1999.** Participatory research in integrated pest management: Lessons from the IPM CRSP. *Agriculture and Human Values* 16: 431-439.
- Pretty, J., and Z. P. Bharucha. 2015.** Integrated pest management for sustainable intensification of agriculture in Asia and Africa. *Insects* 6: 152-182.
- Rahman, M. S., G. W. Norton, and M. H.-A. Rashid. 2018.** Economic impacts of integrated pest management on vegetables production in Bangladesh. *Crop Protection* 113: 6-14.
- Rajotte, E. G., G. W. Norton, G. C. Luther, V. Barrera, and K. Heong. 2005.** IPM transfer and adoption. Globalizing Integrated Pest Management: A Participatory Research Process: 143-157.
- Schreinemachers, P., S. Balasubramaniam, N. M. Boopathi, C. V. Ha, L. Kenyon, S. Praneetvatakul, A. Sirijinda, N. T. Le, R. Srinivasan, and M.-H. Wu. 2015.** Farmers' perceptions and management of plant viruses in vegetables and legumes in tropical and subtropical Asia. *Crop Protection* 75: 115-123.
- Sen, S. 2016.** Development of an IPM approach in controlling insect pest of tomato DEPARTMENT OF ENTOMOLOGY, SHER-E-BANGLA AGRICULTURAL UNIVERSITY, DHAKA.
- Singh, A., L. Sah, Y. D. GC, M. Devkota, L. Colavito, B. Rajbhandari, G. Norton, E. Rajotte, and R. Muniappan. 2018.** Evaluation of pest exclusion net to major insect pest of tomato in Kavre and Lalitpur. Nepalese Journal of Agricultural Sciences 16: 128-137.
- Strand, L. R. 1998.** Integrated pest management for tomatoes, vol. 3274, UCANR Publications.
- Team, R. C. 2013.** R: A language and environment for statistical computing.
- Tolin, S. A., and A. Fayad. 2016.** Virus diseases of tropical vegetable crops and their management, pp. 41-76, Integrated pest management of tropical vegetable crops. Springer.
- Wari, D., R. Okada, M. Takagi, M. Yaguchi, T. Kashima, and T. Ogawara. 2020.** Augmentation and compatibility of *Beauveria bassiana* with pesticides against different growth stages of *Bemisia tabaci* (Gennadius); an in-vitro and field approach. *Pest Management Science*.
- Yule, S., P. Chiemsothab, and R. Srinivasan. 2019.** Detection of Tomato yellow leaf curl Thailand virus transmitted by *Bemisia tabaci* Asia I in tomato and pepper. *Phytoparasitica* 47: 143-153.

- 1 **Table 2.** Disease severity in tomato plants during both dry and rainy seasons in small-farm scales. Dry season: $\chi^2=8.81$; $P=0.01216$; rainy season:
 2 $\chi^2=19.25$, $P < 0.0001$ (mean \pm se).

Plant pathogen	Dry season					Rainy season				
	χ^2	P	Control	Conventional	IPM	χ^2	P	Control	Conventional	IPM
Bacterial wilt	4.19	0.1225	0.12 \pm 0.05	0.12 \pm 0.05	0.05 \pm 0.03	0.99	0.6086	0.15 \pm 0.05	0.17 \pm 0.05	0.12 \pm 0.04
Fusarium wilt	11.09	0.003	0.55 \pm 0.10	0.49 \pm 0.10	0.27 \pm 0.07	17.59	0.0001	0.55 \pm 0.10	0.49 \pm 0.10	0.15 \pm 0.07
Powdery mildew	0.01	0.9944	0.5 \pm 0.04	0.49 \pm 0.04	0.5 \pm 0.04	0.06	0.9692	0.50 \pm 0.04	0.49 \pm 0.04	0.50 \pm 0.04
Damping-off	6.83	0.028	0.27 \pm 0.07	0.25 \pm 0.07	0.12 \pm 0.05	13.5	0.0011	0.27 \pm 0.07	0.25 \pm 0.07	0.12 \pm 0.05
Unknown disease	0.13	0.9351	1.00 \pm 0.11	1.05 \pm 0.10	1.00 \pm 0.11	0	1.00	0.30 \pm 0.11	0.30 \pm 0.10	0.30 \pm 0.11

3 **Table 3.** Arthropod pest populations in tomato plants during both dry and rainy seasons in small-farm scales. Dry season: $\chi^2=10.23; P=0.01216$;
 4 rainy season: $\chi^2=19.25, P < 0.0001$ (mean \pm se).

Arthropod pest	Dry season					Rainy season				
	χ^2	P	Control	Conventional	IPM	χ^2	P	Control	Conventional	IPM
Aphids	2.31	0.3138	0.11 \pm 0.02	0.08 \pm 0.02	0.05 \pm 0.02	NA	NA	NA	NA	NA
Fruit borer	0.01	0.9933	0.42 \pm 0.04	0.41 \pm 0.04	0.41 \pm 0.04	0.8	0.6676	0.42 \pm 0.04	0.41 \pm 0.04	0.41 \pm 0.04
Leaf miner	0.19	0.9057	0.62 \pm 0.04	0.58 \pm 0.04	0.59 \pm 0.04	1.08	0.581	0.62 \pm 0.04	0.58 \pm 0.04	0.59 \pm 0.04
Whitefly	0.24	0.8825	0.9 \pm 0.02	0.85 \pm 0.03	0.93 \pm 0.03	1.33	0.5133	0.98 \pm 0.02	0.87 \pm 0.03	0.84 \pm 0.03
Worm	2.43	0.2954	0.48 \pm 0.048	0.52 \pm 0.04	0.39 \pm 0.04	6.98	0.0304	0.48 \pm 0.04	0.52 \pm 0.04	0.20 \pm 0.04

5 **Table 4.** Analysis of yield, income, costs, and profit from tomato dry season: $\chi^2=3.69$; $P=0.0412$; and rainy season: $\chi^2=26.11$, $P < 0.0001$ (mean ±
6 se). Different letters indicate significant pairwise differences between values within the same column at the $P < 0.05$ level

Season	Treatment	Yield (kg m⁻²)	Income (USD m⁻²)	Costs (USD m⁻²)	Profit (USD m⁻²)
Dry	Control	2.67±0.11 (a)	0.79±0.04 (a)	0.51±0.007 (a)	0.21±0.04 (a)
	Conventional	2.91±0.13 (a)	0.79±0.04 (a)	0.68±0.007 (b)	0.14±0.04 (a)
	IPM	3.10±0.11 (b)	0.84±0.04 (b)	0.72±0.007 (c)	0.12±0.04 (b)
	<i>F</i>	7.78	7.39	264.86	3.69
	<i>DF</i>	22	22	22	22
	<i>P</i>	0.0027	0.0034	<0.0001	0.04
Rainy	Control	1.69±0.12 (a)	1.05±0.08 (a)	0.51±0.003 (a)	0.52±0.08 (a)
	Conventional	2.28±0.12 (b)	1.41±0.08 (b)	0.63±0.003 (b)	0.78±0.08 (b)
	IPM	2.49±0.12 (b)	1.54±0.08 (b)	0.66±0.003 (b)	0.88±0.08 (b)
	<i>F</i>	40.38	41.43	801.10	20.18
	<i>DF</i>	22	22	22	22
	<i>P</i>	<0.0001	<0.0001	<0.0001	<0.0001

7 **Supplementary material**8 **Table S1.** Pest control tools used under IPM during rainy and dry seasons.

Season	Category	Biocontrol agent	Total	Average number
		/Active ingredient	number of farms	of applications (±SEM)
Rainy	Biocontrol of pathogens	<i>Trichoderma viridae</i>	12	2
	Biocontrol of insects	<i>Bacillus subtilis</i>	12	7
	Biocontrol of insects	<i>Bacillus thuringiensis</i>	12	7
	Biocontrol of insects	<i>Beauveria bassiana</i>	12	7
	Bactericide	Copper 17.5%+Streptomyce n 5%+Zine Sulphate 10% WP	12	2
Dry	Biocontrol of pathogens	<i>Trichoderma harzianum</i>	12	2
	Biocontrol of insects	<i>Bacillus subtilis</i>	12	4±0.16

Biocontrol of insects	<i>Bacillus</i> <i>thuringiensis</i>	12	5±0.18
Biocontrol of insects	<i>Beauveria bassiana</i>	12	6
Insecticide	Orange oil	12	2±0.25
	<i>Pseudomonas</i> <i>fluorescens</i>	12	4±0.61
Bactericide	Copper	9	1±0.5
	17.5%+Streptomyci n 5%+Zine		
	Sulphate 10% WP		

10 **Table S2.** Pest control tools used under conventional management during rainy and dry seasons.

Season	Category	Active ingredient	Total	Average number number of farms (±SEM)	Average number of applications
Rainy	Fungicide	Mancozeb 80%	1	2	
		Mancozeb+Metalaxy 18%) + Imidacloprid	1	2	
		Matalaxyl 35WP	7	1.85±0.26	
		Abamectin 0.4g/L + Beta-cypermethrin	4	2±0.70	
		1.6g/L			
		Acetamiprid 150g + Buprofezin 150g	2	1	
	Insecticide	Acetamiprid + Buprofezin	2	2	
		Acetamiprid 20%+Imidacropid	2	2	
		16%			
		Azoxystrobin	2	4	
		Chlorantraniliprole	1	3	
	Copper	12	2.16±0.32		
		17.5%+Streptomycin			

		5%+Zine Sulphate		
		10% WP		
		Copper hydroxide	2	3.5±0.57
		Emamectin Benzoate	4	2.5±0.86
		5% W/W + Lambda		
		cyhalothrin 10%		
		W/W		
		Copper Oxychloride	12	2.75±0.16
		45% + Kasugamycin		
		2% WP		
		Imidacropid	2	2
		Imidacropid 50g/L +	1	2
		Chlorpyrifos Ethyl		
		200g/L		
		Thiamethoxam	6	2±0.57
		250g/kg &		
		Acetamiprid 100g/kg		
Dry	Fungicide	Mancozeb 64%+	1	5
		Metalazyil 8%		
	Bactericide	Mancozeb 80%	3	3±0.57
	Insecticide	Methalazy	1	3
		Pymetrozine 50%	1	2

Thiamethoxam	1	1
250g/kg+Acetamipri		
d 100g/kg		
Emamectin Benzoate	2	3.5±0.5
4%		
Emamectin Benzoate	1	2
5%		
Emamectin Benzoate	4	2.5±0.86
5% W/W + Lambda		
cyhalothrin 10%		
W/W		
Emamectin Benzoate	1	4
5.5%		
Emamectin Benzoate	2	2
50g/Kg+Emamectin		
Benzoate 5% W/W +		
Lambda		
Imidacropid	1	2
Mancozeb 64%+	1	5
Metalazyil 8%		
Mancozeb 80%	3	3±0.57
Methalazy	1	3

Pymetrozine 50% 1 2


Thiamethoxam 1 1

250g/kg+Acetamipri

d 100g/kg

11 **Table S3.** Severity scoring definitions for each stage of the tomato plant.

Severity score	Vegetative	Flowering	Fruit set and harvesting
1 = Little	Few insects or signs of damage on up to 10% of leaves	Few insects or signs of damage on up to 10% of lower and older leaves; up to 20% of flowers dropped	Few insects or signs of damage on up to 10% of lower and older leaves; up to 30% of flowers dropped; few fruits damaged
2 = Moderate	More insects plus 10-20% of leaves damaged by insects or diseases	More insects plus 10-50% of leaves damaged by insects or diseases; 20-30% of flowers dropped	More insects plus 10-50% of leaves damaged by insects or diseases; 30-50% of flowers dropped; 10%-20% of fruit damaged
3 = Severe	Many insects plus 20%+ of leaves damaged by insects or diseases; stunting, stem or root rot and permanent wilt	Many insects plus 50%+ of leaves damaged by insects or diseases; stunting, stem rot or permanent wilt; 30%+ of flowers dropped	Many insects plus 50%+ of leaves damaged by insects or diseases; stunting, stem rot or permanent wilt; 50%+ of flowers dropped; 20%+ of fruit damaged



12 **Figure 1.** Infestation scores for tomato under conventional management, IPM or control (no
13 management) during rainy (A) and dry (B) seasons in Cambodia. Bars represent mean \pm SE.