



REPLY TO ELMENDORF AND ETTINGER:

Photoperiod plays a dominant and irreplaceable role in triggering secondary growth resumption

Jian-Guo Huang^{a,b,1}, Filipe Campelo^c, Qianqian Ma^{a,b}, Yaling Zhang^{a,b}, Yves Bergeron^d, Annie Deslauriers^e, Patrick Fontf^f, Eryuan Liang^g, Harri Mäkinen^h, Walter Oberhuberⁱ, Cyrille B. K. Rathgeber^j, Roberto Tognetti^k, Václav Trem^l, Bao Yang^m, Lihong Zhai^b, Jiao-Lin Zhang^{n,b}, Serena Antonucci^k, J. Julio Camarero^o, Katarina Čufar^p, Henri E. Cuny^q, Martin De Luis^r, Alessio Giovannelli^s, Jožica Gričar^t, Andreas Gruberⁱ, Vladimír Gryc^{u,v}, Aylin Güney^{v,w}, Xiali Guo^{a,b}, Wei Huang^x, Tuula Jyske^h, Jakub Kašpar^l, Gregory King^{f,y}, Cornelia Krause^e, Audrey Lemay^e, Feng Liu^{z,b}, Fabio Lombardi^{aa}, Edurne Martínez del Castillo^r, Hubert Morin^e, Cristina Nabais^c, Pekka Nöjd^h, Richard L. Peters^{f,bb}, Peter Prislan^p, Antonio Saracino^{cc}, Irene Swidrakⁱ, Hanuš Vavřík^u, Joana Vieira^c, Biyun Yu^{a,b}, Shaokang Zhang^{a,b}, Qiao Zeng^{dd}, Emanuele Ziaco^{ee}, and Sergio Rossi^{aa,e}

In their Letter, Elmendorf and Ettinger (1) question the dominant role of photoperiod in driving secondary growth resumption (hereafter referred to as xylem formation onset) of the Northern Hemisphere conifers, recently reported by Huang et al. (2). Their opinions are grounded on the following three aspects, including 1) the seasonality of the photoperiod, 2) the dependence of the predictor variables (e.g., photoperiod, forcing, and chilling) on the response variable (the date of onset of xylem formation, day of the year [DOY]), and 3) the limit value of the obtained models for interannual forecasting. We think they bring up an interesting issue that deserves further discussion and clarification.

Photoperiod is acknowledged to regulate spring bud swelling while wood formation starts (3, 4). Although photoperiod seasonality occurs at each site, its influence is marginal in our study given that the analysis involved comparisons among sites across the Northern Hemisphere. Our conclusion that photoperiod plays a dominant role was built upon the combination of several coherent pieces of evidence, rather than “the crux of Huang et al. ...” as they pointed out. First, we clearly stated that model 2, which modeled DOY as a function of the mean annual temperature of the site (MAT), forcing, chilling, and soil moisture, was considered the best model in terms of parsimony

^aKey Laboratory of Vegetation Restoration and Management of Degraded Ecosystems, South China Botanical Garden, Chinese Academy of Sciences, Guangzhou, 510650, China; ^bCenter of Plant Ecology, Core Botanical Gardens, Chinese Academy of Sciences, Guangzhou, 510650, China; ^cCentre for Functional Ecology, Department of Life Sciences, University of Coimbra, 3000-456 Coimbra, Portugal; ^dForest Research Institute, Université du Québec en Abitibi-Témiscamingue, Rouyn-Noranda, QC J9X5E4, Canada; ^eDépartement des Sciences Fondamentales, Université du Québec à Chicoutimi, Chicoutimi, QC G7H 2B1, Canada; ^fSwiss Federal Research Institute for Forest, Snow and Landscape, CH-8903 Birmensdorf, Switzerland; ^gKey Laboratory of Alpine Ecology and Biodiversity, Key Laboratory of Tibetan Environment Changes and Land Surface Processes, Institute of Tibetan Plateau Research, Chinese Academy of Sciences, Beijing 100101, China; ^hDepartment of Forests, Natural Resources Institute Finland, 02150 Espoo, Finland; ⁱDepartment of Botany, Leopold-Franzens University of Innsbruck, 6020 Innsbruck, Austria; ^jAgroParisTech, Institut National de Recherche pour l'Agriculture, l'Alimentation et l'Environnement, Université de Lorraine, Silva, F-54000 Nancy, France; ^kDipartimento di Agricoltura, Ambiente e Alimenti, Università degli Studi del Molise, 86100 Campobasso, Italy; ^lDepartment of Physical Geography and Geoecology, Charles University, CZ-12843 Prague, Czech Republic; ^mCold and Arid Regions Environmental and Engineering Research Institute, Chinese Academy of Sciences, Lanzhou 730000, China; ⁿChinese Academy of Sciences Key Laboratory of Tropical Forest Ecology, Xishuangbanna Tropical Botanical Garden, Chinese Academy of Sciences, Mengla, Yunnan 666303, China; ^oInstituto Pirenaico de Ecología, Consejo Superior de Investigaciones Científicas, 50192 Zaragoza, Spain; ^pBiotechnical Faculty, University of Ljubljana, 1000 Ljubljana, Slovenia; ^qDepartment of Forest and Carbon Resources, Institut National de Géographie et Forestière, 54250 Champigneulle, France; ^rDepartment of Geography and Regional Planning, Environmental Science Institute, University of Zaragoza, 50009 Zaragoza, Spain; ^sIstituto di Ricerca sugli Ecosistemi Terrestri, Consiglio Nazionale delle Ricerche, 50019 Sesto Fiorentino, Italy; ^tLaboratory for Dendrochronology, Slovenian Forestry Institute, 1000 Ljubljana, Slovenia; ^uDepartment of Wood Science and Wood Technology, Mendel University in Brno, 61300 Brno, Czech Republic; ^vInstitute of Botany, University of Hohenheim, 70593 Stuttgart, Germany; ^wSouthwest Anatolia Forest Research Institute, 07010 Antalya, Turkey; ^xState Key Laboratory for Conservation and Utilization of Subtropical Agro-bioresources, College of Life Sciences, South China Agricultural University, Guangzhou 510642, China; ^yDepartment of Sciences, University of Alberta, Camrose, AB T4V 2R3, Canada; ^zKey Laboratory of Aquatic Botany and Watershed Ecology, Wuhan Botanical Garden, Chinese Academy of Sciences, Wuhan 430074, China; ^{aa}Dipartimento di Agraria, Università Mediterranea di Reggio Calabria, 89122 Reggio Calabria, Italy; ^{bb}Laboratory of Plant Ecology, Department of Plants and Crops, Faculty of Bioscience Engineering, Ghent University, B-9000 Ghent, Belgium; ^{cc}Department of Agricultural Sciences, University of Naples Federico II, I-80055 Portici-Napoli, Italy; ^{dd}Key Lab of Guangdong for Utilization of Remote Sensing and Geographical Information System, Guangdong Open Laboratory of Geospatial Information Technology and Application, Guangzhou Institute of Geography, Guangzhou 510070, China; and ^{ee}DendroLab, Department of Natural Resources and Environmental Science, University of Nevada, Reno, NV 89557

Author contributions: J.-G.H., F.C., Q.M., Y.Z., Y.B., A.D., P.F., E.L., H. Mäkinen, W.O., C.B.K.R., R.T., V.T., B. Yang, L.Z., J.-L.Z., S.A., J.J.C., K.C., H.E.C., M.D.L., A. Giovannelli, J.G., A. Gruber, V.G., A. Güney, X.G., W.H., T.J., J.K., G.K., C.K., A.L., F. Liu, F. Lombardi, E.M.d.C., H. Morin, C.N., P.N., R.L.P., P.P., A.S., I.S., H.V., J.V., B. Yu, S.Z., Q.Z., E.Z., and S.R. performed research; J.-G.H., F.C., and Q.M. analyzed data; and J.-G.H. wrote the paper.

The authors declare no competing interest.

Published under the [PNAS license](#).

¹To whom correspondence may be addressed. Email: huangjg@scbg.ac.cn.

according to minimum Akaike information criterion and Bayesian information criterion, rather than R^2 as referred to in their Letter. Second, photoperiod interacted with MAT and can explain 61.7% of the variance of MAT alone (2). Therefore, we concluded that secondary growth resumption was driven primarily by MAT and photoperiod or by their interaction, which is challenging to be disentangled without experimental data, we agree. In terms of biological functioning, they play an undoubtedly relevant role in secondary growth resumption, as explained in Huang et al. (2) and Delpierre et al. (5). In contrast, Elmendorf and Ettinger do not provide any biological reason why photoperiod would not control secondary growth resumption, but acknowledge that “on a yearly basis plants may rely on cues beyond average temperature.” Finally, following Elmendorf and Ettinger’s approach, we found that their models lack biological foundation and their findings are at least questionable or even unrealistic. Therefore, we moderately adapted their code to avoid unrealistic aspects, resulting in more consistent insights than those they reported (Figs. 1 and 2).

According to traditional definitions, several exogenous factors were calculated and retained in the final models after rigorous statistical screening, such as collinearity tests. More importantly, these models are biologically sound and provide perspectives for driving future mechanistically related experimental designs, although we agree that statistical models are imperfect approximations to reality.

The recently developed microsampling technique used in Huang et al. (2) provides high time-resolution (weekly) data of xylogenesis for refining our knowledge on wood formation during the growing season. Therefore, we advocate applying our weekly data-based modeling results into earth system models to better understanding of carbon, water, and energy cycles at intra-annual scale (6), rather than at interannual scale that can be better assessed by dendroecology (7).

Acknowledgments

This work was funded by the National Natural Science Foundation of China (Grants 41861124001, 41661144007, and 31971499), the International Collaborative Key Project of the Chinese Academy of Sciences (Grant GJHZ1752), and Guangdong Natural Science Foundation (Grant 2019B121202007).




Fig. 1. Simulated phenological data [adapted from Elmendorf and Ettinger (1)], produced using mean annual temperature (MAT), latitude, and seasonal temperature variation (estimated for each site as the long-term mean difference between July and January temperatures). (A) Simulated time-series of temperature by Elmendorf and Ettinger ($\sigma = 4$), without an evident seasonal trend. Lines are simulated temperature series for each site and year, and the color represents the latitude. The black line represents the mean daily temperature for all sites. (B) Simulated time-series of temperature, considering mean annual temperature, temperature seasonality, and with a given level of “noise” ($\sigma = 4$). The detailed equation is as follows: $T_t = MAT - (T_{Jul} - T_{Jan})/2 \times \cos(2 \times \pi/365 \times (t - 15)) + Error$; where T_t is temperature for the DOY of the year (t), MAT is mean annual temperature, T_{Jul} (T_{Jan}) is the long-term mean temperature in July (January), and Error is the noise added ($\sigma = 4$). The vertical line in blue (dash in black) gives the mean day of the year (DOY) observed (simulated) of wood formation onset for all studied sites. The black line represents the mean daily temperature for all sites. (C) The phenological data were simulated using forcing unit (FU) thresholds described in Huang et al. (2). The gray lines represent seasonal forcing accumulation for the site “SIM” (LAT = 48° N), for the period 2002 to 2014. The horizontal black (blue) line is the mean DOY of onset observed (simulated) for the site “SIM.” The dots are simulated dates of wood formation onset at this site. Our simulated data varied forcing units required for onset of wood formation (FU) with latitude, as previously described by Huang et al. (2). (D) A linear relationship is found between DOY at the beginning of wood formation (DOY_{wf}) and MAT; the root-mean-square error (RMSE) of the simulated temperature and simulated (observed) phenology appears in black (blue). The color dots show dates of wood formation onset, with colors indicating latitude (legend on the Right of the panel). (E) Linear relationship between DOY_{wf} and photoperiod. Color lines show continuous patterns in photoperiod across the spring. For the observed phenology, the RMSE (in blue) for the photoperiod (E) is higher than the RMSE for MAT (D), which is held for σ values in the range 0 to 10. For estimated phenology, the RMSE (in black) for the photoperiod (E) is higher than for MAT (D); for time-series of simulated temperature with high noise level ($\sigma > 5$) the photoperiod shows a lower RMSE than MAT, but the simulated DOY_{wf} (horizontal black line in C) deviates from the observed DOY_{wf} (horizontal blue line in C). We found that photoperiod becomes a stronger predictor of phenology for high σ values (>5), while for lower values of σ phenology is mainly predicted by MAT alone.




Fig. 2. Predictive performance of the date of wood formation onset (DOY_{wf}) based on forcing, latitude, mean annual temperature (MAT), and photoperiod. Linear mixed models were fitted on randomly sampled subsets of the data, in each iteration 25% of sites were dropped (to evaluate the importance of environmental variables across spatial gradients in driving phenology; for each iteration the testing dataset consisted of sites in the latitude range used for training to reduce the bias introduced by the low distribution of sites at low and high latitudes) or a single year (to access the importance of environmental conditions on phenology at the temporal scale; only sites with phenological data over 3 y were considered; for each site 1 y was randomly sampled for testing and the remainder used for training). For latitude and MAT, we modeled the DOY_{wf} directly (e.g., $DOY_{wf} \sim MAT + \text{random effects}$), producing a DOY_{wf} prediction for the evaluation subset. For forcing, chilling, and photoperiod, we used the threshold value of the environmental variable at xylogenesis as the response variable (e.g., $FU_{wf} \sim \text{Intercept} + \text{random effects}$), and then used the simulated daily meteorological data with the modeled thresholds to predicted DOY_{wf} (e.g., the first DOY where FU exceeds the predicted FU_{wf} for a given site, species, and year). The predictive performance of each model was defined as the difference in predictive accuracy (root-mean-square error [RMSE] of predicted vs. observed DOY_{wf}) in relation to a null model with no environmental drivers for each iteration ($n = 100$). A negative (positive) $\delta_{RMSE}(\text{model} - \text{null})$ indicates that model with environmental drivers explains more (less) variation than the null model. In the spatial cross-validation (A), the proportion of negative $\delta_{RMSE}(\text{model} - \text{null})$ is 0.01 for MAT, while MAT presents a proportion of 0.11 negative δ_{RMSE} in the temporal analysis (B). Overall, the predictive performance of the models (A and B) is better than Elmendorf and Ettinger reported. Adapted with permission from ref. 1.

- 1 S. C. Elmendorf, A. K. Ettinger, Is photoperiod a dominant driver of secondary growth resumption? *Proc. Natl. Acad. Sci. U.S.A.*, 10.1073/pnas.2019411117 (2020).
- 2 J.-G. Huang *et al.*, Photoperiod and temperature as dominant environmental drivers triggering secondary growth resumption in Northern Hemisphere conifers. *Proc. Natl. Acad. Sci. U.S.A.* **117**, 20645–20652 (2020).
- 3 D. Basler, C. Körner, Photoperiod and temperature responses of bud swelling and bud burst in four temperate forest tree species. *Tree Physiol.* **34**, 377–388 (2014).
- 4 D. Basler, C. Körner, Photoperiod sensitivity of bud burst in 14 temperate forest tree species. *Agric. For. Meteorol.* **165**, 73–81 (2012).
- 5 N. Delpierre *et al.*, Chilling and forcing temperatures interact to predict the onset of wood formation in Northern Hemisphere conifers. *Glob. Change Biol.* **25**, 1089–1105 (2019).
- 6 H. E. Cuny *et al.*, Woody biomass production lags stem-girth increase by over one month in coniferous forests. *Nat. Plants* **1**, 15160 (2015).
- 7 F. O. Babst *et al.*, Twentieth century redistribution in climatic drivers of global tree growth. *Sci. Adv.* **5**, eaat4313 (2019).