



Università degli Studi Mediterranea di Reggio Calabria
Archivio Istituzionale dei prodotti della ricerca

Effects of two reflective materials on gas exchange, yield, and fruit quality of sweet orange tree *Citrus sinensis* (L.) Osb.

This is the peer reviewed version of the following article:

Original

Effects of two reflective materials on gas exchange, yield, and fruit quality of sweet orange tree *Citrus sinensis* (L.) Osb / Gullo, G.; Dattola, A.; Vonella, V.; Zappia, R.. - In: EUROPEAN JOURNAL OF AGRONOMY. - ISSN 1161-0301. - 118:(2020), pp. 1-9. [10.1016/j.eja.2020.126071]

Availability:

This version is available at: <https://hdl.handle.net/20.500.12318/67799> since: 2020-11-19T11:26:00Z

Published

DOI: <http://doi.org/10.1016/j.eja.2020.126071>

The final published version is available online at: <https://www.sciencedirect.com>.

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website

Publisher copyright

This item was downloaded from IRIS Università Mediterranea di Reggio Calabria (<https://iris.unirc.it/>) When citing, please refer to the published version.

(Article begins on next page)

1
2
3
4 Effects of two reflective materials on gas exchange, yield, and fruit quality of sweet
5
6 orange tree (*Citrus sinensis* (L.) Osb.)
7
8

9
10 Gregorio Gullo¹, Antonio Dattola, Vincenzo Vonella, Rocco Zappia
11

12
13 ¹Department AG.R.A.R.I.A. *Mediterranean* University of Reggio Calabria, Reggio Calabria, 89124 Italy,
14

15 UE – ggullo@unirc.it; antonio.dattola@unirc.it; vincenzo.vonella@unirc.it; rocco.zappia@unirc.it;
16

17 www.unirc.it
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32

33 e-mail: ggullo@unirc.it
34
35
36
37
38
39
40
41
42
43
44

45
46 *Abstract*
47
48

49 This study was conducted in an orange orchard (Navelina ISA 315 cv) located in southern Italy.
50

51 Leaf temperatures, Photosynthetic Photon Flux Density (PPFD), and UV-B reached high values
52 during many days in summer, causing photosynthesis (A_n) in the outer layers of the canopy to
53 decrease. This reduction was caused due to depression of stomatal conductance (g_s),
54 photoinhibition, and damage to the photosynthetic system. By contrast, A_n was lower in the inner
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

1 Reflective materials, either kaolin or calcium carbonate, were sprayed onto the canopy of the trees
2 every month from June to September. The behaviour of the treated trees was compared with that of
3 the untreated trees. The light reflected from the kaolin- (TK) and calcium carbonate-treated trees
4 (TCC) reduced the leaf temperature to an optimal value for photosynthesis (below 30 °C), and
5 increased the stomatal conductance (g_s). Furthermore, treatments reducing PPF_D played an
6 important role in protecting the chlorophyll pigments and increasing the maximum efficiency of
7 photosystem II (treated, ~0.67; untreated, 0.48). Consequentially, in the outer layer of the canopy
8 A_n was higher in treated trees (TK, 9.2 $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$; TCC, 7.6 $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$) than in the
9 control trees (TCN, 5.6 $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$). In the inner canopy layer, the reflective action of kaolin
10 and calcium carbonate improved light penetration and photosynthesis compared to the untreated
11 trees. The total average yield per tree over two years was 15% and 25% higher in TCC and TK,
12 respectively, compared with TCN. The fresh weight, total soluble solids, titratable acid, and the
13 ratio between these quantities were better in the treated groups than in the control, whereas no
14 differences were observed for nutraceutical parameters. Kaolin and calcium carbonate also
15 improved the peel colour of the fruits produced by the treated trees. In conclusion, reflective
16 materials improved the physiological, productive, and qualitative performance of orange trees, and
17 kaolin was the more effective of the two reflective materials. Even considering the cost, the use of
18 these reflective materials can be justified for citrus trees that yield high profits, such as the Navelina
19 ISA 315 used in our experiment.

20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48 **Keywords: calcium carbonate, kaolin, nutraceutical, Navelina, PPF_D**

49 50 51 52 53 54 **1. Introduction**

55 Under natural conditions, solar radiation is the most important source of energy for plants
56 (Larcher 1995). Light is a key factor for photosynthesis; however, excess light energy is deleterious
57 to the photosynthetic system. In many areas of the Mediterranean Basin, Photosynthetic Photon
58
59
60
61
62
63
64
65

1 Flux Density (PPFD), a component of light with a wavelength range of 400-760 nm, can exceed the
2 photosynthetic capacity of a leaf for many hours during clear summer days. Moreover, the reduction
3 of the ozone layer has also increased the terrestrial amount of ultraviolet (UV) radiation, a
4 component of light with a wavelength range of 280-400 nm. Increased UV-B radiation (wavelength
5 280 to 315 nm) also affects the photosynthetic process through direct or indirect damage
6
7 (Balakumar et al., 1993; Potters et al., 2009; Lidon et al., 2012; Mpoloka, 2008; Fedina et al.,
8
9 2010).
10
11
12
13
14
15

16 The activity of photosystem II (PSII) declines rapidly when the leaf is exposed to high light
17 intensity. This phenomenon is called photoinhibition (Long et al., 1994), and it involves a slow and
18 reversible reduction of photosynthetic efficiency that leads to a partial loss of capacity to convert
19 radiant energy into dry material (Long et al., 1994; Kitao et al., 2000). Photoinhibition can be
20 caused by ultraviolet light (UV-B), by visible light (V), or by their interaction (Powles, 1984).
21
22
23
24
25
26
27
28

29 Prolonged exposure of leaves to excessive radiation leads to photooxidation. This is a
30 secondary phenomenon occurring after photoinhibition has progressed as a function of light
31 intensity and exposure time (Powles, 1984; Long et al., 1994). Photooxidation causes the
32 destruction of photosynthesizing pigments (Powles, 1984), indicated by discolouration of these
33 pigments, and it may also cause cell death. Pigment bleaching happens when a certain degree of
34 photoinhibition has already occurred (Hendrey et al., 1987).
35
36
37
38
39
40
41
42

43 Under unfavourable conditions, such as thermal stress induced by IR radiation (a component
44 of light with a wavelength range of 760-4000 nm), the light saturation point of leaves occurs at
45 lower intensities compared to optimal conditions. Heat stress negatively influences the
46 carboxylation of ribulose biphosphate (RuBP) in photosynthesis; because of low RuBisCO activity
47 in the thylakoidal stroma, the light saturation lowers, and then photoinhibition phenomena start
48 sooner than in leaves at optimal temperature (Crafts-Brandner and Salvucci, 2000; Cen and Sage,
49 2005). Furthermore, recent studies have shown that thermal stress inhibits repair of the damage
50 caused to PSII by photoinhibition (Sharke, 2005; Allakhverdiev et al., 2008).
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 The leaf controls its temperature by reducing heat absorption through different mechanisms.
2 Among these, transpiration is utilized most often, controlled by adjusting the stomata openings
3 (Allakhverdiev et al., 2008; Yamamoto et al., 2008). However, this mechanism may be insufficient
4 for leaf cooling when leaf overheating is the result of a strong inflow of solar radiation, and the
5 temperature of the surrounding environment is high, even if water availability in the soil is optimal
6 (Nahal, 1981).
7

8
9
10
11
12
13
14 In the canopy profile, PPFD is higher on the canopy surface than on its under-layers because
15 the overlying canopy layers shade the ones underneath (Jackson et al., 1988). As a result, the
16 photosynthetic activity of leaves in the inner layers is low. The shaded environment of the forest
17 understorey is the original habitat of citrus species (Syvertsen and Lloyd, 1994; Davies and Albrigo,
18 1994; Spiegel-Roy and Goldschmidt, 1996), and thus the leaf has a low light saturation point
19 (Kriedemann, 1968; Sinclair and Allen, 1982). However, citrus trees are cultivated in a subtropical
20 climate where high solar radiation leads to high temperatures and light availability.
21

22
23
24
25
26
27
28
29
30
31 In recent years, many agriculture strategies have sought to mitigate the negative effects of
32 high irradiance and extreme temperature. Application of reflective material is one commonly used
33 approach. This treatment reduces heat and light stress in plants by reflecting solar radiation from the
34 foliar surface (Lombardini et al., 2004, Glenn et al. 2005, 2010). The aim of this work was to study
35 how high PPFD and high temperature affected leaf gas exchange, photosynthesis, and yield
36 production in citrus trees in full field, and to test whether new agronomic strategies, such as
37 reflective materials, can improve leaf physiology in the canopy and drive productive performance of
38 citrus trees during hot summers in the Mediterranean Basin.
39
40
41
42
43
44
45
46
47
48
49
50
51
52

53 **2. Materials and methods.**

54 *2.1 Orchard*

55
56
57
58 The experiment was conducted over two years, from 2016 to 2017, on the farm of
59 Cooperative OP Frujt, Serrata (RC), Italy, ([latitude, 38°30'30.1"N – longitude, 16°03'10.4" E](#)).
60
61
62
63
64
65

1 Orange trees (cultivar Navelina ISA 315) grafted onto sour orange trees were planted in the spring
2 of 1998 in pH 6 (subacidic) sandy soil, with 2.23% organic matter and 1.5 g·kg⁻¹ nitrogen content.
3
4 The plants were spaced 6 m × 4 m apart (417 plants ha⁻¹), and a north-to-south orientation was
5
6 adopted. Each tree was trimmed into a globe shape. The canopy volume was calculated following
7
8 the methodology proposed by Barrett and Brown (2012). The orchard was managed using a
9
10 standard integrated pest management system and stable drip irrigation and fertilisation system.
11
12
13

14 2.2 Treatment and experimental design

15
16 Surround WP and Purshade (both from Serbios Srl, Rovigo, Italy) are commercially
17
18 manufactured products containing 95% kaolin (kaolinite) and 95% calcium carbonate (calcite),
19
20 respectively. Both products have good properties for use as a reflective foliar coating. They were
21
22 applied at concentrations of 50 g·L⁻¹ (Surround WP) and 66.8g·L⁻¹ (Purshade). Trees sprayed with
23
24 kaolin and calcium carbonate were compared with control trees (trees sprayed with only water). A
25
26 total of 36 trees, three blocks of 12 plants with four plants per treatment, were arranged in a
27
28 randomised block design. Trees with a volume of approximately 30 m³ were selected. The reflective
29
30 materials, with 3 L of suspension sprayed per tree, uniformly covered the leaves of the treatment
31
32 trees. Treatments were applied using a trailed sprayers, every month during the summer season,
33
34 from June to September. After immature fruits dropped in June (“June drop”), the same number of
35
36 fruits was left on each tree (~460 fruits Tree⁻¹).
37
38
39
40
41
42
43

44 We determined the amount of kaolin and calcium carbonate (g·m⁻²) sprayed on the leaves in
45
46 the outer, intermediate, and inner layers of the canopy using a methodology similar to that adopted
47
48 by Jifon and Syvertsen (2003). Tester plates (polymethyl methacrylate , 3 cm × 3 cm) were
49
50 randomly distributed: ten testers were weighed and hung on each layer of the treated and untreated
51
52 trees. The testers were removed and reweighed after treatment when the reflective materials had
53
54 dried. The amount of reflective material was determined by the weight difference of the testers
55
56 before and after treatment. Both coated and uncoated testers were exposed to sunlight at noon on a
57
58
59
60
61
62
63
64
65

1 sunny day, and PPFD and UV radiations were recorded using PAR and UV sensors, respectively
2 (Spectrum Technologies Inc., Aurora, Illinois, USA), placed 10 cm below the testers.
3
4
5

6 7 *2.3 Measurements*

8
9 PAR and UV radiation were monitored on each tree using sensors installed on the three
10 layers. The sensors were connected to a logger that recorded the measurements every minute. Leaf
11 temperature was also monitored every minute, with fine wire thermocouples (GMR Instruments,
12 Firenze, Italy, UE) connected to a data logger (Spectrum Technologies Inc., Aurora, Illinois, USA).
13
14 The thermocouples were pressed against the abaxial leaf surface and held in place using lightweight
15 clips.
16
17
18
19
20
21
22
23

24 The leaf reflection (whiteness) was measured by a spectrophotometer (CR700; Minolta Cor.,
25 Ramsey, New Jersey, USA). Whiteness measurements (L values ranging from 0 to 100, where
26 black = 0 and white = 100) of the adaxial leaf surfaces were recorded before treatment when the
27 leaf was clean, and after treatment, when the individual leaf was coated with kaolin or calcium
28 carbonate.
29
30
31
32
33
34
35

36 Leaf ACO_2 , g_s , leaf-to-air vapour pressure deficit (LAVPD), leaf transpiration, and water
37 use efficiency (WUE; $ACO_2/Transpiration$) were measured for mature leaves in the outer,
38 intermediate, and inner layers of each tree (3 leaves \times 3 plants \times 3 layers \times 3 blocks). Measurements
39 were taken using a portable photosynthesis system (Li-Cor 6400XT; LI-COR Biosciences, Lincoln,
40 Nebraska, USA) and carried out on a clear, sunny summer day from 11:00 to 13:00, under optimal
41 weather conditions (average CO_2 partial pressure of 38 Pa and average PPFD of $1600 \mu mol \cdot m^{-2} \cdot s^{-1}$)
42 during the last week of the summer months in both years.
43
44
45
46
47
48
49
50
51
52

53 Photoinhibition was also estimated using a chlorophyll fluorometer (Li-Cor 6400-40; LI-
54 COR Biosciences, Lincoln, Nebraska, USA). Measurements were taken after 30 min of dark
55 adaptation using leaf clips on the same leaves used for gas exchange measurements. Leaves from
56
57
58
59
60
61
62
63
64
65

1 the outer layers were detached, washed with distilled water, and the SPAD 502 (Spectrum
2 Technologies Inc., Aurora, Illinois, USA) index was measured.
3

4 The PAR spectrum light was measured at midday (12:00) using a spectroradiometer (PS-
5 300; Apogee Instruments Inc., Logan, Utah, USA). We used an average of 4 points for each layer.
6

7 Soil water potential and plant water potential were continuously monitored (every five
8 minutes, 24 hours/day) with Watermark® sensors (Irrometer Company, Inc., Riverside, U.S.A.)
9 located at soil depths of 30 and 60 cm from the trunk, and with a PSY1 Stem Psychrometer (located
10 at a trunk height of 60 cm), respectively. The water tension value in the soil of the treated and
11 untreated trees was maintained between 30 cbars and field capacity during the experiment.
12
13
14
15
16
17
18
19
20
21
22
23

24 *2.4 Fruit sensorial and nutritional quality*

25 Observations were carried out at the *Colture Arboree* Laboratory of the AGRARIA
26 Department of The Mediterranean University of Reggio Calabria. For each tree, five fruits were
27 randomly selected, and their transversal diameter was measured every 14 days from mid-September
28 until mid-December. Concurrently, fruits from each treatment group and the control were sampled
29 (12 fruits per block). Immediately after harvesting, the colour of the skin and pulp was determined
30 in terms of CIELab and HSB colour spaces. Transversal and longitudinal diameters were measured,
31 and fresh weight (FW), dry weight (DW), and juice yield were determined using an electronic
32 balance (Mettler-Toledo, Greifensee, Switzerland). The juice was evaluated for total soluble solids
33 (SST) using a handheld digital refractometer (PR-1; Atago, Tokyo, Japan). Titratable acidity (TA)
34 was measured using 10 ml orange juice diluted with distilled water (1:1) and titrated to pH 8.2 with
35 0.1 N NaOH (mEq. NaOH 100 g⁻¹ fresh fruit). Ascorbic acid (AA) content was determined using a
36 procedure based on the reduction of 2,6-dichlorophenolindophenol (DIP) dye by ascorbic acid (mg
37 ascorbic acid 100 g⁻¹ FW).
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57

58 For each repetition, six fruits were placed in polyethene bags and frozen at -80 °C, until the
59 analyses of total polyphenol content (TPC) and total antioxidant capacity (TAC) were performed.
60
61
62
63
64
65

1 Pulp of the orange sample was homogenised using an Ultra-Turrax blender (20,000 rpm, T 25
2 Basic; IKA, Werke, Germany, UE). The TPC and TAC were separately analysed using a Lambda
3
4 35 spectrophotometer (PerkinElmer Corporation, Waltham, Massachusetts, USA). Before
5
6 measuring the TPC and TAC, standard curves were prepared for each test. The TPC (mg gallic acid
7
8 equivalents g^{-1} FW) was determined using the Folin–Ciocalteu method (Slinkard and Singleton,
9
10 1997). The TAC was determined using the modified TEAC assay and expressed as mmol Trolox
11
12 equivalents g^{-1} FW (Pellegrini et al., 1999; Re et al., 1999). The TEAC assay included both the
13
14 hydrophilic and the lipophilic contributions of the orange samples (Scalzo et al., 2005).
15
16
17
18
19
20
21

22 *2.5 Harvest*

23
24 The yield per tree (in weight and number of fruits) was determined during harvesting.
25
26
27

28 *Statistical analysis*

29
30 All data were analysed using one-way ANOVA tests for means comparisons with standard
31
32 errors. The data regarding fruit quality and nutritional analysis are reported as the means from both
33
34 years. The differences between plant development and production were calculated according to
35
36 Student-Newman-Keuls tests and were considered significant at $P < 0.05$. The canopy physiology
37
38 data were subjected to ANOVA tests, and the means were separated by Tukey's tests when the
39
40 ANOVAs indicated significant ($P < 0.05$) variable effects. All analyses were performed using the
41
42 IBM[®] SPSS[®] Statistics software, version 22 (SPSS Inc. IBM Company, Armonk, New York, USA).
43
44
45
46
47
48
49
50

51 **3. Results**

52
53 In our experiment, an average of 11.2 g of kaolin and 3.8 g of calcium carbonate per square
54
55 metre were deposited on the testers hanging on the outer layer of the canopy. In the intermediate
56
57 canopy layer, the reflective material on the testers was 50% lower for kaolin and 80% lower for
58
59 calcium carbonate, compared with the outer canopy layer. The lowest amount of reflective material
60
61
62
63
64
65

1 was recorded from the inner canopy layer, at $1.12 \text{ g}\cdot\text{m}^{-2}$ for kaolin and $0.38 \text{ g}\cdot\text{m}^{-2}$ for calcium
2 carbonate (Tab. 1). No differences were recorded in the water status of the treated and untreated
3 trees about stem and soil water potential (data not shown).
4
5
6

7 The leaf temperature and PPFD of the outer layer of the canopy in the control trees followed
8 the same pattern over 24 h. The pattern of leaf temperature from 10:00 to 16:00 was strongly
9 influenced by treatments with kaolin and calcium carbonate compared with the untreated trees in
10 the outer layer of the canopy (Fig. 1). The leaf temperature was $30 \text{ }^{\circ}\text{C}$ from 10:00 to 16:00, with a
11 peak of $34 \text{ }^{\circ}\text{C}$ from 11:00 to 12:00 (Fig. 1). The two treatments lowered the leaf temperatures of the
12 outer canopy layers to below $30 \text{ }^{\circ}\text{C}$ from 9:00 to 18:00. The average leaf temperatures of the outer
13 canopy layers decreased approximately 12% and 9% in TK ($-3.5 \text{ }^{\circ}\text{C}$) and TCC ($-2.9 \text{ }^{\circ}\text{C}$),
14 respectively, compared with the outer canopy layer of the control trees (Fig. 1). No differences in
15 leaf temperatures of the outer canopy layers were observed during the night between the treated and
16 untreated trees (Fig. 1). Thus, the long-wave emittance was unaffected by kaolin, corroborating
17 observations by Jifon and Syvertsen (2003). We observed the same with calcium carbonate
18 treatment. Leaf temperature was optimal in the intermediate and inner layers of the canopy without
19 significant differences between the treated and untreated trees ($\sim 24\text{-}25 \text{ }^{\circ}\text{C}$, data not shown).
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37

38 PPFD changed from 900 to $1200 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ from 11:00 to 16:00, with a peak of 1900
39 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ around noon in the outer canopy of the trees. After removal of the hanging plates from
40 trees, the PPFD transmittance in full-field was 50% and 18% lower under the plates from the outer
41 layer of the canopies of TK and TCC, respectively, compared with those from the control trees.
42 Under the testers from the intermediate layer of the canopy, PPFD decreased by 39% in TK and
43 12% in TCC compared with the control trees. Finally, under the testers from the inner layer of the
44 canopy, PPFD decreased by 28% and 5% in TK and TCC, respectively, compared with the
45 untreated trees (Tab. 1).
46
47
48
49
50
51
52
53
54
55
56
57

58 In the intermediate canopy layer, the PPFD measured under the plates on the trees was 30% and
59 20% higher in TK and TCC, respectively, compared to the control trees. In the inner canopy layer,
60
61
62
63
64
65

1 the PPFD was four times higher in TK ($394 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$) and three times higher in TCC (287
2 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$) compared with the control trees (Tab. 1). Thus, in the middle and inner canopy layers
3
4 of the treated trees, the PPFD received by the leaves was higher compared with the control trees
5
6 (Tab. 1).
7

8
9 In the outer layers of the canopy, the red-to-blue light ratio was 1:1, with no difference
10
11 between treatment and control trees. In the inner layer, the red-to-blue light ratio decreased by 33%
12
13 in the control trees compared to the treated trees (Fig. 3).
14
15

16 In the outer layer of treated trees, the UV-b radiation was 35% lower compared to the outer
17
18 layer of untreated trees. No substantial differences were found between the middle and inner canopy
19
20 layers, or for these layers between the treated and untreated trees (Glenn et al., 2007).
21
22

23 We observed that the light reflected [whiteness (L^*)] from the leaves coated with reflective
24
25 materials was greater than the light reflected from the control leaves (Fig. 2). Leaf whiteness as a
26
27 result of kaolin and calcium carbonate treatment was 71% and 53% higher, respectively, than that
28
29 of the control trees, according to Glenn et al. (1999). Between the two treatments, whiteness was
30
31 significantly higher in TK compared with TCC (Fig. 2).
32
33
34
35

36 Stomatal opening responds to four independent variables: light, CO_2 , temperature, and
37
38 vapour pressure deficit (V_{pd}) (Farquar and Sharkey, 1982; Draper and Smith, 1966). The stomatal
39
40 conductance (g_s) in the outer layer did not decrease, but was 1.5 and 1.7 times higher in TCC and
41
42 TK, respectively than g_s in TC, as reported in Tab. 1. In the intermediate and inner layers of treated
43
44 trees, g_s was also higher compared to control. Among treatments, g_s was higher in TK than in TCC.
45
46 Furthermore, no significant differences were observed among treatment and control tree with
47
48 respect to soil water potential or plant water status (data are not shown).
49
50
51
52

53 C_i was significantly higher in the outer layers of the control trees compared with TK and
54
55 TCC. In the outer layer, A_n was 50% and 40% higher in TK and TCC, respectively, compared with
56
57 the leaves of control trees ($5.8 \mu\text{mol CO}_2 \text{ m}^{-2}\cdot\text{s}^{-1}$) (Tab. 1). A_n was approximately 60% higher in the
58
59 intermediate canopy layer of the treated trees compared with that in the untreated trees. No
60
61
62
63
64
65

1 differences were observed between the kaolin and calcium carbonate treatments. In the inner
2 canopy layer, A_n was 158% and 383% higher in TCC and TK, respectively, compared with the
3 control trees (Tab. 1).
4

5
6
7 The differences among years were significant for some gas exchange parameters, but no differences
8 were observed about "year x treatment" interaction (Tab.1).
9

10
11 The ratio of R/B light increased in the inner layer compared to the outer layer in TK and TCC
12 treatments, whereas it did not change in the inner layers of the untreated trees (Fig. 3). The SPAD
13 index (Tab. 2) increased by 11% in the leaves of TK and TCC plants compared with the leaves of
14 the control plants. The F_v/F_m values of TK and TCC were 0.67 and 0.66, respectively, whereas the
15 ratio reached a critical value in the control trees, approximately 0.48 (Tab. 2). The treatments in the
16 outer canopy layers also induced a positive effect on Φ_{PSII} (Tab. 2), a fluorescence index that
17 allows determination of the amount of light absorbed by the leaf and used for the photochemical
18 reactions of the light phase of photosynthesis (Correia et al., 2012). The Φ_{PSII} was 2 and 1.8 times
19 higher in TCC and TK, respectively, compared to the control trees. qP , the proportion of PSII
20 reaction centres that are open, was also higher in the treated trees because a portion of the absorbed
21 light was used in the photochemical reactions (Tab. 2). In the control trees, most of the energy
22 absorbed was dissipated, as shown by the coefficient of non-photochemical quenching (qN). The
23 differences among years were significant because these parameters are very sensitive to climate
24 factors. However, no differences were observed about interaction "year per treatment" (Tab. 2).
25
26 Compared with the control trees, in the treated trees the WUE was 77% and 54% higher when
27 treated with kaolin and calcium carbonate, respectively (Fig. 4).
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50

51 The presence of reflective materials on the fruit during development and maturation affected
52 its colour. The CIELab colour space for the skin of the fruit, L^* was significantly lower in TCC
53 trees, a^* was 18% higher in the treated trees compared with the control trees, and b^* was lower (less
54 yellow skin) in TCC trees. The $a^*:b^*$ ratio was also higher in the treated trees compared to the
55 control trees, resulting in a darker orange colour of the skin of the fruit from the treated trees
56
57
58
59
60
61
62
63
64
65

1 compared with the untreated trees. This darker colour is recognised as the preferred colour of
2 orange fruits (Tab. 3).
3

4 The HSL colour space was statistically different between the skin of the fruit of the TCC
5 trees compared with both TK-treated and control trees (Tab. 3). The °hue was significantly lower in
6 the treated trees, with no significant differences between treatment groups. Furthermore, the
7 reflective materials had a favourable effect on the skin colour by reducing its temperature. The pulp
8 colour was darker orange in the TCC trees compared with the control trees (Tab. 4). Indeed a*, b*
9 (CIELab colour space), a*:b* ratio, and °hue (HSL colour space) were higher in TCC fruit
10 compared to TK and control fruits. No difference was observed between TK and control fruits for
11 these parameters (Tab. 4). Furthermore, L* and chrominance were lower in the TK fruit than in the
12 TCC and control fruits (Tab. 4).
13
14
15
16
17
18
19
20
21
22
23
24

25 The total average yield per tree over two years was 15% and 25 % higher in TCC and TK,
26 respectively, compared with TC (Tab. 5). The main carpometric indexes (weight and yield in juice)
27 were similar or better in the treated trees compared with the control trees, as shown in Tab. 6. The
28 fruit weight was significantly different between the treated and untreated trees. The fresh fruit
29 weight was 15% and 25% higher in TCC and TK trees, respectively, than in control trees (Tab. 6).
30 The dry weight was also higher in the fruit of the treated trees compared with the control trees, but
31 the dry weight does not change if expressed as the percentage of fresh weight. The percentage of
32 juice yield was not different between the control trees and the TK trees, whereas it was 14% lower
33 in the TCC trees compared with the control trees (Tab. 6). The total amount of juice per fruit was
34 55.54 and 54.79 g per fruit in the control and TCC fruit, respectively, whereas it was higher in the
35 TK fruit (74.39 g per fruit). Therefore, in the TCC trees, the increased fresh fruit weight compared
36 with the control was attributable to the increased thickness of the peel (data not shown).
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54

55 The SST values of the treated groups were similar to that of the control. Furthermore, the
56 TCC group decreased its titratable acidity, resulting in an increased SST:AT ratio, whereas no
57 differences were observed between the TCC and TK groups (Tab. 6). Nutraceutical parameters
58
59
60
61
62
63
64
65

1 were also not different according to the treatments (Tab. 7). The ascorbic acid content was not
2 significantly different between the treatment and control groups (Tab. 7). The TAC and TPC
3 contents in the treatment groups were not significantly different from those in the control group, and
4 the values were similar to those recorded by Ramful et al., 2011. Therefore, in our experiment, we
5 have observed that the correlation between AA and TAC was stronger than the correlation between
6 TAC and TPC. Indeed, Pearson's coefficient was higher in the first correlation ($r = 0.89$) than in the
7 second correlation ($r = 0.56$). Ascorbic acid is the major antioxidant in orange juice (Rice-Evans
8 and Miller, 1996).

9 The yield efficiency of the TK trees was significantly higher than the control trees, whereas
10 no differences were observed between TCC trees, TK, and the control trees (Tab. 5). Ennab et al.
11 (2017) found that kaolin improved some mandarin fruit quality parameters. All carpometric,
12 organoleptic, nutraceutical, and productive parameters shown significant differences among years.
13 Furthermore, no differences were observed between "Year x Treatment" interaction (Tabb. 3,5,6)
14 except the SST, TA parameters of the fruit (Tab. 5).

36 3. Discussion

37 Photosynthesis is a highly temperature-sensitive process (Yordanov et al., 1986; Berry and
38 Bjorkman, 1980; Critchey, 1998). The A_n depression was recorded in the outer layer of control trees
39 where light, temperature, and LVPD were high (Fig. 1, Tab. 1). Increased temperature increases
40 LAVPD. This, in turn, increases the conductance, but only up to a certain point, beyond which
41 stomatal conductance decreases. Indeed, high temperatures decreased stomatal conductance in the
42 outer layer of the canopy in the untreated trees where LAVPD likely reached the limit threshold (4
43 Kpa). This result agrees with previous observations carried out on citrus and other species (Chen
44 and Zhang, 1994; Pons and Welschen, 2003). In contrast, treatments with reflective materials
45 lowered the LVPD in the outer layer of the canopy, increasing g_s and improved A_n (Tab. 1) in
46 agreement with observation on grapefruit by Jifon and Syvertsen (2003). The lower A_n in the outer
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 layer of untreated trees is attributable to both stomatal and non-stomatal factors (Flexas and
2 Medrano, 2002). Indeed, PPFD exceeded $1400 \mu\text{mol m}^{-2} \cdot \text{s}^{-1}$ in the outer layer of the untreated
3 canopy during measurements of gas exchange. This value is higher than the light saturation point of
4 the leaf, which for citrus is about $600 \mu\text{mol m}^{-2} \cdot \text{s}^{-1}$ (Jifon and Syvertsen, 2003). Salvucci et al.,
5 (2001; 2004) have shown that high temperature lowers the leaf light saturation point.
6
7
8
9
10

11 In addition, excess photons that were not used for the process of photosynthesis caused the
12 production of reactive oxygen species (ROS), which in turn induce photooxidative phenomena
13 (Smirnoff, 1993). Anti-stress protection mechanisms involving the antioxidant system counteract
14 ROS (Brosché and Strid, 2003; Frohnmeyer and Staiger, 2003; Fedina et al., 2007). Furthermore,
15 heat stress induced by high leaf temperature also inhibits the repair of damage caused to PSII
16 (Sharkey, 2005; Allakhverdiev et al., 2008). Initially, ROS inhibits the repair of photodamaged PSII
17 by suppressing the synthesis of proteins such as D1 (Murata et al., 2007), causing photoinhibition
18 primarily by lowering photochemical efficiency (Fv/Fm). The reflective materials we used
19 protected the photochemical process from excess energy as shown by the increasing of Fv/Fm ratio
20 according to Sharkey (2005) and Allakhverdiev et al. (2008).
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35

36 ΦPS2 shows the same pattern as Fv/Fm. This parameter is a more reasonable criterion to
37 define photoinhibition than Fv/Fm because it reflects the decrease of photochemical efficiency
38 induced by non-photochemical dissipation of excessive photons (Jia, 2001). The other fluorometric
39 parameters are photochemical quenching (qP) and non-photochemical quenching (qN), which
40 increase and decrease respectively in the outer layer of treated trees, and also confirm that the
41 reflective materials have limited the negative effects of high PPFD into the outer layer of the
42 canopy (Tab.2).
43
44
45
46
47
48
49
50
51
52

53 In the long run, ROS causes photodestruction of the chlorophyll pigments. The hand-held
54 Chl meter (SPAD) gives accurate estimate of foliar chlorophyll (Torres Netto et al., 2002; Chang
55 and Robison, 2003; Marengo et al., 2009) and is a good tool to diagnose the integrity of the
56 photosynthetic system after exposure of the leaf to high PPFD level (Torres Netto et al., 2005). The
57
58
59
60
61
62
63
64
65

1 SPAD index was lower in the outer layer of the untreated canopy than in trees treated with kaolin
2 and calcium carbonate.
3

4 In the intermediate and inner canopy layers, the temperature was within the optimal range for the
5 leaf. Indeed, no differences were observed with regard to LVPD in the internal layer among the
6 treated and untreated trees. Therefore, the different g_s between the treated and untreated canopy can
7 be exclusively ascribed to better light penetration (Tab. 2), as the effect of reflective action of kaolin
8 and calcium carbonate. The treated trees also had better photosynthetic performances compared to
9 the untreated trees, increasing production as fruit size and dry weight.
10
11
12
13
14
15
16
17

18 The Pearson's coefficient showed a negative correlation among the amount of material
19 deposited on the leaves and PPFD into the canopy (Tab. 1) and a very low correlation among the
20 amount of material deposited on the leaves and gas exchange parameters (Tab. 1).
21
22
23
24
25

26 Therefore, the results obtained could be attributed to the reflective action of the material,
27 rather than its quantity (Tab.1).The spectral distribution of solar radiation changes as the light
28 penetrates and scatters within the tree canopy due to the structure and optical properties of the
29 canopy components, such as the leaves, fruits, and branches (Palmer, 1977; Baldini et al., 1997).
30 The higher levels of red light compared to blue light should promote more formation of flowers in
31 the inner layer of the treated trees compared to the untreated trees. Indeed, the increase in yield of
32 the treated trees was statistically significant when compared with the control trees, similar to what
33 was observed in other species treated with reflective materials (*Olea europaea* L. (Glenn et al.,
34 2001), *Malus communis* L. (Puterka et al., 2000; Wünsche et al., 2004; Glenn and Puterka., 2007),
35 *Prunus dulcis* (Rosati et al., 2006), *Pyrus communis* L. (Chamchaiyaporn et al., 2013), *Mangifera*
36 *indica* L. (Abd-Allah et al., 2013;), and *Punica granatum* L. (Palitha et al., 2010; Hegazi et al.,
37 2014).
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54

55 Climate factors affect fruit quality, as was observed by differences in the colour of rinds
56 among citrus fruits cultivated in tropical and subtropical regions (Young and Erickson, 1961).
57 Orange and mandarin fruits do not attain their attractive rind colour under warm temperatures. The
58
59
60
61
62
63
64
65

1 high temperature interferes with the loss of chlorophyll as well as with the build-up of carotenoids.
2 The reflective action of kaolin and calcium carbonate during the first maturation phase (September)
3
4 lowered the rind temperature, improving peel colour. Regarding the internal quality of the fruit,
5
6 high temperatures in untreated trees led to lower soluble solids, acidity, and juice content compared
7
8 to the lower temperatures of the treated trees, as observed by Burns and Albrigo (1997). Therefore,
9
10 treatments create some improved parameters according to Ennab et al. (2017). Furthermore, Kaolin
11
12 can contrast fruit fly (Bernardes Ourique et al., 2018) and some disease (Melgarejo et al., 2004).
13
14
15

16 All observed, parameters are more or less sensitive to climate factors (Utsunomiya et al.,
17
18 1982), and because these last can be similar among the years but not identical, the differences
19
20 among years were significant for some of them, whereas "year x treatment" interaction was
21
22 observed for only a few parameters.
23
24
25
26
27

28 **4. Conclusion**

29 This study suggests that in warm citrus producing regions, where high radiation and VPD
30
31 can limit photosynthetic capacity, reflective materials such as kaolin and calcium carbonate could
32
33 improve leaf carbon uptake potential in the outer layer of the canopy.
34
35
36

37 These materials also improve light penetration inside the canopy where the light is the very limiting
38
39 factor photosynthetic process, as an effect for shading the layers of upper leaves.
40
41
42

43 The treated tree has improved yield production and fruit quality as size, colour and
44
45 maturation index. Among the two reflective materials used, Kaolin gave the best results compared
46
47 with calcium carbonate for its high light reflective actions. However, for their cost, their use can be
48
49 justified in citrus trees that yield high profits, such as the Navelina ISA 315 used in this experiment.
50
51
52
53

54 **Acknowledgements**

55 The research was financed by MIUR: Project PON03PE_00090_03 "Sustainable models and new
56
57 technologies for the valorisation of Mediterranean vegetable chains".
58
59
60
61
62
63
64
65

References

- Abd-Allah, A.S., Abd El-Razek, E., Saleh, M.M., 2013. Effect of sun-block materials on preventing sunburn injury of Keitt mango fruits. *J. Appl. Sci. Res.* 9 (1), 567-571.
- Allakhverdiev, I.S., Kreslavski, V.D., Klimov, V.V., Los, A.D., Carpentier, R., Mohanty, P., 2008. Heat stress: an overview of molecular responses in photosynthesis. *Photosynth. Res.* 98, 541.
- Balakumar, T., Hani Babu Vincent, V., Paliwal, K., 1993. On the interaction of UV-B radiation (280–315 nm) with water stress in crop plants. *Physiol. Plant.* 87 (2), 217-222.
- Baldini, E., Facini, O., Nerozzi, F., Rossi, F., Rotondi, A., 1997. Leaf characteristics and optical properties of different woody species. *Trees-Struct. Funct.* 12, 73-81.
- Barrett, A.S., Brown, L.R., 2012. A novel method for estimating tree dimensions and calculating canopy volume using digital photography. *Afr. J. Range For. Sci.* 29 (3), 153-156.
- Bernardes Ourique, C., Rodrigues Redaelli, L., Douglas Pedrini, C.E., 2018. Effect of kaolin and limestone on infestation of South American fruit fly in citrus orchards. *Biol. Agric. Hortic.* [https://doi: 10.1080/01448765.2018.1512897](https://doi.org/10.1080/01448765.2018.1512897).
- Berry, J., Bjorkman, O., 1980. Photosynthetic response and adaptation to temperature in higher plants. *Annu. Rev. Plant Physiol.* 31, 491-543.
- Brosché, M., Strid, A., 2003. Molecular events following perception of ultraviolet-B radiation by plants. *Physiol. Plant* 117 (1), 1-10.
- Burns, J.K., Albrigo, L.G. 1997. Granulation in grapefruit. *Proceedings of the Florida State Horticultural Society* 110, 204-208.
- Cen, Y.P., Sage, R.F., 2005. The regulation of Rubisco activity in response to variation in temperature and atmospheric CO₂ partial pressure in sweet potato. *Plant Physiol.* 139, 979–990.

1 Chamchaiyaporn, T., Jutamanee, K., Kasemsap, P., Vaithanomsat, P., Henpitak, C., 2013. Effects of
2 kaolin clay coating on mango leaf gas exchange, fruit yield and quality. *Kasetsart J. Nat. Sci.* 47:
3 479-491.
4
5
6
7 Chang, S.X., Robison, D.J., 2003. Nondestructive and rapid estimation of hardwood foliar nitrogen
8 status using the SPAD502 chlorophyll meter. *For. Ecol. Manag.* 6215, 1–8.
9
10
11 Chen, Z.H., Zhang, L.C., 1994. Diurnal variation in photosynthesis efficiency of leaves in Satsuma
12 Mandarin. *Acta Phytophysiol. Sin.* 20, 263-271.
13
14
15
16 Correia, C., Dinis, L.T., Pinheiro, R., Fraga, H., Ferreira, H., Gonçalves, I., Costa, J., Oliveira, A.,
17 Malheiro, A., Moutinho-Pereira, J., 2012. Climate change and adaptation strategies for viticulture.
18
19 *J. Int. Sci. Publ: Agric. Food* 2, 424-429.
20
21
22
23
24 Crafts-Brandner, S.J., Salvucci, E.M., 2000. Rubisco activase constrains the photosynthetic
25 potential of leaves at high temperature and CO₂. *PNAS* 97 (24), 13430-13435.
26
27
28
29 Critchley, C., 1998. Photoinhibition, in: Raguavendra, A.S. (Eds.), *Photosynthesis: a*
30 *comprehensive treatise.* Cambridge University Press, England, pp. 264-272.
31
32
33
34 Davies, F.S., Albrigo, L.G., 1994. *Citrus*, CAB International, Wallingford.
35
36
37 Draper, N.R., Smith, H., 1966. *Applied regression analysis*, John Wiley and Sons, New York.
38
39
40
41 Ennab, H.A., El-Sayed, S.A., Abo El-Enin, M.M., 2017. Effect of kaolin applications on fruit
42 sunburn, yield and fruit quality of Balady mandarin (*Citrus reticulata*, Blanco). *Menoufia J. Plant*
43 *Prod.* 2, 129-138.
44
45
46
47 Farquhar, G.D., Sharkey, T.D., 1982. Stomatal conductance and photosynthesis. *Annu. Rev. Plant*
48 *Physiol.* 33, 317-345.
49
50
51
52 Fedina, I., Velitchkova, M., Georgieva, K., Demirevska, K., Simova, L., 2007. UV-B response of
53 green and etiolated barley seedlings. *Biol. Plant.* 51, 699–706.
54
55
56
57 Fedina, I., Hidema, J., Velitchkova, M., Georgieva, K., Nedeva, D., 2010. UV-B induced stress
58 responses in three rice cultivars. *Biol. Plant.* 54 (3), 571-574.
59
60
61
62
63
64
65

- 1 Flexas, J., Medrano, H., 2002. Drought-inhibition of photosynthesis in C3 plants: stomatal and
2 non-stomatal limitations revisited. *Ann. Bot.* 89 (2), 183–189.
3
4 Frohnmeyer, H., Staiger, D., 2003. Ultraviolet-B radiation-mediated responses in plants. balancing
5 damage and protection. *Plant Physiol.* 133, 1420-1428.
6
7 Glenn, D.M., Puterka, G.J., Vanderzwet, T., Byers, R.E., Feldhake, C., 1999. Hydrophobic particle
8 films: a new paradigm for suppression of arthropod pests and plant diseases. *J. Econ. Entomol.* 92
9 (4), 759–771.
10
11 Glenn, D.M., Puterka, G.J., Drake, S., Unruh, T.R., Knight, A.L., Baherle, P., Prado, E., Baugher,
12 T., 2001. Particle film application influences apple leaf physiology, fruit yield and fruit quality. *J.*
13 *Amer. Soc. Hort. Sci.* 126, 175-181.
14
15 Glenn, D.M., Puterka, G.J., 2007. The use of plastic films and sprayable reflective particle films to
16 increase light penetration in apple canopies and improve apple color and weight. *HortScience* 42
17 (1), 91-96.
18
19 Glenn, D. M., Cooley, N., Walker, R., Clingeleffer, P., 2010. Impact of Kaolin Particle Film and
20 Water Deficit on Wine Grape Water Use Efficiency and Plant Water Relations. *HortScience* 45 (8),
21 1178–1187.
22
23 Hendry, G.A.F., Houghton, J.D., Brown, S.B., 1987. The degradation of chlorophyll – a biological
24 enigma. – *New Phytol.* 107, 255-302.
25
26 Hegazi, A., Samra, N.R., El-Baz, E.E., Khalil, B.M., Gawish, M.S., 2014. Improving fruit quality of
27 Manfalouty and Wonderful pomegranate by using bagging and some spray treatments with
28 gibberellic acid, calcium chloride and kaolin. *J. Plant Production*, 5 (5), 779 792.
29
30 Hendrey, G.A.F., Houghton, J.D., Brown, S.B., 1987. The degradation of chlorophyll: a biological
31 enigma. *New Phytol.* 107, 255-302.
32
33 Jackson, R.D., Kustas, W.P., Choudhury, B.J., 1988. A reexamination of the crop water stress
34 index. *Irrigation Sci.* 9 (4), 309-317.
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 Jia, H.S., 2001 .Relationship Between Photoinhibition and Active Oxygen Species Metabolism in
2 Apple Leaves Under Drought Stress. Ph.D. Dissertation. Shandong Agricultural University,
3 Shandong.
4
5
6
7 Jifon, J.L., Syvertsen, J.P., 2003. Kaolin particle film application can increase photosynthesis and
8 water use efficiency of “Ruby Red” grapefruit leaves. J. Amer. Soc. Hort. Sci. 128, 107-112.
9
10
11 Kitao, M., Lei, T.T., Koike, T., Tobita, H., Maruyama, Y., 2000. Susceptibility to photoinhibition of
12 three deciduous broadleaf tree species with different successional traits raised under various light
13 regimes. Plant Cell Environ. 23 (1), 81-89.
14
15
16
17 Kriedemann, P.E., 1968. Photosynthesis in vine leaves as a function of light intensity, temperature,
18 and leaf age. Vitis 7, 213-220.
19
20
21
22
23 Larcher, W., 1995. Physiological Plant Ecology, third ed. SpringerVerlag, Berlin.
24
25
26 Lidon, F., Reboredo, F., Silva, O.A.O.M., Duarte, M.P., Ramalho, J., 2012. Impact of UV-B
27 radiation on photosynthesis - an overview. Emir. J. Food Agric. 24 (6), 546-556.
28
29
30
31 Lombardini, L., Glenn, D.M., Harris, M.K., 2004. Application of kaolin-based particle film on
32 pecan trees: consequences on leaf gas exchange, stem water potential, nut quality, and insect
33 populations. HortScience 39, 857-858.
34
35
36
37
38 Long, S.P., Humphries, S., Falkowski, L., 1994. Photoinhibition of photosynthesis in nature. Annu.
39 Rev. Plant Physiol. Plant Mol. Biol. 45, 633-662.
40
41
42
43 Marengo, R.A., 2009. Relationship between specific leaf area, leaf thickness, leaf water content and
44 SPAD 502 readings in six Amazonian tree species. Photosynthetica 47, 184–190
45
46
47
48 Melgarejo, P., Martínez, J.J., Hernández, Fca, Martínez-Font, R., Barrows, P., Erez, A., 2004.
49 Kaolin treatment to reduce pomegranate sunburn. Sci. Hortic. 100, 349-353.
50
51
52
53 Mpoloka, S.W., 2008. Effects of prolonged UV-B exposure in plants. Afr. J. Biotechnol. 7 (25),
54 4874-4883.
55
56
57
58 Murata N., Takahashi S., Nishiyama Y., Allakhverdiev S.I. 2007. Photoinhibition of photosystem II
59 under environmental stress. Biochim. Biophys. Acta 1767, 414–421
60
61
62
63
64
65

- 1 Nahal, I., 1981. The Mediterranean climate from a biological viewpoint, in: Di Castri, F., Goodall,
2 D.W., Specht, R.L. (Eds.), *Ecosystems of the World II: Mediterranean-type shrublands*. Elsevier
3 Scientific Publishing Company, Amsterdam, pp. 63–86.
- 4
5
6
7 Palitha, W.J., Magdalena, M., J. Rogers, J., 2010. The effect of maturity, sunburn and application of
8
9 sunscreen on the internal and external qualities of pomegranate fruit grown in Australia. *Sci. Hort.*
10
11 124, 5761.
- 12
13
14 Palmer, J.W. 1977. Light transmittance by apple leaves and canopies. *J. Appl. Ecol.* 14, 505-513.
- 15
16
17 Pons, T.L, Welschen Rob, A.M., 2003. Midday depression of net photosynthesis in the tropical
18
19 rainforest tree *Eperua grandiflora*: contributions of stomatal and internal conductances, respiration
20
21 and rubisco functioning. *Tree Physiol.* 23, 937–947.
- 22
23
24 Powles, S.B., 1984. Photoinhibition of photosynthesis induced by visible light. *Annu. Rev. Plant*
25
26 *Physiol.* 35, 15–44.
- 27
28
29 Pellegrini, N., Re, R., Yang, M., Rice, Evans C., 1999. Screening of dietary carotenoids and
30
31 carotenoid-rich fruit extracts for antioxidant activities applying the ABTS⁺ radical cation
32
33 decolorization assay. *Methods Enzimol.* 299, 589-603.
- 34
35
36 Potters, G., Pasternak, T.P., Guisez, Y., Jansen, M.A.K., 2009. Different stresses, similar
37
38 morphogenic responses: integrating a plethora of pathways. *Plant Cell Environ.* 32 (2), 158-169.
- 39
40
41 Puterka, G.J., Glenn, D.M., Sekutowski, D.G., Unruh, T.R., Jones, S.K., 2000. Progress toward
42
43 liquid formulations of particle films for insect and disease control in pear. *Environ. Entomol.* 29,
44
45 329-339.
- 46
47
48 Ramful, D., Tarnus, E., Aruoma, O.I., Bourdon, E., Bahorun, T., 2011. Polyphenol composition,
49
50 vitamin C content and antioxidant capacity of Mauritian citrus fruit pulps. *Food Res. Int.* 44, 2088–
51
52 2099.
- 53
54
55 Re, R., Pellegrini, N., Proteggente, A., Pannala, A., Yang, M., Rice Evans, C., 1999. Antioxidant
56
57 activity applying an improved ABTS radical cation decolorization assay. *Free Radic. Biol. Med.* 26,
58
59 1231-1237.
- 60
61
62
63
64
65

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- Rice-Evans, C.A., Miller, N.J., Paganga, G., 1996. Structure antioxidant activity relationships of flavonoids and phenolic acids. *Free Radic. Biol. Med.* 20, 933-956.
- Rosati, A., Metcalf, S.G., Buchner, R.P., Fulton, A.E., Lampinen, B.D., 2006. Physiological effects of kaolin applications in well-irrigated and water-stressed walnut and almond trees. *Ann. Bot.* 98, 267-275.
- Salvucci, M.E., Osteryoung, K.W., Crafts-Brandner, S.J., Vierling, E., 2001. Exceptional sensitivity of Rubiscoactivase to thermal denaturation in vitro and in vivo. *Plant Physiol.* 127, 1053-1064.
- Salvucci, M.E., Crafts-Brandner, J.S., 2004. Inhibition of photosynthesis by heat stress: the activation state of Rubisco as a limiting factor in photosynthesis. *Physiol. Plant.* 120, 179–186.
- Scalzo, J., Politi, A., Pellegrini, N., Mezzetti, B., Battino, M., 2005. Plant genotype affects total antioxidant capacity and phenolic contents in fruit. *Nutrition*, 21 (2), 207-213.
- Sharkey, T.D., 2005. Effects of moderate heat stress on photosynthesis: importance of thylakoid reactions, rubisco deactivation, reactive oxygen species, and thermo tolerance provided by isoprene. *Plant Cell Environ.* 28, 269–277.
- Sinclair, T.R., Allen, L.H. Jr., 1982. Carbon dioxide and water vapor exchange of leaves on field-grown citrus trees. *J. Expt. Bot.* 33, 1166– 1175.
- Slinkard, K., Singleton, V.L., 1997. Total phenol analysis: automation and comparison with manual methods. *Am. J. Enol. Viticult.* 28, 49-55.
- Smirnoff, N., 1993. The role of active oxygen in the response of plants to water deficit and desiccation. *New Phytol.* 125, 27-58.
- Spiegel-Roy, P., Goldschmidt, E.E., 1996. *Biology of Citrus*, Cambridge University Press, Cambridge.
- Syvetsen, J.P., Lloyd, J., 1994. Citrus, in: Schaffer, B., Andersen, P.C. (Eds), *Handbook of Environmental Physiology of Fruit Crops*, Vol. II. CRC Press, Boca Raton, pp. 65-99.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Torres Netto, A., Campostrini, E., Oliveira, J.G., Yamanishi, O.K., 2002. Portable chlorophyll meter for the quantification of photosynthetic pigments, nitrogen and the possible use for assessment of the photochemical process in *Carica papaya* L. *Braz. J. Plant Physiol.* 14, 203–210.

Torres Netto, A., Campostrini, E., Oliveira, J.G., Bressan-Smith, R.E., 2005. Photosynthetic pigments, nitrogen, chlorophyll a fluorescence and SPAD-502 readings in *Coffea* leaves. *Sci. Hort.* 104, 199–209.

Utsunomiya, N., Yamada, H., Kataoka, I., Tomana, T., 1982. The effect of fruit temperature on the maturation of Satsuma mandarin (*Citrus unshiu* Marc.) fruits. *J. Jpn. Soc. Hortic. Sci.* 51, 135–141.

Wünsche, J.N., Lombardini, L., Greer, D.H., 2004. Surround particle film applications effects on whole canopy physiology of apple. *Acta Hort.* 636, 565–571.

Yamamoto, Y., Aminaka, R., Yoshioka, M., Khatoun, M., Komayama, K., Takenaka, D., Yamashita, A., Nijo, N., Inagawa, K., Morita, N., Sasaki, T., Yamamoto, Y., 2008. Quality control of photosystem II: impact of light and heat stresses. *Photosynth. Res.* 98, 589.

Yordanov, I.S., Dilova, R., Petkova, T., Pangelova, V., Goltsev, V., Süß, K.H., 1986. Mechanisms of the temperature damage and acclimation of the photosynthetic apparatus. *Photobiochem. Photobiophys.* 12, 147–155.

Young, L. B., Erickson, L. C., 1961. Influence of temperature on color change in Valencia oranges. *Proc. Am. Soc. Hot. Sc.*, 78, 197-200.

Figure

[Click here to download Figure: Figures...docx](#)

Figure 1 – Pattern of temperature ($^{\circ}\text{C}$) and PPFD ($\text{micromol. m}^2. \text{s}^{-1}$) in the outer layers of the canopy in treated (Kaolin= TK; calcium carbonate =TK) and untreated (Control =TC) sweet orange tree, cv Navelina ISA 315. The measures reported representing the average of the observations made in 3 days of full summer and clear sky during the last decade of July in both years of observation.

Figure 2 –Leaf whiteness measured on the adaxial leaf surfaces, in the outer layers of the canopy in treated (Kaolin= TK; calcium carbonate =TK) and untreated (Control =TC) sweet orange tree, cv Navelina ISA 315. The measures reported representing the average of the observations made in 3 days of full summer and clear sky during the last decade of July in both years of observation. Different letters indicate statistically significant differences per $P < 0.05$; .n= 18

Figure 3 – The Red (680 nm)/Blue (435nm) light ratio in the external (E) and inner (I) layers of the canopy of treated (TK; TP) and untreated (TC) tree. Different letters indicate statistically significant differences per $P \leq 0.05$.

Figure 4 - Water use efficiency (WUE) in the outer layers of the canopy in treated (Kaolin= TK; calcium carbonate =TK) and untreated (Control =TC) sweet orange tree, cv Navelina ISA 315. The measures reported representing the average of the observations made in 3 days of full summer and clear sky during the last decade of July on two years. Different letters indicate statistically significant differences per $P \leq 0.05$.

Figure 5- Correlations between Total Antioxidant Capacity (TAC) and Total Polyphenols (P) and between Total Antioxidant Capacity (TAC) and ascorbic acid, in the citrus fruit of sweet orange, cv Navelina ISA 315




Figure 2




Figure 2




Figure 3




Figure 4





Figure 5

Table 1 - PPFD in three layers of canopy, PPFD under plate in full field and on tree, gas exchange (An, gs, LAVPD,Ci), and Pearson's coefficient in treated (Kaolin= TK; calcium carbonate =TCC) and untreated (Control =TCN) sweet orange tree, cv Navelina ISA 315. The measures reported representing the average of the observations made in 3 days of full summer and clear sky during the last decade of July on two years. n=54

	Layer	Reflective material weight $g\ m^{-2}$	PPFD under plate in full field $\mu mol\ m^{-2}\ s^{-1}$	PPFD measured underplate on the tree $\mu mol\ m^{-2}\ s^{-1}$	An $\mu mol\ CO_2\ m^{-2}\ s^{-1}$	gs $mol\ H_2O\ m^{-2}\ s^{-1}$	LAVAP kPa	Ci $\mu mol\ mol^{-1}$
TCN	1	0	1492±102a	1450±98a	5.3±0.2c	0.14±0.1e	4.16±0.01a	310.30±15a
	2	0	1496±109a	480±38e	5.6±0.3c	0.16±0.08d	2.7±0.02c	240.11±22b
	3	0	1488±105a	102±11h	1.2±0.1e	0.10±0.07f	2.5±0.02d	200.11±32d
TCC	1	3.8±0.09c	933±49c	1180±42b	7.6±0.3b	0.22±0.08b	3.18±0.03b	269.02±32b
	2	1.78±0.08e	1233±37b	605±30e	8.8±0.3a	0.20±0.07c	2.8±0.02c	248.30±25b
	3	0.38±0.08f	1398±11b	286±20g	3.1±0.4d	0.16±0.04d	2.6±0.01d	235.15±10c
TK	1	11.2±0.12a	846±55d	915±46c	8.7±0.4a	0.26±0.03a	3.15±0.02b	247.48±21b
	2	5.5±0.11b	1074±22c	720±33d	9.2±0.3a	0.23±0.04b	2.7±0.01c	250.11±18b
	3	1.12±0.08d	1350±19b	394±21f	5.8±0.2c	0.19±0.03c	2.5±0.01d	230.12±22c
Treatment					*	*	*	*
Year					*	*	*	*
Year x Treatment			n.s.	n.s.	n.s.	n.s.	n.s.	n.s.
Pearson Coefficient TCC weight vs			0.88	0.95	0.215	0.222	0.341	0.125
Pearson Coefficient TK weight vs			0.89	0.96	0.186	0.325	0.284	0.189

Different letters in the column indicate statistically significant differences per * $P\leq 0.05$, ** $P\leq 0.01$ *** $P\leq 0.001$; n.s. = not significant

Table 2 – Effects of kaolin (TK) and calcium carbonate (TCC) sprays on SPAD index and main indices of fluorescence on *Citrus sinensis* L. leaf compared to the control (TC) in the outer layers. The measures reported representing the average of the observations made in 3 days of full summer and clear sky during the last decade of July on two years. (SE= standard error)

	SPAD index	±SE	Fv/Fm	±SE	ΦPSII	±SE	qN	±SE	qP	±SE
TCN	75.445a	2.952	0.48b	0.052	0.102b	0.011	2.07a	0.204	0.298b	0.035
TCC	83.663b	0.813	0.66 a	0.019	0.214a	0.010	1.63b	0.100	0.389a	0.031
TK	83.690b	1.381	0.67a	0.014	0.184a	0.013	1.74b	1.063	0.490a	0.020
Treatment	*		***		***		n.s.		***	
Year	*		**		**		*		**	
Treatment x year	n.s.		n.s.		n.s.		n.s.		n.s.	

Different letters in the column indicate statistically significant differences per * $P \leq 0.05$, ** $P \leq 0.01$, *** $P \leq 0.001$; n.s. = not significant

Table 3 - Effect of treatment on fruit peel and pulp colour characteristics (CIE Lab and HSB colour space) in treated (Kaolin= TK; calcium carbonate =TCC) and untreated (Control =TCN) sweet orange tree, cv Navelina ISA 315.. The measures reported representing the average of the observations made in the last three sampling dates on two years. (SE= standard error)

Treatments	L*	±SE	a*	±SE	b*	±SE	a/b	±SE	Chroma	±SE	°Hue	±SE
TCN (peel)	68.060 a	0.363	17.102 b	1.152	62.247 a	0.486	0.262 b	0.019	66.459 a	0.577	75.229 a	1.295
TCC (peel)	66.710 b	0.358	20.061 a	1.151	59.765 b	0.593	0.327 a	0.021	65.039 b	0.644	72.980 b	1.077
TK (peel)	68.677 a	0.342	19.838 a	1.054	61.271 ab	0.597	0.316 a	0.019	66.072 a	0.628	73.296 b	0.984
<i>Treatment</i>	*		**		**		*		*		**	
<i>Year</i>	*		*		*		*		*		*	
<i>Treatment x Year</i>	n.s.		n.s.		n.s.		n.s.		n.s.		n.s.	
TCN (Pulp)	48.838a	0.134	6.586b	0.177	29.586 ab	0.164						
TCC (Pulp)	48.857a	0.158	7.222a	0.174	30.096 a	0.194						
TK (Pulp)	47.442b	0.166	6.618b	0.200	29.044 b	0.190						
<i>Treatment</i>	*		*		*		*		*		*	
<i>Year</i>	*		*		*		*		*		*	
<i>Treatment x Year</i>	n.s.		n.s.		n.s.		n.s.		n.s.		n.s.	

Different letters in the column indicate statistically significant differences per * $P \leq 0.05$, ** $P \leq 0.01$, *** $P \leq 0.001$; ; n.s. = not significant

Table 4 – Fruit drop pre-harvest (%), fruit at harvest, yield per plant, canopy volume and yield efficiency in treated (Kaolin= TK; calcium carbonate =TCC) and untreated (Control =TCN) sweet orange tree, cv Navelina ISA 315. The measures reported representing the average of the observations made at harvest in two years. (SE= standard error)

Treatment	Fruit drop during summer %	±SE	Fruit number.tree ⁻¹	±SE	Kg.tree ⁻¹	±SE	Volume m ³ .tree ⁻¹	Yield Efficiency Kg.m ⁻³
TCN	4.470a	0.10	406	2.1	139.186c	0.10	24.2	5.78b
TCC	3.530b	0.09	410	3.2	160.375b	0.09	25.1	6.40ab
TK	2.118c	0.08	416	2.4	177.591a	0.08	25.2	7.05a
<i>Year (A)</i>	*		n.s.		n.s.		n.s	n.s.
<i>Treatment (B)</i>	**		n.s.		**			**
<i>Interaction (AXB)</i>	n.s.		n.s.		n.s.			n.s.

Different letters in the column indicate statistically significant differences per * $P \leq 0.05$, ** $P \leq 0.01$; *** $P \leq 0.001$; ; n.s. = not significant

Table 5 - Effect of treatment on fruit weight (FW)t, dry weight (DW), juice yield, total soluble solids (TSS), titratable acidity (TA) and SST/AT in treated (Kaolin=TK; calcium carbonate =TCC) and untreated (Control =TCN) sweet orange tree, cv Navelina ISA 315. The measures reported representing the average of the observations made in the last three sampling dates on two years. (SE= standard error)

Treatment	FW g	±ES	DW g	±ES	DW %	±ES	Juice yield %	±ES	SST	±ES	TA %	±ES	SST/TA	±ES
TCN	145.781c	63.329	26.136b	7.844	15.348ns	0.226	38.315a	1.083	10.465ns	0.278	1.761a	0.157	8.407b	1.116
TCC	166.774b	83.017	32.924a	12.151	16.068	0.558	33.011b	0.894	10.912	0.215	1.354b	0.097	9.692a	0.798
TK	181.711a	78.144	32.927a	9.735	15.105	0.285	41.104a	0.975	10.348	0.145	1.622a	0.079	8.319b	0.822
<i>Year (A)</i>	n.s.		n.s.		n.s.		n.s.		n.s.		***		**	
<i>Treatment (B)</i>	***		***		n.s.		***		n.s.		***		***	
<i>Interaction (AxB)</i>	n.s.		n.s.		n.s.		n.s.		n.s.		*		*	

Different letters in the column indicate statistically significant differences per * $P \leq 0.05$, ** $P \leq 0.01$; *** $P \leq 0.001$; ; n.s. = not significant

Table 6- Total Antioxidant Capacity (TAC), Total Polyphenols (TP), and ascorbic acid o(AA) in orange cv Navelina ISA 315 by the treated (TCC and TK) and untreated tree (TCN). The measures reported representing the average of the observations made in the last three sampling dates on two years. (SE= standard error).

	TAC <i>μmoli Trolox/g FW</i>	±ES	TPC <i>mg GAE/g FW</i>	±ES	Ascorbic acid <i>mg ascorbic acid/100 ml</i>	±ES
<i>TCN</i>	4.863ns	0.208	1.158ns	0.051	66.330ns	2.350
<i>TCC</i>	4.856	0.166	1.125	0.041	68.748	1.851
<i>TK</i>	4.399	0.341	1.093	0.076	62.802	1.804
<i>Year (A)</i>	n.s.		n.s.		n.s.	
<i>Treatment (B)</i>	n.s.		n.s.		**	
<i>Interaction (AxB)</i>	n.s.		n.s.		n.s.	

Different letters in the column indicate statistically significant differences per * $P \leq 0.05$, ** $P \leq 0.01$; *** $P \leq 0.001$; n.s. = not significant

***Credit Author Statement**

Credit Author Statement	Gullo G	Dattola A.	Vonella V.	Zappia R.
Conceptualization	X			
Methodology	X			
Formal Analysis	X	X		
Validation	X	X	X	X
Software	X	X	X	
Writing original draft	X	X		
Writing Review and Editing	x	X		X
Data Curation			X	X
Visualization	X	X		X
Project Administration	X			X
Funding Acquisition				X

Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: