



Università degli Studi Mediterranea di Reggio Calabria
Archivio Istituzionale dei prodotti della ricerca

Impact of invasive alien plants on native plant communities and Natura 2000 habitats: State of the art, gap analysis and perspectives in Italy

This is the peer reviewed version of the following article:

Original

Impact of invasive alien plants on native plant communities and Natura 2000 habitats: State of the art, gap analysis and perspectives in Italy / Lazzaro, Lorenzo; Bolpagni, Rossano; Buffa, Gabriella; Gentili, Rodolfo; Lonati, Michele; Stinca, Adriano; Acosta, Alicia Teresa Rosario; Adorni, Michele; Aleffi, Michele; Allegrezza, Marina; Angiolini, Claudia; Assini, Silvia; Bagella, Simonetta; Bonari, Gianmaria; Bovio, Maurizio; Bracco, Francesco; Brundu, Giuseppe; Caccianiga, Marco; Carnevali, Lucilla; Di Cecco, Valter; Ceschin, Simona; Ciaschetti, Giampiero; Cogoni, Annalena; Foggi, Bruno; Frattaroli, Anna Rita; Genovesi, Piero; Gigante, Daniela; Lucchese, Fernando; Mainetti, Andrea; Mariotti, Mauro; Minissale, Pietro; Paura, Bruno; Pellizzari, Mauro; Perrino, Enrico Vito; Pirone, Gianfranco; Poggio, Laura; Poldini, Livio; Poponessi, Silvia; Prisco, Irene; Prosser, Filippo; Puglisi, Marta; Rosati, Leonardo; Selvaggi, Alberto; Sottovia, Lucio; Spampinato, Giovanni; Stanisci, Angela; Venanzoni, Roberto; Viciani, Daniele; Vidali, Marisa; Villani, Mariacristina; Lastrucci, Lorenzo. - In: JOURNAL OF ENVIRONMENTAL MANAGEMENT. - ISSN 0301-4797. - 274:(2020), p. 111140. [10.1016/j.jenvman.2020.111140]

Availability

DOI: <http://doi.org/10.1016/j.jenvman.2020.111140>

The final published version is available online at:<https://www.sciencedirect>.

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website

Publisher copyright

This item was downloaded from IRIS Università Mediterranea di Reggio Calabria (<https://iris.unirc.it/>) When citing, please refer to the published version.

(Article begins on next page)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48

Affiliations:

- 1 Italian Society for Vegetation Science (SISV), Via Scopoli 22-24, I-27100 Pavia, Italy
- 2 Department of Biology, University of Florence, Via G. La Pira 4, I-50121 Firenze, Italy
- 3 Department of Chemistry, Life Sciences and Environmental Sustainability, University of Parma, Parco Area delle Scienze 11/a, I-43124 Parma, Italy
- 4 Department of Environmental Sciences, Informatics and Statistics, Ca' Foscari University of Venice, Via Torino 155, I-30172 Venezia, Italy
- 5 Department of Earth and Environmental Sciences, University of Milan-Bicocca, Piazza della Scienza 1, I-20126 Milano, Italy
- 6 Department of Agricultural, Forest and Food Sciences, University of Torino, Largo Paolo Braccini 2, I-10095 Grugliasco, Italy
- 7 Department of Environmental, Biological and Pharmaceutical Sciences and Technologies, University of Campania "Luigi Vanvitelli", Via A. Vivaldi 43, I-81100 Caserta, Italy
- 8 Department of Sciences, University of Roma Tre, Viale G. Marconi 446, I-00146 Roma, Italy
- 9 Via degli Alpini 7, I-43037 Lesignano de' Bagni (PR), Italy
- 10 School of Biosciences and Veterinary Medicine, Plant Diversity & Ecosystems Management Unit, Bryology Laboratory & Herbarium, University of Camerino, Via Pontoni 5, I-62032 Camerino (MC), Italy
- 11 Department of Agricultural, Food and Environmental Sciences, Marche Polytechnic University, Via Breccie Bianche, I-60131 Ancona, Italy
- 12 Department of Life Sciences, Via P.A. Mattioli 4, I-53100, Siena, Italy
- 13 Department of Earth and Environmental Sciences, University of Pavia, Via S. Epifanio 14, I-27100 Pavia, Italy
- 14 Department of Chemistry and Pharmacy, University of Sassari, Via Piandanna 4, I-07100 Sassari, Italy
- 15 Faculty of Science and Technology, Free University of Bozen-Bolzano, Piazza Università, 5, I-39100, Bozen-Bolzano, Italy
- 16 Comitato Scientifico, Museo Regionale di Scienze Naturali "Efisio Noussan", Loc. Tache, I-11010 Saint-Pierre (AO), Italy
- 17 Department of Agriculture, University of Sassari, Viale Italia 39, I-07100 Sassari, Italy
- 18 Department of Biosciences, University of Milan, Via Celoria 26, I-20133 Milano, Italy
- 19 Istituto Superiore per la Protezione e la Ricerca Ambientale (ISPRA), Via V. Brancati 60, I-00144 Roma, Italy
- 20 Department of Life, Health & Environmental Sciences, University of L'Aquila, Piazzale Salvatore Tommasi 1, I-67100 L'Aquila, Italy
- 21 Majella National Park, via Badia 28, I-67039 Sulmona (AQ), Italy
- 22 Department of Life and Environmental Sciences, Botany section, University of Cagliari, V.le S. Ignazio 13, I-09123 Cagliari, Italy
- 23 Department of Agricultural, Food and Environmental Sciences, University of Perugia, Borgo XX giugno 74, I-06121 Perugia, Italy
- 24 Department of Earth, Environment and Life Sciences, University of Genova, Corso Europa 26, I-16132 Genova, Italy
- 25 Department of Biological, Geological and Environmental Sciences, University of Catania, Via A. Longo 19, I-95125 Catania
- 26 Department of Agriculture, Environment and Food Sciences, via De Sanctis snc, I-86100 Campobasso, Italy
- 27 Istituto Comprensivo "Bentivoglio", Via Salvo D'Acquisto 5/7, I-44028 Poggio Renatico (FE), Italy

49 28 CIHEAM, Mediterranean Agronomic Institute of Bari, Via Ceglie 9, I-70010 Valenzano (BA), Italy
50 29 Scientific Research and Biodiversity Service, Gran Paradiso National Park, Fraz. Valnontey 44, I-
51 11012 Cogne (AO), Italy
52 30 Department of Life Sciences, University of Trieste, Via L. Gorgieri 5, I-34127 Trieste, Italy
53 31 Department of Chemistry, Biology and Biotechnology, University of Perugia, Polo Didattico, via
54 del Giochetto 6, Ed. A, I-06126 Perugia Italy
55 32 Fondazione Museo Civico di Rovereto, Largo S. Caterina 41, I-38068 Rovereto (TN), Italy
56 33 School of Agricultural, Forestry, Food and Environmental Sciences, University of Basilicata, via
57 dell'Ateneo Lucano 10, I-85100 Potenza, Italy
58 34 Istituto per le Piante da Legno e l'Ambiente, Corso Casale 476, I-10132 Torino, Italy
59 35 Ufficio Biodiversità e Rete Natura 2000, Provincia Autonoma di Trento, Via R. Guardini 75, I-
60 38121 Trento, Italy
61 36 Department of Agriculture, Mediterranean University of Reggio Calabria, loc. Feo di Vito, I-
62 89122 Reggio Calabria, Italy
63 37 Department of Bioscience and Territory, University of Molise, via Duca degli Abruzzi s.n.c., I-
64 86039 Termoli, Italy
65 38 Botanical Garden of Padua, University of Padua, Via Orto Botanico 15, I-35121 Padova, Italy
66 39 University Museum System, Natural History Museum of the University of Florence, Botany, Via
67 G. La Pira 4, I-50121 Florence, Italy

68
69 * Author for correspondence: lorenzo.lazzaro@unifi.it
70
71

72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97

Abstract

Invasive alien plants are a major threat to biodiversity and they contribute to the unfavourable conservation status of habitats of interest to the European Community. In order to favour implementation of European Union Regulation no. 1143/2014 on invasive alien species, the Italian Society of Vegetation Science carried out a large survey led by a task force of 49 contributors with expertise in vegetation across all the Italian administrative regions. The survey summed up the knowledge on impact mechanisms of invasive alien plants in Italy and their outcomes on plant communities and the EU habitats of Community Interest, in accordance with Directive no. 92/43/EEC. The survey covered 241 alien plant species reported as having deleterious ecological impacts. The data collected illustrate the current state of the art, highlight the main gaps in knowledge, and suggest topics to be further investigated. In particular, the survey underlined competition as being the main mechanism of ecological impact on plant communities and Natura 2000 habitats. Of the 241 species, only *Ailanthus altissima* was found to exert an ecological impact on plant communities and Natura 2000 habitats in all Italian regions; while a further 20 species impact up to ten out of the 20 Italian administrative regions. Our data indicate that 84 out of 132 Natura 2000 Habitats (64 %) are subjected to some degree of impact by invasive alien plants. Freshwater habitats and natural and semi-natural grassland formations were impacted by the highest number of alien species, followed by coastal sand dunes and inland dunes, and forests. Although not exhaustive, this research is the first example of nationwide evaluation of the ecological impacts of invasive alien plants on plant communities and Natura 2000 Habitats.

Keywords:

Competition; Ecological impact; Expert survey; Impact mechanism; Impact outcome; Natura 2000 network

98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145

1. Introduction

Biological invasions are one of the most important drivers of biodiversity loss and ecosystem degradation worldwide (Seebens et al., 2017). The establishment and spread of invasive alien species (IAS) have affected multiple ecosystem processes, including community composition, biotic interactions, and functions and services (Vilà and Hulme, 2017). Furthermore, IAS can also impact important socio-economic assets, reducing the efficiency of natural resource exploitation, affecting infrastructure effectiveness, and imposing costly management efforts (Bacher et al., 2017). IAS are indeed one of the major drivers of changes in European habitats, and increase the probability of unfavourable conservation status of natural habitats (Maes, 2013), causing a general deterioration of biodiversity and the alteration of habitat structure and functions in plant communities (Pyšek et al., 2012; Gigante et al., 2018). Therefore, it is an urgent and complex goal in invasion biology to understand the mechanisms underlying biological invasions, one crucial to predicting habitat invasibility (i.e. susceptibility to invasions) and recognising community response to invasion in order to implement actions for the restoration and long-term management of invaded habitats.

Despite there being a general awareness of the effects of invasive alien plants (IAPs), in Europe there is still a lack of exhaustive works investigating the effects of IAPs on native plant communities and on the habitats of Community Interest listed in the Habitats Directive (Council Directive no. 92/43/EEC, hereafter N2000 Habitats) (Guerra et al., 2018). The N2000 network proved to be crucial for preserving the EU's biodiversity, although there are increasing calls for improvements and adjustments (Trochet and Schmeller, 2013; Friedrichs et al., 2018). The importance of the N2000 network in tackling the risks posed by biological invasions was underlined by the European Commission in the *EU 2020 Biodiversity Strategy*, and further emphasised in the recent *EU 2030 Biodiversity Strategy*. However, there is still no common approach for protecting the N2000 network, and its efficacy in decreasing the vulnerability to invasive alien species is in large part still unknown (Guerra et al., 2018; Mazaris and Katsanevakis, 2018). Data on the presence and impact of IAS on the N2000 network is crucial to counter their detrimental impacts, and of pivotal importance when considering the effects of climate change, which are likely to increase the uncertainty associated with IAS performance (Guerra et al., 2018). Indeed, protected areas and the N2000 network could become valuable tools to tackle the spread of invasive species, especially in the light of future climate change (Gallardo et al. 2017).

In Italy, research on IAPs has gained momentum in the last 20 years (Lazzaro et al., 2019). Celesti-Grapow et al. (2009) published the first comprehensive checklist, recently updated by Galasso et al. (2018a). Several national projects and studies have been carried out in the last ten years (e.g. Malavasi et al., 2018; Celesti-Grapow et al., 2016; Lazzaro et al., 2019). Nevertheless, scientific literature concerning the impacts of IAPs in Italy is still patchy and there are no data for some taxa and N2000 habitats.

In this context, the Italian Institute for Environmental Protection and Research (ISPRA) commissioned a project to collect information on the impacts of IAPs on biodiversity and on the N2000 network in Italy. The aim was to support national implementation of Regulation (EU) 1143/2014 on IAS. The study, designed and conducted by the Italian Society of Vegetation Science (SISV), had two main aims: develop of a check-list of alien-dominated plant communities in Italy (Viciani et al., 2020); and assess of the ecological impacts of IAPs on N2000 Habitats in Italy. To achieve this second goal, we carried out a survey of the literature and of expert opinion that involved a large number of botanist members of SISV. Specifically, this survey aimed at i) verifying and listing IAPs known to affect native plant communities and N2000 Habitats on a regional and a national scale, and ii) assessing the mechanism by which species make an impact and the possible

impacts exerted (i.e. impact mechanisms and impact outcomes, respectively, according to Blackburn et al., 2014) and iii) determining whether impact outcomes are exerted on specific N2000 Habitats. A supplementary aim was iv) to verify the presence of specific patterns of invasion in N2000 Habitats, hypothesising that the life form and the time since its first introduction may play a pivotal role in the threat posed by IAPs to Italian N2000 Habitats.

2. Methods

2.1 Definitions and context

We adopted the Regulation (EU) no. 1143/2014 definition of IAS as those “*whose introduction or spread has been found to threaten or adversely impact upon biodiversity and related ecosystem services*”. We focused on Ecological impacts, defined as “*a measurable change to the properties of an ecosystem by an alien species*”, considering only deleterious impacts, meant as “*any impact that changes the environment in such a way as to reduce native biodiversity or alter ecosystem function to the detriment of the incumbent native species*” (Blackburn et al., 2014). We included both natural and semi-natural ecosystems, considering only impacts affecting the native biota and/or ecosystem processes. Conversely, we did not consider either impacts on native species at the individual or population level, or any effect on human society (thus excluding any economic/social and health effects of IAPs).

We adopted the impact scheme of the Global Invasive Species Database (GISD 2020), as described in Blackburn et al. (2014), taking into account all possible impact mechanisms identified by the scheme, and all the ecological impact outcomes at the ecosystem/habitat level (see Table 1).

Table 1. Impact mechanisms and outcomes adopted in the survey of Italian IAPs. Each mechanism listed in the left column may result in one or more of the outcomes listed in the right column (Blackburn et al., 2014).

Impact mechanism	Impact outcomes exerted at the ecosystem/habitat level
Competition	Modification of hydrology/water regulation, purification and quality /soil moisture
Predation	Primary production alteration
Hybridization	Modification of nutrient pool and fluxes
Disease transmission	Modification of natural benthic communities
Parasitism	Modification of food web
Poisoning/Toxicity	Reduction in native biodiversity
Bio-fouling	Unspecified ecosystem modification
Grazing/Herbivory/Browsing	Habitat degradation
Rooting/Digging	Habitat or <i>refugia</i> replacement/loss
Trampling	Physical disturbance
Flammability	Modification of fire regime
Interaction with other invasive species	Modification of successional patterns
Others	Soil or sediment modification: erosion
	Soil or sediment modification: bioaccumulation
	Soil or sediment modification: modification of structure
	Soil or sediment modification: modification of pH, salinity or organic substances
	Other

The nomenclature of IAPs follows Galasso et al. (2018a). As the assessment of the effects of IAPs on habitats of community interest made up a pivotal part of our data collection, to define the habitats we followed the “Italian Interpretation Manual of the 92/43/EEC Habitats Directive” (Biondi et al., 2009) and the Interpretation manual of European Union EU28 (European Commission, 2013).

2.2 Survey strategy

Our evaluation of the current impact outcomes of IAPs in Italy was based on a survey of the literature and expert opinion. The working group was composed of 49 members of the SISV (the co-authors of the present work – mainly technicians or academic botanists, with expertise in vegetation science, N2000 Habitats and IAPs), who provided data and their knowledge of the situation at the local (regional) level or regarding specific IAPs.

We provided each expert with a spreadsheet template that included specific guidelines on the type of data required and how to fill in the spreadsheet (see Appendix 1). The template included an initial list of 184 IAPs established in Italy, taken from the National Alien Plant Species Data Base (Lazzaro et al., 2019). The contributors were asked to provide data on impact mechanisms and outcomes in their region in accordance with Blackburn et al. (2014), see Table 1. Particularly, for each species of the list, experts were asked to provide the following information: a) impact mechanism, b) impact outcomes, c) impact outcomes with specific reference to N2000 Habitats, d) data source, specifying whether it originated from i) scientific literature, ii) technical reports or grey literature or iii) expert assessment, and e) the level of uncertainty of the data provided. In addition, contributors were encouraged to add to the list any further IAPs found to have an impact.

The survey strategy adopted in the study followed the framework of the consensus-building approach (see Vanderhoeven et al., 2017), in which several rounds of structured questionnaires, with subsequent aggregation of responses followed by feedback to the experts, are used to reduce inconsistencies among assessors. In our case, in a first round of evaluation, the contributors were asked to fill in the template individually, after reading the guidelines and the referenced documentation. Data from this preliminary collection (which ended on 2017 July 31) were aggregated and presented to all contributors during a two-day workshop (2017 October 16-17), to discuss possible shortcomings and identify knowledge gaps and dissimilarities in the data collection. After the workshop, we opened a second call (conclusion in 2017 December) to allow all the contributors to homogenize the data provided and overcome the shortcomings that emerged during the workshop.

2.3 Data analyses

2.3.1 Breakdown of results on IAPs and impacts outcomes and mechanisms

The data obtained on the impacts (mechanisms and outcomes) of IAPs on plant communities and N2000 Habitats were merged into a single table. The data were cleaned and standardized, spelling mistakes were corrected and duplicate records were deleted. Data were organized to depict the overall patterns of IAP distribution across Italian regions, and to outline patterns of impact mechanisms and outcomes on native plant communities and N2000 Habitats at species and administrative regions levels.

2.3.2 Patterns of invasion on N2000 Habitats

With the aim of detecting specific trends of invasion, for all IAPs assessed as having an impact on N2000 Habitats at a national level, we collected further data regarding a) the life form, b) the number of administrative regions colonized (according to Galasso et al., 2018a and subsequent updates, Galasso et al., 2018b; 2018c) and c) the date of the first introduction in Italy (information retrieved from literature and technical sources, see Appendix 2). However, although of paramount

219 importance, data on the time of introduction is not entirely reliable, since it may correspond
 220 either to the date of the first introduction in botanical gardens or to the first detection in nature.
 221 Accordingly, we chose to reclassify neophytes into three main groups: 1) introduced between
 222 1492 and 1800, 2) introduced between 1800 and 1950 and 3) introduced from 1950 to date
 223 (2020). This grouping (hereafter named introduction period) reflects the main changes in global
 224 human flows, passing from the age of geographical discoveries to the XIX century (1492–1799),
 225 from colonialism to the industrial revolution and the two world wars (1800–1950), and finally from
 226 the time of the economic boom to globalisation (1951–2020).
 227 We excluded from the analysis all the species introduced before 1492 (archaeophytes) (only eight
 228 species among those exerting impacts on N2000 habitats: *Abutilon theophrasti*, *Arundo donax*,
 229 *Cuscuta cesatiana*, *Cyperus esculentus*, *Cyperus serotinus*, *Isatis tinctoria* subsp. *tinctoria*, *Ricinus*
 230 *communis*, *Sorghum halepense*), and *Salvinia molesta*, whose presence in Italy is doubtful.
 231 We analysed impacts on single N2000 habitats (as indicated by the entire N2000 code, e.g. 1210)
 232 and then on macro-categories, as indicated by the first number of the N2000 code (e.g. 1: coastal
 233 habitats, 2: dune habitats and so on; see European Commission, 2013). To avoid possible biases
 234 due to the uneven number of habitats across administrative regions, the analyses were conducted
 235 at the national level (i.e. we used number of habitats and habitat macro-categories invaded by the
 236 species in Italy).
 237 To investigate the correlation between the number of habitats, the number of macro-categories of
 238 habitats and the number of colonized administrative regions, we ran a correlation analysis for
 239 each introduction period, we calculated the pairwise Spearman's rank correlation coefficient
 240 (Spearman's ρ) and evaluated its significance by means of the asymptotic t approximation.
 241 Finally, we ran a series of generalized linear models (GLM) to study the effect of the introduction
 242 period and of the life form categories on the number of invaded N2000 habitats, the number of
 243 invaded N2000 habitat macro-categories, and the number of invaded administrative regions.
 244 Given the overdispersion of our data, we adopted a quasi-Poisson distribution and evaluated the
 245 significance of the terms with an ANOVA table. All the analyses were conducted in R environment
 246 vers. 3.6.1 (R Core Team 2019).

247 3. Results

248 3.1 Breakdown of results on IAPs and impacts outcomes and mechanisms

249 We collected data on 241 IAPs, 57 more than the 184 originally indicated in the template (see
 250 Appendix 2). Only a few species were reported in a high number of administrative regions and by
 251 several contributors. In general, degree of knowledge varied substantially between regions, as
 252 shown by the variation in numbers of regional records on impact mechanisms and outcomes (Figs.
 253 1A and 1B). The number of IAPs assessed as having an ecological impact (Fig. 1C) and the
 254 distribution of impacts on N2000 habitats (Fig. 1D), varied widely between administrative regions.
 255 Similarly, compared to the total number of habitats harboured in a given region, the percentage of
 256 those exposed to some degree of ecological impact was very variable, with Lombardy and Friuli-
 257 Venezia Giulia having more than half of the habitats impacted by IAPs, followed by Molise and
 258 Sardinia (Fig. 1E). Variability also characterized the relationship between species and their impact
 259 mechanisms and outcomes. Indeed, most IAPs had very few reports of mechanisms of impact—220
 260 out of 241 species had less than five reports of impacts—while only a handful of species had a high
 261 number of records. A similar situation was found as far as impact outcomes are concerned: most
 262 of the species had very few reports and very few species had a high number of records (see Fig. 2).
 263 Of the impact mechanisms, “*Competition*” was the most frequent, being common to around 83%
 264 of reports, followed by “*Unknown*” mechanism (4%), “*Interaction with other invasive species*” (4%)
 265 and “*Poisoning/toxicity*” (3%).

266 *Ailanthus altissima*, listed in all Italian administrative regions, was the species with the highest
267 number of records of impact mechanisms. Competition was the main impact mechanism assigned
268 to the species, followed by “Rooting/digging”. Further species with a very high number of records,
269 such as *Senecio inaequidens*, *Robinia pseudoacacia*, *Helianthus tuberosus* and *Sorghum halepense*,
270 were all assessed as invaders with a high degree of impact in many administrative regions (see Fig
271 3A for main mechanisms of the first 23 species). Nevertheless, despite a high number of impact
272 reports, most data—62%—came from expert assessments that were not experimentally verified
273 (Fig. 4 A). Only 25% of the reports were retrieved from the scientific literature and 13% from
274 technical reports and grey literature.

275 Species most frequently recorded for impact outcomes differed from those most frequently
276 recorded for impact mechanisms: *Robinia pseudoacacia* was the species with the highest number
277 of impact outcome records, followed by *Acacia saligna*, *Amorpha fruticosa*, *Arundo donax*,
278 *Ailanthus altissima*, *Carpobrotus edulis*, *C. acinaciformis*, *Helianthus tuberosus*, *Senecio*
279 *inaequidens*, and *Solidago gigantea* (Fig 3B). As with impact mechanisms, most of the records
280 (78%) were from expert assessments, while 13% were retrieved from technical reports and grey
281 literature, and only 9% from the scientific literature (Fig. 4B).

282 Reduction in native biodiversity was by far the most reported outcome, followed by general
283 habitat degradation, loss of habitat and refugia, and modification of successional patterns (Fig.
284 5A). The ranking of threats posed to N2000 Habitats showed the same order. Indeed, a reduction
285 in native biodiversity was cited for nearly all N2000 habitats present in Italy (81 out of 84),
286 followed by the same outcomes named above (Fig. 5B).

287 Nonetheless, impacts were unevenly distributed, especially in terms of the number of IAPs
288 impacting specific habitats, and less in terms of the number of administrative regions in which the
289 target N2000 habitat is impacted (Fig. 6A-B). N2000 Habitat 3270 was by far the one impacted by
290 the highest number of invasive species (79 species), followed by N2000 Habitat 6430. At the
291 macro-category level, freshwater habitats (N2000 Habitats 3xxx) and natural and semi-natural
292 grassland formations (N2000 Habitats 6xxx) were impacted by the highest number of alien species,
293 followed by coastal sand dunes and inland dunes (N2000 Habitats 2xxx), and forests (N2000
294 Habitats 9xxx). In terms of the regional distribution of impacted habitats, freshwater habitats were
295 generally affected in many regions, with coastal sand dunes, coastal and halophytic habitats
296 (N2000 Habitats 1xxx) and forests also being frequently affected (Fig 6 B).

297 **3.2 Patterns of invasion on N2000 Habitats**

298 The 241 IAPs recorded in our survey included 167 neophytes invading the N2000 habitats; 29
299 species were introduced before 1800, 84 between 1800 and 1950, and 54 after 1950 (Appendix 2).
300 Therophytes (56 species) and phanerophytes (45) were the most frequent life forms, followed by
301 hemicryptophytes (19), geophytes (17), chamaephytes (16), and hydrophytes (14). *Ailanthus*
302 *altissima*, *Robinia pseudoacacia*, *Senecio inaequidens*, *Amorpha fruticosa*, and *Carpobrotus edulis*
303 were the most frequent invaders both in N2000 habitats (28, 25, 23, 17, 16 each respectively) and
304 in habitat macro-categories (7, 7, 8, 6, 5 respectively), although with slightly different rankings
305 (see Appendix 2). The number of habitats and that of habitat macro-categories invaded correlated
306 strongly for all three periods (p value < 0.001 , see Table 2), although the correlation decreased
307 slightly from the first introduction period (before 1800; $p = 0.930$) to the last (from 1950 to date; p
308 $= 0.845$). Showing a more strongly decreasing trend, the number of regions colonised was
309 significantly correlated with the number of habitats ($p = 0.523$, p value $= 0.003$) and macro-
310 habitats invaded ($p = 0.489$, p value $= 0.007$) only for the first introduction period (before 1800);
311 while no significant correlation was found for the other two introduction periods (Table 2).
312 Life form categories significantly affected the distribution of species in terms of number of invaded
313 N2000 habitats and macro-habitats, as well as in terms of number of colonised administrative

regions (Table 3). Overall, chamaephytes were the most widespread invaders, invading the highest number of habitats (and habitat macro-categories), followed by geophytes, phanerophytes and therophytes; hydrophytes were specific to a small number of habitats (Figs. 7A, C). On the other hand, therophytes had spread into the highest number of regions, together with geophytes (Fig. 7E). Introduction period strongly affected the number of habitats, of habitat macro-categories, and of administrative regions invaded (Table 3). Indeed, the longer a species had been introduced, the higher the number of invaded habitats, macro-habitats and administrative regions (7 B, D, F).

Table 2 Matrices of correlation between the number of N2000 habitats (Habitats), number of macro-categories of N2000 habitats (Macro-habitats) and number of invaded administrative regions (Regions), for each introduction period. In each correlation matrix, the upper triangle (numbers in plain text) displays Spearman's rank correlation coefficient (Spearman's rho), while the lower triangle (numbers in italic) displays its significance.

Introduction Period		Habitats	Macro-habitats	Regions
1492 – 1799	Habitats	-	0.930	0.523
	Macro-habitats	<0.001	-	0.489
	Regions	0.003	0.007	-
1800 – 1950	Habitats	-	0.860	0.129
	Macro-habitats	<0.001	-	0.187
	Regions	0.129	0.187	-
1951 – present	Habitats	-	0.845	0.077
	Macro-habitats	<0.001	-	0.096
	Regions	0.579	0.488	-

Table 3 Analysis of the deviance table for the generalized linear models analysing the effect of life form categories and introduction period on the number of invaded N2000 Habitats (Habitats), number of macro-categories of N2000 Habitat (Macro-habitats) and number of invaded administrative regions (Regions). χ^2 = Likelihood ratio Chi-square; Df = Degree of freedom; Significance codes: P value < 0.001 '***'; P value < 0.05 '*'.

Response	Term	χ^2	Df	P value	
Habitats	life form	24.576	5	<0.001	***
	introduction period	28.551	2	<0.001	***
	life form×introduction period	12.005	10	0.285	
Macro-habitats	life form	23.720	5	<0.001	***
	introduction period	28.723	2	<0.001	***
	life form×introduction period	15.669	10	0.109	
Regions	life form	13.725	5	0.017	*

introduction period	52.092	2	<0.001	***
life form×introduc tion period	15.691	10	0.109	

4. Discussion

Our data showed that the general impact of IAPs on native plant communities and N2000 Habitats has still only been partially unravelled at the national level. Specifically, our study brought to light two main problems. One is that that very few data are available on the mechanisms by which IAPs exert their impact. This lack greatly reduces our ability to implement effective adaptive strategies to counteract the spread and the effects of IAPs. In addition to this, the data that are available are very unevenly distributed between regions, further reducing our capacity to understand the nation-wide effects of IAPs. Filling these gaps calls for an urgent nation-wide collaborative initiative with coordinated action programs and standard methodologies. The initiative would ideally be conducted under the auspices of ISPRA or of the Italian Ministry of the Environment and Preservation of Land and Sea, which provided funding and motivation for the present study. Also, the establishment of a national collaboration between numerous Italian research groups to participate in joint projects at the EU level (i.e. within EU LIFE programme or Horizon Europe 2021-2027), or within national scientific societies like SISV or the Italian Botanical Society (SBI), is essential to complete the picture of alien species invasion in Italy.

The differences between the impacts recorded for different Italian regions are consistent with the findings of the main catalogues of alien plants in Italy (Galasso et al., 2018a; Celesti-Grapow et al., 2009). The highest number of records was observed in the largest and most densely populated regions (i.e. Lombardy, Piedmont, and Tuscany), where human-driven land cover changes like urbanization, industrialisation, road infrastructures, and agriculture, cause higher rates of introduction (McLean et al., 2017), thereby facilitating biological invasion. The intensification of agricultural use of land has been proven to play a crucial role in the introduction, establishment and spread of IAPs due to a decline in biodiversity caused by oversimplified landscape matrices (Walker et al., 2009; Buffa et al., 2018). This is especially true for lowland riverscapes which suffer from the deterioration of water, of sediments and of hydrological regimes (Bolpagni and Piotti, 2015; Bolpagni et al., 2013), situations that have been identified as common key factors driving the establishment and spread of IAPs in newly invaded areas (Aronson et al., 2017).

However, comparatively large and densely populated regions (e.g. Lazio, Emilia-Romagna) did not show the same degree of invasion. This difference could be the result of contrasting levels of awareness between the different administrative regions in Italy. A minority of regions have already adopted specific regulatory frameworks to address the issue of biological invasions (Brundu et al., 2020), while the other regions still pay little attention to this matter. Only Lombardy, Piedmont, Aosta Valley have a list of restricted IAPs, approved by regional laws, and working groups dedicated to IAPs. In Friuli-Venezia Giulia and Tuscany only few IAPs are considered in regional laws. Finally, Liguria has established a surveillance network and a permanent working group on IAS within the Italian-French ALIEM Project. To our knowledge, all the other Italian administrative regions lack a local regulatory framework on IAPs, even if the recent promulgation of legislative decree no. 230/2017 calls for a comprehensive framework to tackle this issue. We are convinced that a decisive contribution to this matter would come from the establishment of a national list of invasive alien species of Member State (Italy) concern (see Art. 12 of Regulation (EU) no. 1143/2014), under the guidance of the Italian Ministry of the Environment and Preservation of Land and Sea. To this end, an important action aimed at the

374 individuation of candidate species for the implementation of such a list (see Lazzaro et al., 2019) is
375 currently being carried out by ISPRA, within the Life ASAP project (LIFE15 GIE/IT/001039).
376 The high number of expert-based assessments in our survey underscores the major difficulty in
377 retrieving suitable and reliable literature on the impacts of IAPs in Italy. Indeed, direct evidence is
378 frequently also lacking for very well-studied species, generally considered a priori to be a serious
379 threat to biodiversity. The gaps in knowledge of the different taxa hinder the study (and
380 management) of the impacts associated with biological invasions, with most papers focusing on a
381 narrow set of already studied species (Hulme et al., 2013; Latombe et al., 2017). This is especially
382 serious because data on impacts are necessary to lay the basis for any generalisation about
383 biological invasions and are mandatory for risk assessment and management (Bolpagni et al.,
384 2014a; Lazzaro et al., 2015).

385 For example, information concerning *A. altissima* mainly comes from regional reports and
386 checklists which often lack direct measurements of the cited impacts (see Badalamenti et al.,
387 2016; Maiorca et al., 2007). Although the impacts caused by this species are relatively well-studied
388 (Castro-Díez et al., 2019), impacts in Italy are only documented for Sardinia (Traveset et al., 2008;
389 Vilà et al., 2006) and the Karst area in northeastern Italy (Uboni et al., 2019) and there is very little
390 literature on impacts outcomes on plant communities or on N2000 Habitats. Among the most
391 studied species, *Robinia pseudoacacia* stands out as a major invasive tree in Europe (Vítková et al.,
392 2017). Many studies in Italy have focused on the impacts of this species (Nascimbene et al., 2012,
393 2015; Benesperi et al., 2012; Lazzaro et al., 2018; Sitzia et al., 2018; Campagnaro et al., 2018;
394 Gentili et al., 2019). Most authors have found evidence that the rapid expansion of this species in
395 Italy is causing the progressive decline of native forests, with loss of species richness and diversity
396 and a shift in species composition towards nitrophilous assemblages (Benesperi et al., 2012;
397 Lazzaro et al., 2018; Allegrezza et al., 2019). In contrast, other authors have shown that secondary
398 *Robinia* forests, growing on abandoned lands, may host compositionally heterogeneous plant
399 communities and may contribute to some degree to regional biodiversity (Campagnaro et al.,
400 2018). Nevertheless, as pointed out also by Lazzaro et al. (2018), it is worth mentioning that in
401 many cases *Robinia* forests replace habitats considered of community interest in Europe (i.e.
402 N2000 Habitats 9260, 91B0, 91M0, 91AA*; Montecchiari et al., 2020, and 92A0 among others).
403 *Robinia pseudoacacia* is also an important forest species, so that one option would be to apply
404 forestry best practice to avoid its escape from areas set aside for cultivation.

405 *Robinia pseudoacacia* is also predicted to be one of the most competitive species in a climate
406 change scenario (Kleinbauer et al., 2010), and recently Nascimbene et al. (2020) showed the
407 effects of the interaction between climate change and invasion by *R. pseudoacacia* on the
408 endangered lichen species *Lobaria pulmonaria*. Biological invasions and climate change (often
409 referred to as “double trouble”) are considered two of the key drivers of biodiversity loss, whose
410 interaction will lead to a magnification of the threats to biodiversity worldwide (Mainka and
411 Howard, 2010).

412 Among the highly invasive tree species, in Northern Italy *Prunus serotina* and *Quercus rubra* have
413 also been reported to greatly impact native communities and ecosystem components at the soil
414 level (Gentili et al., 2019; Vegini et al., 2020). *Acacia* is another genus well-studied in Italy and
415 worldwide. The negative impacts on plant communities of these nitrogen-fixing trees are well
416 documented for several species (e.g., *A. dealbata*: Lazzaro et al., 2014; Minuto et al., 2020; *A.*
417 *pycnantha*: Lazzaro et al., 2015). Likewise, the impacts of *A. saligna* on coastal dune N2000
418 habitats (Del Vecchio et al., 2013; Bonari et al., 2017; Calabrese et al., 2017).

419 *Carpobrotus acinaciformis*, *C. edulis* and their hybrids (Campoy et al., 2018), is another group of
420 invasive species widely investigated in Italy, whose impacts on both biodiversity (Santoro et al.,
421 2012; Jucker et al., 2013) and soil conditions (Zedda et al., 2010; Santoro et al., 2011; Badalamenti

et al., 2016) are well depicted. Their pattern of occurrence at the community level (Carboni et al., 2010; Sperandii et al., 2017) as well as their habitat preference, including N2000 Habitats 2120, 2210 and 2250*, have also been studied (Sarmati et al., 2019).

Some specific studies have focused on the impact of different IAPs (e.g. *Ambrosia psilostachya*, *Cenchrus longispinus*, *Erigeron canadensis*, *Oenothera stueckii*, *Senecio inaequidens*) on sand dune ecosystems in northeastern Italy, showing significant negative effects on species richness, species diversity and evenness, and plant community composition, with effects increasing from N2000 drift line habitats (1210) to fixed-dune habitats (2130) (Del Vecchio et al., 2015).

Unfortunately, once these few well-studied IAPs are excluded, most of the records of impact on plant communities and N2000 habitats collected in the present general assessment derive from expert evaluations. This applies in particular to the species listed among the Invasive Alien plant Species of Union Concern (*sensu* Regulation (EU) no. 1143/2014), both because their spatial distribution in Italy is still scattered or localized (e.g. *Alternanthera philoxeroides*, *Pontederia crassipes*) and especially because data on their impacts is still missing (with only a few exceptions; see e.g. Lastrucci et al. (2017) for *Myriophyllum aquaticum*). The lack of information may also be due to the difficulty in obtaining reliable data for plants and vegetation in aquatic ecosystems, normally extremely time- and money-consuming to sample (Azzella et al., 2017). Indeed, half of the IAPs of Union concern are aquatic or wetland plants (19 out of 36). Despite their high number in the EU list confirming the general poor state of conservation of inland waters (Brundu, 2014; Lastrucci et al., 2017), aquatic IAPs have been so far neglected or little investigated in Italy.

However, several studies confirmed their pivotal role in reducing local biodiversity. Bolpagni (2013a; 2013b) and Bolpagni et al. (2017) found that *Lagarosiphon major* and *Elodea nuttallii* create extensive submerged meadows that almost completely replace native macrophyte communities belonging to N2000 habitats 3140 and 3150. *Nelumbo nucifera* and *Ludwigia hexapetala* seem to actively compete with native species (Bolpagni et al., 2014b; Villa et al., 2017; 2018; Tóth et al., 2019) due to their enhanced competitive ability for limiting resources and their tolerance to edaphic conditions variability (Tóth et al., 2019). Some specific studies focused on the impact of *Lemna minuta* on freshwater ecosystems. The results highlighted substantial negative effects on water quality and on aquatic plant and animal communities (Ceschin et al., 2019; 2020), showing that *L. minuta* causes drastic alterations to the local vegetation, often replacing native species. Similar results have been found for the most common duckweed, *L. minor*, which assimilates available nutrients faster than native species and shows a higher relative growth rate (Ceschin et al., 2016a, 2016b), characteristics that make this species highly competitive. However, the scarcity of knowledge on the majority of aquatic IAPs hinders both a correct assessment of the environmental impacts and the planning of actions to be carried out for effective recovery of impacted ecosystems.

Our survey also evidenced a group of well-known and widespread invasive herbs (i.e., *Amaranthus retroflexus*, *Ambrosia artemisiifolia*, *Artemisia verlotiorum*, *Bidens frondosa*, *Erigeron canadensis*), that are very competitive in disturbed habitats (agricultural areas, roadsides, ruderal areas) also thanks to their high propagule pressure. Ruderal species are highly opportunistic and it is worth mentioning the presence of *Phytolacca americana* and *Solanum chenopodioides* in an old-growth *Quercus ilex* urban forest disturbed by a severe windstorm in Southern Italy (Bonanomi et al., 2018). Opportunistic alien species, which invest in rapid growth and in sexual reproduction, are a challenging issue in invasion biology since their invasiveness may change during different successional stages (Domènech and Vilà, 2006). For instance, the annual *A. artemisiifolia* is completely suppressed by late colonists and perennial species after a three-year succession (Gentili et al., 2017). Conversely, the invasion of *Solidago canadensis* has been reported to modify

the trajectory of vegetation succession and to exert a higher negative effect on native diversity in older successional communities (Fenesi et al., 2015a; 2015b).

Finally, this survey showed that the time elapsed since introduction has a highly significant effect on the number of habitats invaded. This highlights the importance of better investigating species of relatively recent introduction, which are not yet truly invasive, but which have considerable potential to invade given the size of the introduced populations. These include some species of *Eucalyptus*, especially *E. camaldulensis*, which in recent years has begun to spread in river habitats in Sicily (Badalamenti et al., 2018). In this, as in other cases, biological evolution acts on the introduced species and may enable some alien plants to occupy a broad range of novel habitats until they become invasive (Oduor et al., 2016). Tree species should be carefully monitored because although they take time to become invasive, when they do, the impact is high because of their large biomass.

5. Conclusions

Our survey is the first attempt to assess the impact of the most harmful IAPs on plant communities and on N2000 habitats throughout Italy. The study highlighted numerous knowledge gaps which, however, replicate the gaps in plant invasion science ascertained at a global level. The differences in levels of knowledge between the Italian administrative regions is likely linked to dissimilarities in awareness, as shown by regional differences in policies and legislation. Thus, a primary aim of the scientific community should be to even up disparities in knowledge at the regional level. Most importantly, we highlighted a glaring lack of evidences, even for well-known invasive species. We also exposed a lack of data on the impacts of IAPs on N2000 habitats, which ought to be a primary focus of conservation efforts. Thus, filling the knowledge gap is a mission of primary importance, to provide data both for risk analysis and to support decision makers. Our results show that Italy needs a coordinated nation-wide strategy to evaluate and manage the risk of invasion in the N2000 network. This need is even more urgent in the light of the compounded effects of biological invasion and climate change, which biodiversity management planning and policy should take into careful consideration.

Funding

This work was carried out as part of the project “Implementazione del regolamento 1143/2014 sulle specie esotiche” funded by the Italian Ministry of the Environment and Preservation of Land and Sea.

Author contributions

L.Las., R.Bol., L.Laz., G.Buf. and F.Bra. were involved in designing the methodology and coordinated the work for the data collection on behalf of Italian Society of Vegetation Science.

L.Laz., L.Las., R.Bol., G.Buf., R.Gen., M.Lon. and A.Sti. were involved in data analysis and interpretation and drafting the manuscript. All authors provided data for the survey as associated of SISV, critically revised the draft of the manuscript and gave final approval for publication.

507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554

References

Allegrezza, M., Montecchiari, S., Ottaviani, C., Pelliccia, V., Tesei, G., 2019. Syntaxonomy of the *Robinia pseudoacacia* communities in the central peri-adriatic sector of the Italian peninsula. *Plant Biosystems - An International Journal Dealing with all Aspects of Plant Biology* 153, 616–623. <https://doi.org/10.1080/11263504.2019.1610108>

Aronson, M.F.J., Patel, M.V., O'Neill, K.M., Ehrenfeld, J.G., 2017. Urban riparian systems function as corridors for both native and invasive plant species. *Biological Invasions* 19, 3645–3657. <https://doi.org/10.1007/s10530-017-1583-1>

Azzella, M.M., Bresciani, M., Nizzoli, D., Bolpagni, R., 2017. Aquatic vegetation in deep lakes: macrophyte co-occurrence patterns and environmental determinants. *Journal of Limnology* 76(s1), 97–108. <https://doi.org/10.4081/jlimnol.2017.1687>

Bacher, S., Blackburn, T.M., Essl, F., Genovesi, P., Heikkilä, J., Jeschke, J.M., Jones, G., Keller, R., Kenis, M., Kueffer, C., Martinou, A.F., Nentwig, W., Pergl, J., Pyšek, P., Rabitsch, W., Richardson, D.M., Roy, H.E., Saul, W.-C., Scalera, R., Vilà, M., Wilson, J.R.U., Kumschick, S., 2017. Socio-economic impact classification of alien taxa (SEICAT). *Methods in Ecology and Evolution* 9, 159–168. <https://doi.org/10.1111/2041-210x.12844>

Badalamenti, E., Cusimano, D., Mantia, T.L., Pasta, S., Romano, S., Troia, A., Ilardi, V., 2018. The ongoing naturalisation of *Eucalyptus* spp. in the Mediterranean Basin: new threats to native species and habitats. *Australian Forestry* 81, 239–249. <https://doi.org/10.1080/00049158.2018.1533512>

Badalamenti, E., Gristina, L., Laudicina, V.A., Novara, A., Pasta, S., Mantia, T.L., 2016. The impact of *Carpobrotus* cfr. *acinaciformis* (L.) L. Bolus on soil nutrients microbial communities structure and native plant communities in Mediterranean ecosystems. *Plant and Soil* 409, 19–34. <https://doi.org/10.1007/s11104-016-2924-z>

Benesperi, R., Giuliani, C., Zanetti, S., Gennai, M., Lippi, M.M., Guidi, T., Nascimbene, J., Foggi, B., 2012. Forest plant diversity is threatened by *Robinia pseudoacacia* (black-locust) invasion. *Biodiversity and Conservation* 21, 3555–3568. <https://doi.org/10.1007/s10531-012-0380-5>

Biondi, E., Blasi, C., Burrascano, S., Casavecchia, S., Copiz, R., Vico, E.D., Galdenzi, D., Gigante, D., Lasen, C., Spampinato, G., Venanzoni, R., Zivkovic, L., 2009. Italian Interpretation Manual of the 92/43/EEC Directive habitats. Accessed March 24, 2020. URL: <http://vnr.unipg.it/habitat/>

Blackburn, T.M., Essl, F., Evans, T., Hulme, P.E., Jeschke, J.M., Kühn, I., Kumschick, S., Marková, Z., Mrugała, A., Nentwig, W., Pergl, J., Pyšek, P., Rabitsch, W., Ricciardi, A., Richardson, D.M., Sendek, A., Vilà, M., Wilson, J.R.U., Winter, M., Genovesi, P., Bacher, S., 2014. A Unified Classification of Alien Species Based on the Magnitude of their Environmental Impacts. *PLoS Biology* 12, e1001850. <https://doi.org/10.1371/journal.pbio.1001850>

Bolpagni, R., 2013a. Multimetric indices based on vegetation data for assessing ecological and hydromorphological quality of a man-regulated lake. *Annali di Botanica* 3, 87–95. <https://doi.org/10.4462/annbotrm-10236>

Bolpagni, R., 2013b. Macrophyte richness and aquatic vegetation complexity of the lake Idro (Northern Italy). *Annali di Botanica* 3, 35–43. <https://doi.org/10.4462/annbotrm-10207>

Bolpagni, R., Azzella, M.M., Agostinelli, C., Beghi, A., Bettoni, E., Brusa, G., De Molli, C., Formenti, R., Galimberti, F., Cerabolini, B.E.L., 2017. Integrating the water framework directive into the habitats directive: Analysis of distribution patterns of lacustrine EU habitats in lakes of Lombardy (Northern Italy). *Journal of Limnology* 76, 75–83. <https://doi.org/10.4081/jlimnol.2017.1627>

Bolpagni, R., Bartoli, M., Viaroli, P., 2013. Species and functional plant diversity in a heavily impacted riverscape: Implications for threatened hydro-hygrophilous flora conservation. *Limnologica* 43, 230–238. <https://doi.org/10.1016/j.limno.2012.11.001>

555 Bolpagni, R., Bresciani, M., Laini, A., Pinardi, M., Matta, E., Ampe, E.M., Giardino, C., Viaroli, P.,
 556 Bartoli, M., 2014b. Remote sensing of phytoplankton-macrophyte coexistence in shallow
 557 hypereutrophic fluvial lakes. *Hydrobiologia* 737, 67–76. [https://doi.org/10.1007/s10750-013-1800-](https://doi.org/10.1007/s10750-013-1800-6)
 558 [6](https://doi.org/10.1007/s10750-013-1800-6)
 559 Bolpagni, R., Laini, A., Soana, E., Tomaselli, M., Nascimbene, J., 2014a. Growth performance of
 560 *Vallisneria spiralis* under oligotrophic conditions supports its potential invasiveness in mid-
 561 elevation freshwaters. *Weed Research* 55, 185–194. <https://doi.org/10.1111/wre.12128>
 562 Bolpagni, R., Piotti, A., 2015. The importance of being natural in a human-altered riverscape: role
 563 of wetland type in supporting habitat heterogeneity and the functional diversity of vegetation.
 564 *Aquatic Conservation: Marine and Freshwater Ecosystems* 26, 1168–1183.
 565 <https://doi.org/10.1002/aqc.2604>
 566 Bonanomi, G., Incerti, G., El-Gawad, A.M.A., Sarker, T.C., Stinca, A., Motti, R., Cesarano, G.,
 567 Teobaldelli, M., Saulino, L., Cona, F., Chirico, G.B., Mazzoleni, S., Saracino, A., 2018. Windstorm
 568 disturbance triggers multiple species invasion in an urban Mediterranean forest. *iForest -*
 569 *Biogeosciences and Forestry* 11, 64–71. <https://doi.org/10.3832/ifor2374-010>
 570 Bonari, G., Acosta, A.T.R., Angiolini, C., 2017. Mediterranean coastal pine forest stands:
 571 Understorey distinctiveness or not? *Forest Ecology and Management* 391, 19–28.
 572 <https://doi.org/10.1016/j.foreco.2017.02.002>
 573 Brundu, G., 2014. Plant invaders in European and Mediterranean inland waters: profiles
 574 distribution, and threats. *Hydrobiologia* 746, 61–79. <https://doi.org/10.1007/s10750-014-1910-9>
 575 Brundu, G., Minicante, S.A., Barni, E., Bolpagni, R., Caddeo, A., Celesti-Gradow, L., Cogoni, A.,
 576 Galasso, G., Iiriti, G., Lazzaro, L., Loi, M.C., Lozano, V., Marignani, M., Montagnani, C., Siniscalco, C.,
 577 2020. Managing plant invasions using legislation tools: an analysis of the national and regional
 578 regulations for non-native plants in Italy. *Annali di Botanica* 10, 1–12.
 579 <https://doi.org/10.13133/2239-3129/16508>
 580 Buffa G., Del Vecchio S., Fantinato E., Milano V., 2018. Local versus landscape-scale effects of
 581 anthropogenic land-use on forest species richness. *Acta Oecologica* 86, 49–56.
 582 Calabrese, V., Frate, L., Iannotta, F., Prisco, I., Stanisci, A., 2017. *Acacia saligna*: an invasive species
 583 on the coast of Molise (southern Italy). *Forest@ - Rivista di Selvicoltura ed Ecologia Forestale* 14,
 584 28–33. <https://doi.org/10.3832/efor2211-013>
 585 Campagnaro, T., Nascimbene, J., Tasinazzo, S., Trentanovi, G., Sitzia, T., 2018. Exploring patterns
 586 drivers and structure of plant community composition in alien *Robinia pseudoacacia* secondary
 587 woodlands. *iForest - Biogeosciences and Forestry* 11, 586–593. [https://doi.org/10.3832/ifor2687-](https://doi.org/10.3832/ifor2687-011)
 588 [011](https://doi.org/10.3832/ifor2687-011)
 589 Campoy, J.G., Acosta, A.T.R., Affre, L., Barreiro, R., Brundu, G., Buisson, E., González, L., Lema, M.,
 590 Novoa, A., Retuerto, R., Roiloa, S.R., Fagúndez, J., 2018. Monographs of invasive plants in Europe:
 591 *Carpobrotus*. *Botany Letters* 165, 440–475. <https://doi.org/10.1080/23818107.2018.1487884>
 592 Carboni, M., Santoro, R., Acosta, A.T.R., 2010. Are some communities of the coastal dune zonation
 593 more susceptible to alien plant invasion? *Journal of Plant Ecology* 3, 139–147.
 594 <https://doi.org/10.1093/jpe/rtp037>
 595 Castro-Díez, P., Vaz, A.S., Silva, J.S., Loo, M., Álvaro Alonso, Aponte, C., Álvaro Bayón, Bellingham,
 596 P.J., Chiufo, M.C., DiManno, N., Julian, K., Kandert, S., Porta, N.L., Marchante, H., Maule, H.G.,
 597 Mayfield, M.M., Metcalfe, D., Monteverdi, M.C., Núñez, M.A., Ostertag, R., Parker, I.M., Peltzer,
 598 D.A., Potgieter, L.J., Raymundo, M., Rayome, D., Reisman-Berman, O., Richardson, D.M., Roos,
 599 R.E., Saldaña, A., Shackleton, R.T., Torres, A., Trudgen, M., Urban, J., Vicente, J.R., Vilà, M., Ylioja,
 600 T., Zenni, R.D., Godoy, O., 2019. Global effects of non-native tree species on multiple ecosystem
 601 services. *Biological Reviews*. <https://doi.org/10.1111/brv.12511>

602 Celesti-Grapow, L., Alessandrini, A., Arrigoni, P.V., Banfi, E., Bernardo, L., Bovio, M., Brundu, G.,
 603 Cagiotti, M.R., Camarda, I., Carli, E., Conti, F., Fascetti, S., Galasso, G., Gubellini, L., La Valva, V.,
 604 Lucchese, F., Marchiori, S., Mazzola, P., Peccenini, S., Poldini, L., Pretto, F., Prosser, F., Siniscalco,
 605 C., Villani, M.C., Viegi, L., Wilhalm, T., Blasi, C., 2009. Inventory of the non-native flora of Italy.
 606 Plant Biosystems - An International Journal Dealing with all Aspects of Plant Biology 143, 386–430.
 607 <https://doi.org/10.1080/11263500902722824>
 608 Celesti-Grapow, L., Bassi, L., Brundu, G., Camarda, I., Carli, E., D'Auria, G., Guacchio, E.D., Domina,
 609 G., Ferretti, G., Foggi, B., Lazzaro, L., Mazzola, P., Peccenini, S., Pretto, F., Stinca, A., Blasi, C., 2016.
 610 Plant invasions on small Mediterranean islands: An overview. Plant Biosystems - An International
 611 Journal Dealing with all Aspects of Plant Biology 150, 1119–1133.
 612 <https://doi.org/10.1080/11263504.2016.1218974>
 613 Ceschin, S., Abati, S., Leacche, I., Iamónico, D., Iberite, M., Zuccarello, V., 2016a. Does the alien
 614 *Lemna minuta* show an invasive behavior outside its original range? Evidence of antagonism with
 615 the native *L. minor* in central Italy. International Review of Hydrobiology 101, 173–181.
 616 <https://doi.org/10.1002/iroh.201601841>
 617 Ceschin, S., Abati, S., Traversetti, L., Spani, F., Grosso, F.D., Scalici, M., 2019. Effects of the invasive
 618 duckweed *Lemna minuta* on aquatic animals: evidence from an indoor experiment. Plant
 619 Biosystems - An International Journal Dealing with all Aspects of Plant Biology 153, 749–755.
 620 <https://doi.org/10.1080/11263504.2018.1549605>
 621 Ceschin, S., Bella, V.D., Piccari, F., Abati, S., 2016b. Colonization dynamics of the alien macrophyte
 622 *Lemna minuta* Kunth: a case study from a semi-natural pond in Appia Antica Regional Park (Rome
 623 Italy). Fundamental and Applied Limnology / Archiv für Hydrobiologie 188, 93–101.
 624 <https://doi.org/10.1127/fal/2016/0870>
 625 Ceschin, S., Ferrante, G., Mariani, F., Traversetti, L., Ellwood, N.T.W., 2020. Habitat change and
 626 alteration of plant and invertebrate communities in waterbodies dominated by the invasive alien
 627 macrophyte *Lemna minuta* Kunth. Biological Invasions 22, 1325–1337.
 628 <https://doi.org/10.1007/s10530-019-02185-5>
 629 Del Vecchio, S., Acosta, A., Stanisci, A., 2013. The impact of *Acacia saligna* invasion on Italian
 630 coastal dune EC habitats. Comptes Rendus Biologies 336, 364–369.
 631 <https://doi.org/10.1016/j.crv.2013.06.004>
 632 Del Vecchio, S., Pizzo L., Buffa G., 2015. The response of plant community diversity to alien
 633 invasion: evidence from a sand dune time series. Biodiversity and Conservation 24, 371–392.
 634 <https://doi.org/10.1007/s10531-014-0814-3>
 635 Domènech, R., Vilà, M., 2006. The role of successional stage vegetation type and soil disturbance
 636 in the invasion of the alien grass *Cortaderia selloana*. Journal of Vegetation Science 17, 591–598.
 637 <https://doi.org/10.1111/j.1654-1103.2006.tb02483.x>
 638 European Commission, 2013. Interpretation manual of European Union habitats EUR 28.
 639 Environment DG. Accessed March 24, 2020. URL:
 640 http://ec.europa.eu/environment/nature/legislation/habitatsdirective/docs/Int_Manual_EU28.pdf
 641 [f](http://ec.europa.eu/environment/nature/legislation/habitatsdirective/docs/Int_Manual_EU28.pdf)
 642 Fenesi, A., Geréd, J., Meiners, S.J., Tóthmérész, B., Török, P., Ruprecht, E., 2015a. Does disturbance
 643 enhance the competitive effect of the invasive *Solidago canadensis* on the performance of two
 644 native grasses? Biological Invasions 17, 3303–3315. <https://doi.org/10.1007/s10530-015-0954-8>
 645 Fenesi, A., Vágási, C.I., Beldean, M., Földesi, R., Kolcsár, L.-P., Shapiro, J.T., Török, E., Kovács-
 646 Hostyánszki, A., 2015b. *Solidago canadensis* impacts on native plant and pollinator communities in
 647 different-aged old fields. Basic and Applied Ecology 16, 335–346.
 648 <https://doi.org/10.1016/j.baae.2015.03.003>

649 Friedrichs, M., Hermoso, V., Bremerich, V., Langhans, S.D., 2018. Evaluation of habitat protection
650 under the European Natura 2000 conservation network. The example for Germany. PLoS One 13,
651 e0208264. <https://doi.org/10.1371/journal.pone.0208264>

652 Galasso, G., Conti, F., Peruzzi, L., Ardenghi, N.M.G., Banfi, E., Celesti-Grapow, L., Albano, A.,
653 Alessandrini, A., Bacchetta, G., Ballelli, S., Mazzanti, M.B., Barberis, G., Bernardo, L., Blasi, C.,
654 Bouvet, D., Bovio, M., Cecchi, L., Guacchio, E.D., Domina, G., Fascetti, S., Gallo, L., Gubellini, L.,
655 Guiggi, A., Iamónico, D., Iberite, M., Jiménez-Mejías, P., Lattanzi, E., Marchetti, D., Martinetto, E.,
656 Masin, R.R., Medagli, P., Passalacqua, N.G., Peccenini, S., Pennesi, R., Pierini, B., Podda, L., Poldini,
657 L., Prosser, F., Raimondo, F.M., Roma-Marzio, F., Rosati, L., Santangelo, A., Scoppola, A.,
658 Scortegagna, S., Selvaggi, A., Selvi, F., Soldano, A., Stinca, A., Wagensommer, R.P., Wilhalm, T.,
659 Bartolucci, F., 2018a. An updated checklist of the vascular flora alien to Italy. Plant Biosystems - An
660 International Journal Dealing with all Aspects of Plant Biology 152, 556–592.
661 <https://doi.org/10.1080/11263504.2018.1441197>

662 Galasso, G., Domina, G., Adorni, M., Ardenghi, N.M.G., Bonari, G., Buono, S., Cancellieri, L.,
663 Chianese, G., Ferretti, G., Fiaschi, T., Forte, L., Guarino, R., Labadessa, R., Lastrucci, L., Lazzaro, L.,
664 Magrini, S., Minuto, L., Mossini, S., Olivieri, N., Scoppola, A., Stinca, A., Turcato, C., Nepi, C., 2018b.
665 Notulae to the Italian alien vascular flora: 5. Italian Botanist 5, 45–56.
666 <https://doi.org/10.3897/italianbotanist.5.25910>

667 Galasso, G., Domina, G., Alessandrini, A., Ardenghi, N.M.G., Bacchetta, G., Ballelli, S., Bartolucci, F.,
668 Brundu, G., Buono, S., Busnardo, G., Calvia, G., Capece, P., D'Antraccoli, M., Nuzzo, L.D., Fanfarillo,
669 E., Ferretti, G., Guarino, R., Iamónico, D., Iberite, M., Latini, M., Lazzaro, L., Lonati, M., Lozano, V.,
670 Magrini, S., Mei, G., Mereu, G., Moro, A., Mugnai, M., Nicoletta, G., Nimis, P.L., Olivieri, N., Pennesi,
671 R., Peruzzi, L., Podda, L., Probo, M., Prosser, F., Enri, S.R., Roma-Marzio, F., Ruggero, A., Scafidi, F.,
672 Stinca, A., Nepi, C., 2018c. Notulae to the Italian alien vascular flora: 6. Italian Botanist 6, 65–90.
673 <https://doi.org/10.3897/italianbotanist.6.30560>

674 Gallardo, B., Aldridge, D.C., González-Moreno, P., Pergl, J., Pizarro, M., Pyšek, P., Thuiller, W.,
675 Yesson, C., Vilà, M., 2017. Protected areas offer refuge from invasive species spreading under
676 climate change. Global Change Biology 23, 5331–5343. <https://doi.org/10.1111/gcb.13798>

677 Gentili, R., Ferrè, C., Cardarelli, E., Montagnani, C., Bogliani, G., Citterio, S., & Comolli, R., 2019.
678 Comparing Negative Impacts of *Prunus serotina*, *Quercus rubra* and *Robinia pseudoacacia* on
679 Native Forest Ecosystems. Forests 10, 842. <https://doi.org/10.3390/f10100842>

680 Gentili, R., Montagnani, C., Gilardelli, F., Guarino, M.F., Citterio, S., 2017. Let native species take
681 their course: *Ambrosia artemisiifolia* replacement during natural or artificial succession. Acta
682 Oecologica 82, 32–40. <https://doi.org/10.1016/j.actao.2017.05.007>

683 Gigante, D., Acosta, A.T.R., Agrillo, E., Armiraglio, S., Assini, S., Attorre, F., Bagella, S., Buffa, G.,
684 Casella, L., Giancola, C., Giusso del Galdo, G.P., Marcenò, C., Pezzi, G., Prisco, I., Venanzoni, R.,
685 Viciani, D., 2018. Habitat conservation in Italy: the state of the art in the light of the first European
686 Red List of Terrestrial and Freshwater Habitats. Rendiconti Lincei. Scienze Fisiche e Naturali 29,
687 251–265.

688 GISD, 2020. Global Invasive Species Database. Accessed March 24, 2020. URL:
689 <http://www.iucngisd.org/gisd/>

690 Guerra, C., Baquero, R.A., Gutiérrez-Arellano, D., Nicola, G.G., 2018. Is the Natura 2000 network
691 effective to prevent the biological invasions? Global Ecology and Conservation 16, e00497.
692 <https://doi.org/10.1016/j.gecco.2018.e00497>

693 Hulme, P.E., Pyšek, P., Jarošík, V., Pergl, J., Schaffner, U., Vilà, M., 2013. Bias and error in
694 understanding plant invasion impacts. Trends in Ecology & Evolution 28, 212–218.
695 <https://doi.org/10.1016/j.tree.2012.10.010>

696 Jucker, T., Carboni, M., Acosta, A.T.R., 2013. Going beyond taxonomic diversity: deconstructing
 697 biodiversity patterns reveals the true cost of iceplant invasion. *Diversity and Distributions* 19,
 698 1566–1577. <https://doi.org/10.1111/ddi.12124>
 699 Kleinbauer, I., Dullinger, S., Peterseil, J., Essl, F., 2010. Climate change might drive the invasive tree
 700 *Robinia pseudacacia* into nature reserves and endangered habitats. *Biological Conservation* 143,
 701 382–390. <https://doi.org/10.1016/j.biocon.2009.10.024>
 702 Lastrucci, L., Lazzaro, L., Dell’Olmo, L., Foggi, B., Cianferoni, F., 2017. Impacts of *Myriophyllum*
 703 *aquaticum* invasion in a Mediterranean wetland on plant and macro-arthropod communities.
 704 *Plant Biosystems - An International Journal Dealing with all Aspects of Plant Biology* 152, 427–435.
 705 <https://doi.org/10.1080/11263504.2017.1303002>
 706 Latombe, G., Pyšek, P., Jeschke, J.M., Blackburn, T.M., Bacher, S., Capinha, C., Costello, M.J.,
 707 Fernández, M., Gregory, R.D., Hobern, D., Hui, C., Jetz, W., Kumschick, S., McGrannachan, C., Pergl,
 708 J., Roy, H.E., Scalera, R., Squires, Z.E., Wilson, J.R.U., Winter, M., Genovesi, P., McGeoch, M.A.,
 709 2017. A vision for global monitoring of biological invasions. *Biological Conservation* 213, 295–308.
 710 <https://doi.org/10.1016/j.biocon.2016.06.013>
 711 Lazzaro, L., Bolpagni, R., Barni, E., Brundu, G., Blasi, C., Siniscalco, C., Celesti-Grapow, L., 2019.
 712 Towards alien plant prioritization in Italy: methodological issues and first results. *Plant Biosystems*
 713 - An International Journal Dealing with all Aspects of Plant Biology 153, 740–746.
 714 <https://doi.org/10.1080/11263504.2019.1640310>
 715 Lazzaro, L., Giuliani, C., Benesperi, R., Calamassi, R., Foggi, B., 2015. Plant species loss and
 716 community nestedness after leguminous tree *Acacia pycnantha* invasion in a Mediterranean
 717 ecosystem. *Folia Geobotanica* 50, 229–238. <https://doi.org/10.1007/s12224-015-9222-z>
 718 Lazzaro, L., Giuliani, C., Fabiani, A., Agnelli, A.E., Pastorelli, R., Lagomarsino, A., Benesperi, R.,
 719 Calamassi, R., Foggi, B., 2014. Soil and plant changing after invasion: The case of *Acacia dealbata*
 720 in a Mediterranean ecosystem. *Science of The Total Environment* 497-498, 491–498.
 721 <https://doi.org/10.1016/j.scitotenv.2014.08.014>
 722 Lazzaro, L., Mazza, G., d’Errico, G., Fabiani, A., Giuliani, C., Inghilesi, A.F., Lagomarsino, A., Landi, S.,
 723 Lastrucci, L., Pastorelli, R., Roversi, P.F., Torrini, G., Tricarico, E., Foggi, B., 2018. How ecosystems
 724 change following invasion by *Robinia pseudoacacia*: Insights from soil chemical properties and soil
 725 microbial nematode, microarthropod and plant communities. *Science of The Total Environment*
 726 622-623, 1509–1518. <https://doi.org/10.1016/j.scitotenv.2017.10.017>
 727 Mainka, S.A., Howard, G.W., 2010. Climate change and invasive species: double jeopardy.
 728 *Integrative Zoology*, 5: 102-111. doi:10.1111/j.1749-4877.2010.00193.x
 729 Maes, J., 2013. A model for the assessment of habitat conservation status in the EU. JRC scientific
 730 and policy report. European Commission Joint Research Centre Institute for Environment and
 731 Sustainability, Luxembourg, Publication Office of the European Commission.
 732 Maiorca, G., Spampinato, G., Crisafulli, A., Cameriere, P., 2007. Flora vascolare e vegetazione della
 733 Riserva Naturale Regionale Foce del Fiume Crati (Calabria Italia meridionale). *Webbia* 62, 121–174.
 734 <https://doi.org/10.1080/00837792.2007.10670821>
 735 Malavasi, M., Acosta, A.T.R., Carranza, M.L., Bartolozzi, L., Basset, A., Bassignana, M., Campanaro,
 736 A., Canullo, R., Carruggio, F., Cavallaro, V., Cianferoni, F., Cindolo, C., Cocciuffa, C., Corriero, G.,
 737 D’Amico, F.S., Forte, L., Freppaz, M., Mantino, F., Matteucci, G., Pierri, C., Stanisci, A., Colangelo, P.,
 738 2018. Plant invasions in Italy: An integrative approach using the European LifeWatch infrastructure
 739 database. *Ecological Indicators* 91, 182–188. <https://doi.org/10.1016/j.ecolind.2018.03.038>
 740 Mazaris, A.D., Katsanevakis, S., 2018. The threat of biological invasions is under-represented in the
 741 marine protected areas of the European Natura 2000 network. *Biological Conservation* 225, 208–
 742 212. <https://doi.org/10.1016/j.biocon.2018.07.007>

743 McLean, P., Gallien, L., Wilson, J.R.U., Gaertner, M., Richardson, D.M., 2017. Small urban centres
 744 as launching sites for plant invasions in natural areas: insights from South Africa. *Biological*
 745 *Invasions* 19, 3541–3555. <https://doi.org/10.1007/s10530-017-1600-4>
 746 Minuto, L., Casazza, G., Dagnino, D., Guerrina, M., Macrì, C., Zappa, E., Mariotti, M.G., 2020.
 747 Reproductive traits of the invasive species *Acacia dealbata* in the Northern Mediterranean basin.
 748 *Annali di Botanica* 10, 13–20. <https://doi.org/10.13133/2239-3129/15642>
 749 Montecchiari, S., Tesei, G., Allegrezza, M. 2020. Effects of *Robinia pseudoacacia* coverage on
 750 diversity and environmental conditions of central-northern Italian *Quercus pubescens* sub-
 751 Mediterranean forests (Habitat code 91AA*): A threshold assessment. *Annali di Botanica*, 10, 33–
 752 54. <https://doi.org/10.13133/2239-3129/16447>.
 753 Nascimbene, J., Benesperi, R., Casazza, G., Chiarucci, A., & Giordani, P., 2020. Range shifts of
 754 native and invasive trees exacerbate the impact of climate change on epiphyte distribution: The
 755 case of lung lichen and black locust in Italy. *Science of The Total Environment* 735, 139537.
 756 Nascimbene, J., Lazzaro, L., Benesperi, R., 2015. Patterns of β -diversity and similarity reveal biotic
 757 homogenization of epiphytic lichen communities associated with the spread of black locust
 758 forests. *Fungal Ecology* 14, 1–7. <https://doi.org/10.1016/j.funeco.2014.10.006>
 759 Nascimbene, J., Nimis, P.L., Benesperi, R., 2012. Mature non-native black-locust (*Robinia*
 760 *pseudoacacia* L.) forest does not regain the lichen diversity of the natural forest. *Science of The*
 761 *Total Environment* 421-422, 197–202. <https://doi.org/10.1016/j.scitotenv.2012.01.051>
 762 Oduor, A.M.O., Leimu, R., van Kleunen, M., 2016. Invasive plant species are locally adapted just as
 763 frequently and at least as strongly as native plant species. *Journal of Ecology* 104, 957–968.
 764 <https://doi.org/10.1111/1365-2745.12578>
 765 Pyšek, P., Jarošík, V., Hulme, P. E., Pergl, J., Hejda, M., Schaffner, U., & Vilà, M., 2012. A global
 766 assessment of invasive plant impacts on resident species, communities and ecosystems: the
 767 interaction of impact measures, invading species' traits and environment. *Global Change Biology*
 768 18(5), 1725–1737.
 769 R Core Team, 2019. R: A language and environment for statistical computing. R Foundation for
 770 Statistical Computing, Vienna, Austria. URL <https://www.r-project.org/>
 771 Santoro, R., Carboni, M., Carranza, M.L., Acosta, A.T.R., 2012. Focal species diversity patterns can
 772 provide diagnostic information on plant invasions. *Journal for Nature Conservation* 20, 85–91.
 773 <https://doi.org/10.1016/j.inc.2011.08.003>
 774 Santoro, R., Jucker, T., Carranza, M., Acosta, A., 2011. Assessing the effects of *Carpobrotus*
 775 invasion on coastal dune soils. Does the nature of the invaded habitat matter? *Community Ecology*
 776 12, 234–240. <https://doi.org/10.1556/comec.12.2011.2.12>
 777 Sarmati, S., Conti, L., Acosta, A.T.R., 2019. *Carpobrotus acinaciformis* vs *Carpobrotus edulis*: Are
 778 there any differences in their impact on coastal dune plant biodiversity? *Flora* 257, 151422.
 779 <https://doi.org/10.1016/j.flora.2019.151422>
 780 Seebens, H., Blackburn, T.M., Dyer, E.E., Genovesi, P., Hulme, P.E., Jeschke, J.M., Pagad, S., Pyšek,
 781 P., Winter, M., Arianoutsou, M., Bacher, S., Blasius, B., Brundu, G., Capinha, C., Celesti-Grapow, L.,
 782 Dawson, W., Dullinger, S., Fuentes, N., Jäger, H., Kartesz, J., Kenis, M., Kreft, H., Kühn, I., Lenzner,
 783 B., Liebhold, A., Mosena, A., Moser, D., Nishino, M., Pearman, D., Pergl, J., Rabitsch, W., Rojas-
 784 Sandoval, J., Roques, A., Rorke, S., Rossinelli, S., Roy, H.E., Scalera, R., Schindler, S., Štajerová, K.,
 785 Tokarska-Guzik, B., van Kleunen, M., Walker, K., Weigelt, P., Yamanaka, T., Essl, F., 2017. No
 786 saturation in the accumulation of alien species worldwide. *Nature Communications* 8, 14435.
 787 <https://doi.org/10.1038/ncomms14435>
 788 Sitzia, T., Campagnaro, T., Kotze, D.J., Nardi, S., Ertani, A., 2018. The invasion of abandoned fields
 789 by a major alien tree filters understory plant traits in novel forest ecosystems. *Scientific Reports* 8,
 790 8410. <https://doi.org/10.1038/s41598-018-26493-3>

791 Sperandii, M.G., Prisco, I., Acosta, A.T.R., 2017. Hard times for Italian coastal dunes: insights from a
792 diachronic analysis based on random plots. *Biodiversity and Conservation* 27, 633–646.
793 <https://doi.org/10.1007/s10531-017-1454-1>

794 Traveset, A., Brundu, G., Carta, L., Mprezetou, I., Lambdon, P., Manca, M., Médail, F., Moragues,
795 E., Rodríguez-Pérez, J., Siamantziouras, A.-S.D., Suehs, C.M., Troumbis, A.Y., Vilà, M., Hulme, P.E.,
796 2008. Consistent performance of invasive plant species within and among islands of the
797 Mediterranean basin. *Biological Invasions* 10, 847–858. [https://doi.org/10.1007/s10530-008-9245-](https://doi.org/10.1007/s10530-008-9245-y)
798 [y](https://doi.org/10.1007/s10530-008-9245-y)

799 Trochet, A., Schmeller, D., 2013. Effectiveness of the Natura 2000 network to cover threatened
800 species. *Nature Conservation* 4, 35–53. <https://doi.org/10.3897/natureconservation.4.3626>

801 Tóth, V.R., Villa, P., Pinardi, M., Bresciani, M., 2019. Aspects of Invasiveness of *Ludwigia* and
802 *Nelumbo* in Shallow Temperate Fluvial Lakes. *Frontiers in Plant Science* 10.
803 <https://doi.org/10.3389/fpls.2019.00647>

804 Uboni, C., Tordoni, E., Brandmayr, P., Battistella, S., Bragato, G., Castello, M., Colombetta, G.,
805 Poldini, L., Bacaro, G., 2019. Exploring cross-taxon congruence between carabid beetles
806 (Coleoptera: Carabidae) and vascular plants in sites invaded by *Ailanthus altissima* versus non-
807 invaded sites: The explicative power of biotic and abiotic factors. *Ecological Indicators* 103, 145–
808 155. <https://doi.org/10.1016/j.ecolind.2019.03.052>

809 Vanderhoeven, S., Branquart, E., Casaer, J., D’hondt, B., Hulme, P.E., Shwartz, A., Strubbe, D.,
810 Turbé, A., Verreycken, H., Adriaens T., 2017. Beyond protocols: improving the reliability of expert-
811 based risk analysis underpinning invasive species policies. *Biological Invasions* 19, 2507–2517
812 (2017). <https://doi.org/10.1007/s10530-017-1434-0>

813 Vegini, E., Lastrucci, L., Lazzaro, L., Cardarelli, E., Martignoni, M., 2020. Impact of *Prunus serotina*
814 Ehrh. invasion on heathland vegetation: a case of study in North-Western Italy. *Biologia* 75, 327–
815 336. <https://doi.org/10.2478/s11756-019-00408-7>

816 Viciani, D., Vidali, M., Gigante, D., Bolpagni, R., Villani, M., Acosta, A.T.R., Adorni, M., Aleffi, M.,
817 Allegrezza, M., Angiolini, C., Assini, S., Bagella, S., Bonari, G., Bovio, M., Bracco, F., Brundu, G.,
818 Buffa, G., Caccianiga, M., Carnevali, L., Ceschin, S., Ciaschetti, G., Cogoni, A., Cecco, V.D., Foggi, B.,
819 Frattaroli, A.R., Genovesi, P., Gentili, R., Lazzaro, L., Lonati, M., Lucchese, F., Mainetti, A., Mariotti,
820 M., Minissale, P., Paura, B., Pellizzari, M., Perrino, E.V., Pirone, G., Poggio, L., Poldini, L., Poponessi,
821 S., Prisco, I., Prosser, F., Puglisi, M., Rosati, L., Selvaggi, A., Sottovia, L., Spampinato, G., Stanisci, A.,
822 Stinca, A., Venanzoni, R., Lastrucci, L., 2020. A first checklist of the alien-dominated vegetation in
823 Italy. *Plant Sociology*. In Press.

824 Vilà, M., Tessier, M., Suehs, C.M., Brundu, G., Carta, L., Galanidis, A., Lambdon, P., Manca, M.,
825 Médail, F., Moragues, E., Traveset, A., Troumbis, A.Y., Hulme, P.E., 2006. Local and regional
826 assessments of the impacts of plant invaders on vegetation structure and soil properties of
827 Mediterranean islands. *Journal of Biogeography* 33, 853–861. [https://doi.org/10.1111/j.1365-](https://doi.org/10.1111/j.1365-2699.2005.01430.x)
828 [2699.2005.01430.x](https://doi.org/10.1111/j.1365-2699.2005.01430.x)

829 Villa, P., Pinardi, M., Bolpagni, R., Gillier, J.-M., Zinke, P., Nedelcuț, F., Bresciani, M., 2018.
830 Assessing macrophyte seasonal dynamics using dense time series of medium resolution satellite
831 data. *Remote Sensing of Environment* 216, 230–244. <https://doi.org/10.1016/j.rse.2018.06.048>

832 Villa, P., Pinardi, M., Tóth, V.R., Hunter, P.D., Bolpagni, R., Bresciani, M., 2017. Remote sensing of
833 macrophyte morphological traits: implications for the management of shallow lakes. *Journal of*
834 *Limnology*. <https://doi.org/10.4081/jlimnol.2017.1629>


835 Vilà, M., Hulme, P.E., 2017. *Impact of Biological Invasions on Ecosystem Services*. Springer
836 International Publishing. <https://doi.org/10.1007/978-3-319-45121-3>

837 Vítková, M., Müllerová, J., Sádlo, J., Pergl, J., Pyšek, P., 2017. Black locust (*Robinia pseudoacacia*)
838 beloved and despised: A story of an invasive tree in Central Europe. Forest Ecology and
839 Management 384, 287–302. <https://doi.org/10.1016/j.foreco.2016.10.057>
840 Walker, K.J., Preston, C.D., Boon, C.R., 2009. Fifty years of change in an area of intensive
841 agriculture: plant trait responses to habitat modification and conservation Bedfordshire, England.
842 Biodiversity and Conservation 18, 3597–3613. <https://doi.org/10.1007/s10531-009-9662-y>
843 Zedda, L., Cogoni, A., Flore, F., Brundu, G., 2010. Impacts of alien plants and man-made
844 disturbance on soil-growing bryophyte and lichen diversity in coastal areas of Sardinia (Italy). Plant
845 Biosystems - An International Journal Dealing with all Aspects of Plant Biology 144, 547–562.
846 <https://doi.org/10.1080/11263501003638604>
847

848
849
850
851
852
853
854
855

FIGURES


Figure 1. Distribution of records on the impact of IAPs in Italy *per* administrative region. A) Distribution of records on impact mechanisms. B) Number of IAPs exerting any type of impact mechanism. C) Distribution of records on impact outcomes on plant communities. D) Number of N2000 Habitat types exposed to some degree of ecological impact by IAPs. E) Percentage of N2000 Habitats exposed to some degree of ecological impact by IAPs on the total number of Habitat harboured in the region.



856

857
858
859
860


Figure 2. Number of data collected regarding the presence of impact mechanism (empty green circles, dashed line) and impact outcomes (full blue circles, solid line) exerted by IAPs in Italy.



861
862

863
864
865

Figure 3 A) Number of regional records with specific impact mechanisms for the first 23 IAPs. B) Number of total records of impact outcomes for the first 23 IAPs.




866
867

868
869
870


871
872

Figure 4. Source of the reports (in percentage) concerning A) impact mechanisms and B) impact outcomes for all the species and for the first ten IAPs (in descending order by number of records).



873
874
875
876
877


Figure 5. Impact outcomes. A) Total records (i.e. species × N2000 habitat × administrative region). B) Total number of N2000 Habitats suffering from a specific impact outcome (irrespective of species and administrative regions).



878
879

880
881
882
883
884

Figure 6. Data on the impact outcomes recorded per N2000 Habitat (*sensu* Habitat directive 92/43/EEC). A) Number of species exerting some degree of ecological impact for each target N2000 habitat and B) number of administrative regions in which the target N2000 habitat is impacted.



885

886


887

888

889

890

Figure 7. Effect of the life form categories (A, C, E) and introduction period (B, D, F) on the number of N2000 Habitats (Habitats), number of macro-categories of N2000 Habitats (Macro-habitats) and the number of invaded administrative regions (regions), respectively. C = chamaephytes; G = geophytes; H = hemicryptophytes; I = hydrophytes; P = phanerophytes; T = terophytes.



891

892