

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/284182018>

Molecular diagnosis of native and quarantine pest thrips (Thysanoptera, Thripidae) of southern European citrus orchards.

Article in *Bulletin of Insectology* · June 2016

CITATIONS

0

READS

179

3 authors:



[Alessandra De Grazia](#)

Mediterranean University of Reggio Calabria

17 PUBLICATIONS 12 CITATIONS

[SEE PROFILE](#)



[Rita Marullo](#)

Mediterranean University of Reggio Calabria

76 PUBLICATIONS 620 CITATIONS

[SEE PROFILE](#)



[Gerald B. Moritz](#)

Institute of Biology

120 PUBLICATIONS 519 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Invasive thrips species to southern European Regions [View project](#)



Implementation of integrated thrips and tospovirus management strategies in smallholder vegetable cropping systems of eastern Africa [View project](#)

Molecular diagnosis of native and quarantine pest thrips of southern European citrus orchards

Alessandra DE GRAZIA¹, Rita MARULLO¹, Gerald MORITZ²

¹Dipartimento di Agraria, Università degli Studi Mediterranea di Reggio Calabria, Reggio Calabria, Italy

²Institute of Biology, Martin-Luther-University Halle-Wittenberg, Faculty of Natural Sciences I, Halle, Germany

Abstract

The ITS-RFLP technique of the amplified internal transcribed spacer regions of ribosomal DNA has been used to establish the molecular identification of the most common pest thrips species of citrus orchards. Both native and quarantine thrips species have been tested and molecular keys are proposed through ITS-RFLP techniques. The restriction enzymes produced patterns that allowed unambiguous identification of seven thrips species examined (visual key). The sequences of species can also be verified using a molecular key that permits identification of a species through such parameters as primer pair used, restriction enzyme, length of PCR-product, fragments obtained. This study has shown that the use of genetic markers can be a valid alternative for quarantine workers and for epidemiological researchers, to whom the correct identification of pest species through classic morphological methods could be either very difficult and time-consuming or visually impossible.

Key words: molecular identification, ITS-RFLP, visual key, thrips, citrus pest.

Introduction

The insect order Thysanoptera includes in the family Thripidae about 50 species associated with economic crops and also quarantine species. Problems related to quarantine species are mainly due to identification: difficulties related to their small size, variability of body colour, morphological patterns, secondary sexual character states and the cryptic behaviour of some species which exhibit large differences in genetic compositions, habitat preference, *Tospovirus* transmission efficiency (Brunner *et al.*, 2004; Hoddle *et al.*, 2008; Brunner and Frey, 2010; Rugman-Jones *et al.*, 2010; Jacobson *et al.*, 2013). Classical morphological identification keys require experience and are difficult for non-specialists such as most quarantine workers. Moreover, these difficulties are greater when an identification of either larval stages or eggs that infest trade vegetables is required. Larval identification keys are rare (Kucharczyk, 2010; Vierbergen *et al.*, 2010) and practically useless in terms of fast and accurate pest identification, but during the past 15 years identification systems which combine classic morphological knowledge with molecular identification methods have been produced (Moritz *et al.*, 2000; 2004; 2009; 2010; 2013; Tyagi *et al.*, 2015). Furthermore, improving molecular techniques have led to molecular keys that provide quick and correct identification ways for adults and all ontogenetic stages of the most common pest thrips species (Brunner *et al.*, 2002; Toda *et al.*, 2002; Kox *et al.*, 2005; Rugman-Jones *et al.*, 2006; Timm *et al.*, 2008; Fekrat *et al.*, 2015; Przybylska *et al.*, 2015). The internal transcribed spacer (ITS), the non-coding fragment of the nuclear ribosomal region, has been one of the most widely used markers in thrips species identification (Rugman-Jones *et al.*, 2006; Farris *et al.*, 2010).

In the present article, a molecular technique is developed to identify the most common harmful thrips species on citrus plants in the southern areas of Italy and

the Mediterranean Basin, together with a few exotic thrips species that are the most important quarantine pests for these crops. Results of molecular identification study of seven pest thrips species on citrus (Marullo and De Grazia, 2012) are presented and discussed; four of them are present in the Mediterranean Region, i.e. *Heliethrips haemorrhoidalis* (Bouche), *Frankliniella occidentalis* (Pergande), *Pezothrips kellyanus* (Bagnall), *Thrips tabaci* Lindeman, whereas three species, *Frankliniella bispinosa* (Morgan), *Scirtothrips aurantii* Faure, and *Scirtothrips citri* (Moulton) are considered quarantine species for EU territories.

Materials and methods

Collections of thrips species

Seventy individual thrips identified as seven different species were collected for this study (collecting details are summarized in table 1). The specimens of *F. bispinosa*, *S. citri* and *S. aurantii* from field populations, were provided by Prof. J Funderburk (University of Florida, USA) and Prof. L. Mound (CSIRO, Australia), respectively.

Molecular technique

Ten specimens of each species studied have been tested using a protocol which provides a few phases:

- DNA extraction: DNA has been extracted from single specimens using the Qiagen “DNeasy Tissue” kit (Germany) which allows the carcass of each specimen to be retrieved and mounted onto microscope slides. With the aid of a sterilized needle, we incised the specimen's abdomen and placed it in a 1.5 ml tube containing 20 µl proteinase K. After thoroughly mix by vortexing, the tube was incubated overnight at 55 °C. Then, we added 200 µl Buffer AL (provided with the kit) to the sample and vortexed it. After a short incubation (15 minutes at 70 °C) we added 200 µl ethanol (100%)

Table 1. Sampling details for the collected species.

Species	Locality	Date of collection	Geographic coordinates	Host-plant
<i>Frankliniella occidentalis</i>	Reggio Calabria (South Italy)	23/05/2012	38°6'41.40"N 15°39'43.56"E 31 m asl	<i>Solanum lycopersicum</i>
<i>Frankliniella bispinosa</i>	Miami (Florida, USA)	15/04/2011	25°46'27"N 80°11'37"W 25 m asl	<i>Lagerstroemia speciosa</i>
<i>Pezothrips kellyanus</i>	Reggio Calabria (South Italy)	30/04/2012	38°6'41.40"N 15°39'43.56"E 31 m asl	<i>Citrus limon</i>
<i>Heliothrips haemorrhoidalis</i>	Lamezia Terme (South Italy)	20/09/2011	38°58'0.84"N 16°18'36.00"E 216 m asl	<i>Viburnum</i> spp.
<i>Thrips tabaci</i>	Lamezia Terme (South Italy)	22/05/2012	38°58'0.84"N 16°18'36.00"E 216 m asl	<i>Allium caepa</i>
<i>Scirtothrips aurantii</i>	Canberra (Australia)	12/10/2011	35°17'00"S 149°07'41"E 571 m asl	<i>Citrus</i> spp.
<i>Scirtothrips citri</i>	Davis (California, USA)	13/10/2011	38°32'41"N 121°44'25"W 17 m asl	<i>Citrus</i> spp.

and mix all again. We inserted the mixture into DNeasy Mini spin column (provided with the kit) placed in a 2 ml collection tube and centrifuged. We discarded flow-through and transferred the spin column to a new 2 ml tube where we added 500 µl Buffer AW1 (provided) and after centrifugation, we again discarded flow-through and transferred the spin column to a new 2 ml tube in which we added 500 µl Buffer AW2 (provided) and centrifuged to dry the DNeasy membrane. We placed the DNeasy Mini spin column in a clean 1.5 ml tube and we added 70 µl sterile, ultra pure water directly onto the DNeasy membrane. Finally we incubated at room temperature for 1 minute, and centrifuged to elute DNA;

- PCR conditions: the ITS regions have been amplified with 2 primer pairs (CS249/CS250: 5'-TCGTAACAAGGTTTCCG-3'; 5'-GTTAGTTTCTTTTCCCTC-3'; 18SMP/28SMP: 5'-TGAACCTGCGGAAGGAT-3'; 5'-TCTCACCTBACTGAGG-3') and the polymerase chain reaction mixture contained: 10 µl of template; 5.0 µl of 10x PCR-buffer; 0.4 µl of 25 mM dNTPs; 4.0 µl of 25 mM MgCl₂; 1.5 µl of each primer; 0.2 µl of Taq-polymerase and sterile distilled water to a final volume of 50 µl. The amplification is carried out in an Eppendorf Mastercycler gradient. The DNA has been denatured at 95 °C for 3 minutes, followed by 30 cycles of denaturation at 95 °C for 45 seconds. Annealing (the temperature at which the primer anneals or binds to the target site) at 47 °C (CS249-CS250) or 54 °C (18SMP-28SMP) for 45 seconds. and elongation at 72 °C for 2 minutes. The last cycle is followed by 4 minutes of incubation at 72 °C to complete any partially synthesized strands;

- Restriction: 7 µl of the PCR-products have been digested with 3 units of five enzymes (RsaI, HaeIII, MspI,

HinfI and AluI) (Promega) and observed by electrophoresis in a 2% agarose gel with ethidium bromide to visualize the DNA restriction fragments (Moritz *et al.*, 2000).

Results

The technique of RFLP of the amplified internal transcribed spacer region of ribosomal DNA (ITS1 and ITS2) enables the production of specific DNA fragment patterns for the identification of thrips species. The identification of a species has to be done through comparison of different patterns obtained with the same couple of primers (visual key) (figures 1-2).

Figure 1 shows the differences between *F. bispinosa*, *P. kellyanus*, *S. citri* and *T. tabaci* whose DNA was amplified using the primers 18SMP-28SMP. In all the species, patterns are represented by a sequence of bands for each restriction enzyme used (table 2). In particular, lane 3 corresponds to restriction fragments obtained by digestion of rDNA of *F. bispinosa* using RsaI enzyme; lane 4 coincides in restriction fragments by digestion of rDNA with HaeIII enzyme; lane 5 shows the bands obtained by MspI enzyme; lane 6 and 7 represent restriction fragments obtained with HinfI and AluI, respectively. Similarly, lanes 11-15, 19-23 and 27-31, represent the bands obtained through the digestion of rDNA of *P. kellyanus*, *T. tabaci*, *S. citri*, respectively.

Figure 2 reproduces the patterns of *F. bispinosa* and *F. occidentalis* by using the primers CS249-CS250. Each of the seven species constitutes a distinct pattern. Figure 3 shows the patterns of the second instar larvae of *F. occidentalis* and *P. kellyanus* obtained by digestion of rDNA with 5 restriction enzymes (RsaI, HaeIII, MspI, HinfI and AluI).




Figure 1. Agarose gel showing the ITS-RFLP patterns of *F. bispinosa* (lanes 3-7), *P. kellyanus* (lanes 11-15), *T. tabaci* (lanes 19-23) and *S. citri* (lanes 27-31) obtained by digestion of amplified ITS regions of rDNA (18SMP-28SMP) with *RsaI*, *HaeIII*, *MspI*, *HinfI* and *AluI*. Lanes 1, 8, 9, 16, 17, 24, 25 and 32, 100 bp DNA ladder. Lanes 2, 10, 18 and 26, PCR products of *F. bispinosa*, *P. kellyanus*, *T. tabaci* and *S. citri* respectively.

Table 2. Size (in base pairs) of the ITS-RFLP restriction fragments of each species tested.

Species	Primer Pair	PCR-product	<i>RsaI</i>	<i>HaeIII</i>	<i>MspI</i>	<i>HinfI</i>	<i>AluI</i>
<i>S. aurantii</i>	18SMP-28SMP	1252	693, 176, 141	555, 506, 432, 278, 167, 144	663, 352, 225	647, 244, 145	628, 495, 167
<i>S. citri</i>	18SMP-28SMP	1420	692, 569, 523, 192	292, 256, 213, 154, 131	942, 901, 555, 349, 229, 156	526, 274, 175, 127	583, 345, 216
<i>F. occidentalis</i>	18SMP-28SMP	1360	358, 305, 254	562, 408	553, 369, 223, 141	460, 310, 253, 191	485, 280
<i>F. bispinosa</i>	18SMP-28SMP	1380	455, 300, 202	640, 428, 177	596, 330, 290	708, 332, 285	431, 394
<i>P. kellyanus</i>	18SMP-28SMP	1299	800, 288, 187	224, 214, 170, 146, 127	771, 257, 162, 104	485, 290, 264, 198, 176, 127	782, 654, 485, 446, 360, 328
<i>T. tabaci</i>	18SMP-28SMP	1619	481, 398, 211, 146	451, 365, 263, 193	1141, 435	1010, 408, 305	300, 255, 217, 198
<i>H. haemorrhoidalis</i>	CS249-CS250	865	690	315, 249, 173	700	374	390
<i>F. occidentalis</i>	CS249-CS250	1400	464, 390, 276	595, 521, 204	591, 457, 260, 182	476, 320, 294	543, 281
<i>F. bispinosa</i>	CS249-CS250	1349	478, 239, 130, 112, 102	688, 473, 114, 106	608, 277, 249, 148	732, 320, 279	465, 400, 180
<i>P. kellyanus</i>	CS249-CS250	1370	850, 314, 195	235, 193, 165, 144	810, 271, 164	495, 279, 223, 187, 154	819, 368, 192

Sequences of species determined by ITS-RFLP analysis were used to develop two molecular identification keys to confirm the identification of a species through a series of parameters as: primer pair used, restriction enzyme, length of PCR-product, fragments obtained. In a first key (table 3) six species studied were included, considering differences in size of the PCR products and restriction fragment lengths using the primer pair 18SMP-28SMP. These primers did not work on *H. haemorrhoidalis*. Therefore, a second key

for the identification of four species (table 4) was obtained using the primer pair CS249-CS250. Keys require the user to make a choice between only two data at a time. The choices are numbered and refer to the size of the ITS band and the number of fragments obtained by a precise restriction enzyme. In addition, for a further confirmation of the correct identification of a species it is possible to connect to the web-data base: <http://moritz.zoologie.uni-halle.de/> (Moritz, 2010).




Figure 2. Agarose gel showing the ITS-RFLP patterns of *F. bispinosa* (lanes 3-7) and *F. occidentalis* (lanes 11-15), obtained by digestion of amplified ITS regions of rDNA (CS249-CS250) with *Rsa*I, *Hae*III, *Msp*I, *Hinf*I and *Alu*I. Lanes 1, 8, 9 and 16, 100 bp DNA ladder. Lanes 2 and 10, PCR products of analysed species.




Figure 3. Agarose gel showing the ITS-RFLP patterns of larval stages of *F. occidentalis* (lanes 3-7) and *P. kellyanus* (lanes 11-15) obtained by digestion of amplified ITS regions of rDNA (CS249-CS250) with *Rsa*I, *Hae*III, *Msp*I, *Hinf*I and *Alu*I. Lanes 2 and 10 represent PCR products. Lanes 1 and 16, 100 bp DNA ladder.

Table 3. Molecular key to thrips species based on ITS-RFLP products obtained with 18SMP-28SMP.

1	Size of the ITS band > 1260 bp	2
	Size of the ITS band ≤ 1260 bp	<i>S. aurantii</i>
2	Size of the ITS band > 1400 bp	3
	Size of the ITS band ≤ 1400 bp	4
3	PCR product restricted with <i>Msp</i> I gives > 4 bands	<i>S. citri</i>
	PCR product restricted with <i>Msp</i> I gives ≤ 3 bands	<i>T. tabaci</i>
4	PCR product restricted with <i>Hae</i> III gives 2 bands	<i>F. occidentalis</i>
	PCR product restricted with <i>Hae</i> III gives ≥ 3 bands	5
5	PCR product restricted with <i>Msp</i> I gives 3 bands	<i>F. bispinosa</i>
	PCR product restricted with <i>Msp</i> I gives ≥ 4 bands	<i>P. kellyanus</i>

Table 4. Molecular key to thrips species based on ITS-RFLP products obtained with CS249-CS250.

1	Size of the ITS band > 1000 bp	2
	Size of the ITS band < 1000 bp	<i>H. haemorrhoidalis</i>
2	Size of the ITS band > 1350 bp	3
	Size of the ITS band < 1350 bp	4
3	PCR product restricted with <i>Rsa</i> I gives 3 bands	4
	PCR product restricted with <i>Rsa</i> I gives > 3 bands	<i>F. bispinosa</i>
4	PCR product restricted with <i>Msp</i> I gives 3 bands	<i>P. kellyanus</i>
	PCR product restricted with <i>Hae</i> III gives > 3 bands	<i>F. occidentalis</i>

Discussion

DNA patterns generated by ITS-RFLP have potential as useful tool in the identification of pest thrips species. Furthermore, we have successfully tested this method on instar stages, for which morphological keys are scanty or difficult to utilize, and a similar system has been applied to identify single eggs (Moritz *et al.*, 2004). The technique might improve the efficacy of pest quarantine diagnoses. The identification to species level of larval stages is usually avoided by quarantine workers. Larval thrips are often mistaken for Collembola, or adults are confused with Staphylinid beetles (Vierbergen, 1995). Such misidentifications might be important in the case of thrips species that have a predatory behaviour, such as *Karnyothrips* or *Scolothrips* species (Marullo and De Grazia, 2013). Therefore, a need exists for a system of identifying adult and larval stages of pest thrips species. Recently, the molecular technique known as loop-mediated isothermal amplification (LAMP) was employed to discriminate the presence of DNA of *T. tabaci* from other species, using material from different stages of development (Fekrat *et al.*, 2015) and to detect *Thrips palmi* Karny, a pest species under quarantine regulation in the European Union. These studies highlighted the sensitivity and the usefulness of LAMP to identify all the stages of the life cycle, including the preimaginal stage, which otherwise could be only differentiated by expert taxonomists (Przybylska *et al.*, 2015).

In this study, two molecular keys have been presented, one with six of the seven species investigated and the other was developed to include *H. haemorrhoidalis* that was not represented in the first key. Production of a molecular key that involves members of different genera can be difficult, and certain constraints have to be considered. In the present work analysis, a few technical problems were encountered, such as the absence of results for *H. haemorrhoidalis* using the primer pair 18SMP-28SMP in order to amplify the ITS regions. Also, the primer pair CS249-CS250 was ineffective for *T. tabaci*, *S. citri* and *S. aurantii*. A further weakness of the ITS-RFLP method is that it still requires morphological identification of specimens, as similarly asserted by Rugman-Jones *et al.* (2006) for the most important pest species of the genus *Scirtothrips*. This aspect might be an hindrance for non-specialist taxonomists not familiar with thrips' dichotomous and electronic keys (Hoddle *et al.*, 2012; ThripsWiki, 2014).

Another critical consideration is that studies based on PCR are able to detect only a few species at a time. It is essential to develop a more efficient method for simultaneous screening of mass samples. Possibly, DNA barcoding methods in combination with the mitochondrial COI gene could be a future way to molecular studies (Mehle and Trdan, 2012). An integrated approach of morphology and DNA barcoding for the invasive pest thrips, *Thrips parvispinus* (Karny) is proposed from Tyagi *et al.*, 2015, in order to support how DNA barcoding data may be immense use for accurate species identification and phylogenetic analysis (Mound *et al.*, 2010).

Moreover, recent studies (Chung *et al.*, 2011; Lee *et al.*, 2013; Yeh *et al.*, 2015) report the microarray assay as a valid method for insect pest identification. Particularly, Yeh *et al.* (2015), using DNA microarray based on species-specific primers referring to ITS1 sequences of 15 agriculturally important thrips, provided an efficient tool for the simultaneous identification and monitoring of a number of thrips species. In conclusion, the availability of a molecular key could be extremely useful in the identification of species. The advantage of such an identification system over morphology-based taxonomic methods and other molecular techniques is that it is quick, specific, requires only basic laboratory skills, and can be performed with DNA extracted from single individuals, which can be preserved, mounted on slides and used for future reference. On balance, molecular methods for identification of thrips species represent an important possibility when classical morphological methods are difficult, time consuming or visually impossible.

Acknowledgements

The Authors are grateful to Laurence Mound, Australian National Insect Collection, CSIRO, Canberra (Australia) for the text revision. The scientific work, in particular the stage of A. De Grazia at Molecular Laboratory of the Biology Institute, Halle-Wittenberg University (Germany), has been carried on through the financial support of GEISCA Project 2010/2011 (MIUR, Ministero dell'Università e della Ricerca Scientifica e Tecnologica, Roma).

References

- BRUNNER P. C., FREY J. E., 2010.- Habitat-specific population structure in native western flower thrips *Frankliniella occidentalis* (Insecta, Thysanoptera).- *Journal of Evolutionary Biology*, 23: 797-804.
- BRUNNER P. C., FLEMING C., FREY J. E., 2002.- A molecular identification key for economically important thrips species (Thysanoptera: Thripidae) using direct sequencing and a PCR-RFLP-based approach.- *Agricultural and Forest Entomology*, 4: 127-136.
- BRUNNER P. C., CHATZIVASSILIOU E. K., KATIS N. I., FREY J. E., 2004.- Host-associated genetic differentiation in *Thrips tabaci* (Insecta: Thysanoptera), as determined from mtDNA sequence data.- *Heredity*, 93: 364-370.
- CHUNG I. H., KANG S., KIM Y. R., KIM J. H., JUNG J. W., LEE S., LEE S. H., 2011.- Development of a low density DNA microarray for diagnosis of target-site mutations of pyrethroid and organophosphate resistance mutations in the whitefly *Bemisia tabaci*.- *Pest Management Science*, 67: 1541-1548.
- FARRIS R. E., RUIZ-ARCE R., CIOMPERLIK M., VASQUEZ J. D., DE LEON R., 2010.- Development of a ribosomal DNA ITS2 marker for the identification of the thrips, *Scirtothrips dorsalis*.- *Journal of Insect Science*, 10: 1-15.
- FEKRAT L., ZAKI AGHI M., TAHAN V., 2015.- Application of the LAMP Assay as a diagnostic technique for rapid identification of *Thrips tabaci* (Thysanoptera:Thripidae).- *Journal of Economic Entomology*, 108 (3): 1337-1343.
- HILLIS D. M., HUELSENBECK J. P., CUNNINGHAM C. W., 1994.- Application and accuracy of molecular phylogenies.- *Science*, 264: 671-677.

- HODDLE M. S., HERATY J. M., RUGMAN-JONES P. F., MOUND L. A., STOUTHAMER R., 2008.- Relationships among species of *Scirtothrips* (Thysanoptera: Thripidae, Thripinae) using molecular and morphological data.- *Annals of the Entomological Society of America*, 101: 491-500.
- HODDLE M. S., MOUND L. A., PARIS D., 2012.- *Thrips of California*.- [online] URL: <http://www.lucidcentral.com>
- JACOBSON A. L., BOOTH W., VARGO E. I., KENNEDY G. G., 2013.- *Thrips tabaci* population genetic structure and polyploidy in relation to competency as a vector of Tomato spotted Wilt Virus.- *PLoS One*, 8 (1): e54484.
- KOX L. F. F., VAN DEN BELD H. E., ZIJLSTRA C., VIERBEGEN G., 2005.- Realtime PCR assay for the identification of *Thrips palmi*.- *EPP0 Bulletin*, 35: 141-148.
- KUCHARCZYK H., 2010.- *Comparative morphology of the second larval instar of the Thrips genus species (Thysanoptera: Thripidae) occurring in Poland*.- Wydawnictwo Mantis, Olsztyn, Poland.
- LEE W. S., CHOI H., KANG J., KIM J. H., LEE S. H., LEE S., HWANG S. Y., 2013.- Development of a DNA microarray for species identification of quarantine aphids.- *Pest Management Science*, 69: 1399-1406.
- MARULLO R., DE GRAZIA A., 2012.- Thripidae, pp. 109-118. In: *Integrated control of citrus pests in the Mediterranean region* (VACANTE V., GERSON U., Eds).- Bentham Science Publishers, United Arab Emirates.
- MARULLO R., DE GRAZIA A., 2013.- Territorial distribution, classification and relationships amongst Italian Thysanoptera.- *Bulletin of Insectology*, 66 (1): 127-134.
- MEHLE N., TRDAN S., 2012.- Traditional and modern methods for the identification of thrips (Thysanoptera) species.- *Journal of Pest Science*, 85: 179-190.
- MORITZ G., 2010.- *Database ThripsNet*.- [online] URL: <http://moritz.zoologie.uni-halle.de/>
- MORITZ G., DELKER C., PAULSEN M., MOUND L. A., BURGERMEISTER W., 2000.- Modern methods for identification of Thysanoptera.- *EPP0 Bulletin*, 30: 591-593.
- MORITZ G., MOUND L. A., MORRIS D. C., GOLDARAZENA A., 2004.- *Pest thrips of the world*. CD-ROM.- The University of Queensland, Brisbane, Australia.
- MORITZ G., O'DONNELL C., PARRELLA M., 2009.- *Pest thrips of North America*.- CBIT, The University of Queensland, Brisbane, Australia.
- MORITZ G., BRANDT S., SSERWAGI P., MYAMBA A., WAIGANJO M., SUBRAMANIAN S., 2010.- Entwicklung eines LucID 3.5-Identifikations- und Informationssystems für Schad-Thrips in Ostafrika. - *Julius-Kühn-Archiv*, 428: 250-251.
- MORITZ G., BRANDT S., TRIAPTSYN S., SUBRAMANIAN S., 2013.- Identification and information tools for pest thrips in East Africa.- QBIT, The University of Queensland, Brisbane, Australia.
- MORSE J. G., HODDLE M. S., 2006.- Invasion biology of thrips.- *Annual Review of Entomology*, 51: 67-89.
- MOUND L. A., WHEELER G. S., WILLIAMS D. A., 2010.- Resolving cryptic species with morphology and DNA; thrips as a potential biocontrol agent of Brazilian pepper tree, with a new species and overview of *Pseudophilothrips* (Thysanoptera).- *Zootaxa*, 2432: 59-68.
- PRZYBYLSKA A., FIEDLER Z., KUCHARCZYK H., OBRZEPALSKA-STEPLEWSKA A., 2015.- Detection of the quarantine species *Thrips palmi* by loop-mediated isothermal amplification.- *PLoS ONE*, 10 (3): e0122033.
- RUGMAN-JONES P. F., HODDLE M. S., MOUND L. A., STOUTHAMER R., 2006.- Molecular identification key for pest species of *Scirtothrips* (Thysanoptera:Thripidae).- *Journal of Economic Entomology*, 99 (5): 1813-1819.
- THRIPSWIKI, 2014.- *ThripsWiki - providing information on the World's thrips*.- [online] URL: <http://thrips.info/wiki/>
- TIMM A. E., STILLER M., FREY J. E., 2008.- A molecular identification key for economically important thrips species (Thysanoptera:Thripidae) in southern Africa.- *African Entomology*, 16 (1): 68-75.
- TODA S., KOMAZAKI S., 2002.- Identification of thrips species (Thysanoptera:Thripidae) on Japanese fruit trees by polymerase chain reaction and restriction fragment length polymorphism of the ribosomal ITS2 region.- *Bulletin of Entomological Research*, 92: 359-363.
- TYAGI K., KUMAR V., SINGHA D., CHAKRABORTY R., 2015.- Morphological and DNA barcoding evidence for invasive pest thrips, *Thrips parvispinus* (Thripidae: Thysanoptera), newly recoded from India.- *Journal of Insect Science*, 15 (1): 105.
- VIERBEGEN G., 1995.- International movement, detection and quarantine of Thysanoptera pests, pp. 119-132. In: *Thrips biology and management* (PARKER B. L., SKINNER M., LEWIS T., Eds).- Plenum Press, New York, USA.
- VIERBEGEN G. B., KUCHARCZYK H., KIRK W. D. J., 2010.- A key to the second instar larvae of the Thripidae of the Western Palaearctic region (Thysanoptera).- *Tijdschrift voor Entomologie*, 153: 99-160.
- YEH W. B., TSENG M. J., CHANG N. T., WU S. Y., TSAI Y. S., 2015.- Agronomically important thrips: development of species-specific primers in multiplex PCR and microarray assay using internal transcribed spacer 1 (ITS1) sequences for identification.- *Bulletin of Entomological Research*, 105: 52-59.

Authors' addresses: Alessandra DE GRAZIA (corresponding author, alessandra.degrazia@unirc.it), Rita MARULLO, Dipartimento di Agraria, Università degli Studi Mediterranea di Reggio Calabria, Località Feo di Vito I-89060 Reggio Calabria, Italy; Gerald MORITZ, Institute of Biology, Martin-Luther University, Halle-Wittenberg, Faculty of Natural Sciences 1, Heinrich-Damerow-Str.4, 06120 Halle, Germany.

Received August 31, 2015. Accepted November 18, 2015.