



Research paper

Synchronisms and correlations of spring phenology between apical and lateral meristems in two boreal conifers

Serena Antonucci^{1,2,5}, Sergio Rossi^{2,3}, Annie Deslauriers², Fabio Lombardi¹, Marco Marchetti¹ and Roberto Tognetti^{1,4}

¹Dipartimento di Bioscienze e Territorio, Università degli Studi del Molise, Pesche (IS) 86090, Italy; ²Département des Sciences Fondamentales, Université du Québec à Chicoutimi, Chicoutimi (QC), G7HSB1, Canada; ³Key Laboratory of Vegetation Restoration and Management of Degraded Ecosystems, Provincial Key Laboratory of Applied Botany South China Botanical Garden, Chinese Academy of Sciences, Guangzhou, China; ⁴The EFI Project Centre on Mountain Forests (MOUNTFOR), Edmund Mach Foundation, San Michele all'Adige (TN) 38010, Italy; ⁵Corresponding author (s.antonucci1@studenti.unimol.it)

Received February 5, 2015; accepted August 1, 2015; published online September 15, 2015; handling Editor Annikki Mäkelä

Phenological synchronisms between apical and lateral meristems could clarify some aspects related to the physiological relationships among the different organs of trees. This study correlated the phenological phases of bud development and xylem differentiation during spring 2010–14 in balsam fir (*Abies balsamea* Mill.) and black spruce [*Picea mariana* Mill. (BSP)] of the Monts-Valin National Park (Quebec, Canada) by testing the hypothesis that bud development occurs after the reactivation of xylem growth. From May to September, we conducted weekly monitoring of xylem differentiation using microcores and bud development with direct observations on terminal branches. Synchronism between the beginning of bud development and xylem differentiation was found in both species with significant correlations between the phases of bud and xylem phenology. Degree-day sum was more appropriate in assessing the date of bud growth resumption, while thermal thresholds were more suitable for cambium phenology. Our results provide new knowledge on the dynamics of spring phenology and novel information on the synchronisms between two meristems in coniferous trees. The study demonstrates the importance of precisely defining the phases of bud development in order to correctly analyse the relationships with xylem phenology.

Keywords: balsam fir, black spruce, bud break, cambium, temperature, xylogenesis.

Introduction

Phenology is one of the most responsive plant traits driven by climate (Badeck et al. 2004), making trees reliable indicators of climate change (Donnelly et al. 2004). Phenological observations record earlier events in terms of bud break, shooting, leafing and flowering, which may be related to the recent spring warming and extended growing seasons (Chuine and Beaubien 2001, Menzel et al. 2006). In spring, bud development is mainly regulated by temperature (Sarvas 1972), although other interacting factors such as chilling and photoperiod can play an important role (Heide 1993, Häkkinen et al. 1998, Partanen et al. 2001, Rossi et al. 2007). Air temperature is a good predictor of bud development and shoot elongation, once ontogenetic development begins (Schwalm and Ek 2001).

Bud swelling is a phenological event of growth resumption composed of sequential events during which the embryonic shoots and leaves rapidly proliferate and emerge from the bud scales. Generally, studies on apical meristem have identified bud swelling with the phase of leaf unfolding (Moser et al. 2010, Cuny et al. 2012, Takahashi and Koike 2014), while few studies have taken into account all bud development phases (Dhont et al. 2010, Huang et al. 2014, Rossi and Bousquet 2014).

As with bud development, stem growth also occurs through different phases. Secondary xylem production is a complex process that derives from cell periclinal and cataclinal division in the vascular cambium. The differentiation of annual rings in conifers involves the formation of tracheids that pass through several phases before reaching their final form (Deslauriers et al. 2003).

Several studies have revealed an early onset of cambium activity and xylem cell production as well as a longer duration of xylem formation, and this has often been related to the influences of the ongoing climate change (Huang et al. 2011, Boulouf-Lugo et al. 2012). The activity of meristems involves a number of biochemical processes resulting in a sequence of phases of development and maturation. These phases are identified according to the morphological or anatomical changes and can last from a few days, in primary meristems, to several weeks, in the case of secondary meristems (Rossi et al. 2013, Rossi 2015).

Aloni (2015) showed how leaf development (related to the primary meristems) and biomass accumulation (related to the secondary, or lateral, meristems) are regulated by a number of environmental factors. These factors influence not only wood production and its biochemical traits but also the amount and differentiation of vascular cells. Moreover, physiological factors, such as carbohydrate availability and hormone distribution, could determine the dynamics of wood formation (Uggla et al. 1998, 2001, Rossi et al. 2008). Studies on bud and cambium phenology have revealed different patterns and links between primary and secondary meristems. In conifers, the last phase of bud development, often identified as bud break, has been found to occur either before or after the onset of xylem cell production (Ladefoged 1952, O'Reilly and Owens 1989, Rensing and Owens 1994, Rossi et al. 2009, Cuny et al. 2012), while in diffuse porous species, cambial reactivation takes place immediately after bud break (Schmitt et al. 2000, Čufar et al. 2008). Thus, the relationships between apical and lateral meristems are still unclear, at least in conifer species. Understanding these relationships is crucial for identifying the timings of carbon uptake and seasonal exchanges of water, nutrients and gases between the land surface and atmosphere (Chen and Xu 2012). In order to elucidate the role of meristems as indicators of the effects of climate change, in particular warming temperature, the relationships between primary and secondary growth need to be defined for a mechanistic understanding of the growth dynamics in all tree organs (Huang et al. 2014).

There are still few ecophysiological studies aiming to explain the relationships between bud and cambial phenology and their synchronism over the growing season (Deslauriers et al. 2009, Huang et al. 2014). Characterization of the relationships between environmental factors, particularly temperature, and meristem activity is crucial to understanding the fate and differentiation of meristem cells and their coordination under environmental stresses. Heat accumulation above a threshold temperature, commonly expressed in degree-days, has often been used to predict or verify the effect of temperature on biological processes (Baskerville and Emin 1969). In this study, having monitored and compared the timings of bud and xylem development in two conifer species of the boreal forest, we identified the spring temperature thresholds and cumulated degree-days

required for triggering the first phases of bud and cambium development. The study was conducted during five growing seasons, investigating in detail all the phenological phases for growth resumption of bud and cambium and their climatic drivers, to improve the understanding of seasonal patterns and responses to inter-annual and long-term variation in climate. The hypothesis that the beginning of bud development occurs after the reactivation of xylem growth was tested.

Materials and methods

Study site and tree selection

This study was conducted in the Monts-Valin National Park (Quebec, Canada). Two sites at different altitudes were selected in the black spruce–feather moss bioclimatic domain, Gaspard (GP) (Lat. 48.58° N, Long. 70.88) and Lagacé (LA) (Lat. 48.60° N, Long. 70.83), located at 227 and 900 m above sea level (a.s.l.), respectively (Table 1). The climate in the region is dominated by dry and moderately polar air masses in winter and moist maritime air masses in summer (Sheridan 2002). Absolute minimum temperature reaches -41°C , and mean air temperature ranges between -1.4 and 2.5°C , with annual precipitation higher than 1000 mm. The sites experience between 204 and 256 frost days per year.

Primary and secondary growth was monitored weekly from May to September, during the 2010–14 growing seasons, in balsam fir [*Abies balsamea* (L.) Mill.] and black spruce [*Picea mariana* (Mill.) BPS]. Three trees per species were randomly selected in each site, choosing healthy dominant or co-dominant individuals.

Bud phenology

Two north-facing and two south-facing branches per tree were selected in the bottom part of the canopy. On each branch, the phases of bud development were recorded on terminal buds according to Rossi and Bousquet (2014): (i) open bud, with the scales starting to separate and a pale spot visible at the tip; (ii) elongated bud, with lengthening scales; (iii) swollen bud, with smooth and pale coloured scales but no visible needle; (iv) translucent bud, with needles visible through the scales; (v) split bud, with open scales but needles still clustered; and (vi) exposed shoot, with needles completely emerged from the surrounding scales and spreading outwards. All data were computed in days of the year (DOY), by averaging the data observed in the field.

Table 1. Mean (standard deviation) of the characteristics of the trees in the two study sites.

Species	Site	Stand age (years)	Height (m)	Diameter at breast height (cm)
Spruce	LA	72.7 (20.3)	4.6 (1.04)	14.0 (0.03)
	GP	39.7 (3.7)	14.2 (1.6)	24.7 (1.4)
Fir	LA	25.3 (2.5)	7.6 (0.7)	16.8 (4.0)
	GP	49.7 (3.7)	17.6 (2.6)	32.2 (9.1)

Cambial phenology

Wood microcores (2 mm in diameter) were extracted from the stem at breast height, using a Trephor tool (Rossi et al. 2006a). The samples contained the preceding four to five tree rings and the developing annual layer with the cambial zone and adjacent phloematic tissue. The microcores were placed in Eppendorf microtubes with an ethanol solution (50% water) and stored at 5 °C to avoid tissue deterioration. Microcores were dehydrated with successive immersions in ethanol and d-limonene and then embedded in paraffin according to Rossi et al. (2006a). In order to obtain thin slices, the samples were cut (7 µm in thickness) using a rotary microtome. The sections were stained with cresyl violet acetate (0.16% in water) and examined within 10–15 min under visible and polarized light at 400–500× magnification to differentiate the developing and mature xylem cells.

The radial number of cells was counted along three radial rows, according to the criteria described in Rossi et al. (2006b). Observations under polarized light discriminated the zones of enlarging and cell wall thickening of tracheids. The progress of cell wall lignification was detected with cresyl violet acetate reacting with lignin (Antonova and Shebeko 1981). Lignification was shown by a colour change from violet to blue. The colour change over the whole cell wall revealed the end of lignification and the tracheid reaching maturity (Gričar et al. 2005). The dormant cambium consisted of three to four cells. In spring, when the diameter of the new xylem cells was at least twice that of a cambial cell, the onset of cell division was considered to occur. Xylogenesis was considered to have started when at least one radial file of enlarging cells was observed (Rossi et al. 2008) (Figure 1). In late summer, when no further cell was observed in wall thickening and lignification, xylem formation was considered complete.

Weather stations

In both sites, a standard weather station was installed in a forest gap to measure temperature and precipitation. Data were collected every 15 min during the 5 years of monitoring and recorded as hourly averages by means of CR10X data loggers (Campbell Scientific Corporation, Logan, UT, USA). Daily mean values were calculated with the time series obtained from the 24 measurements per day.

Data analysis

Timings of bud phenology were compared between species, DOY and years with analysis of covariance (ANCOVA). Phases of bud and xylem phenology were compared using analysis of variance (ANOVA), while the Tukey test was applied for multiple comparisons. The correlations between phenological phases were tested through Pearson correlations.

Degree-days sum (DDS) at the onset of bud and cambium phenology were calculated as a sum of the positive differences between mean daily air temperature and the threshold value




Figure 1. Beginning of xylem differentiation with dividing cambium and cells in enlargement phases (DOY 160, Site LA). Ph, phloem; Cz, cambial zone; Ec, enlarging cells.

0 °C (Man and Lu 2010). This threshold was used for calculating DDS according to Man and Lu (2010), who suggested 0 °C as an optimal base temperature for black spruce. Thermal thresholds (THR) for bud and xylem phenology were assessed using logistic regressions to calculate the probability of active meristem growth at a given measured daily temperature. Binary responses were coded as non-active (value 0) or active (value 1) growth, and temperature thresholds were calculated when the probability of active meristem growth was 0.5 (Rossi et al. 2008). Coefficient of variation was used for comparing DDS and THR, given that these measurements take non-negative values. All statistical analyses were conducted using JMP (Version 11, SAS Institute, Inc., Cary, NC, USA).

Results

Bud phenology

The phases of bud phenology occurred earlier in fir than in spruce at both altitudes, though showing different trends in the investigated years. The period of bud development was shorter in fir than in spruce and longer in 2010 and 2013 than in other years. In 2014, the trend of the two species was similar at lower elevation and almost identical at higher elevation. Occasionally, some intermediate phases were not recorded, especially at low elevation, because they occurred within the same sampling week. The first bud development phase occurred in the second half of May. The two species showed a week of difference for Phase 1, occurring on average on DOY 138 and 145 in fir and spruce, respectively. Phase 6 occurred ~4 and 5 weeks after Phase 1 in fir and spruce, respectively (Figure 2). In LA (900 m




Figure 2. Average of days of occurrence of the phases of bud development in spruce and fir observed during 2010–14 at two altitudes (1, open bud; 2, elongated bud; 3, swollen bud; 4, translucent bud; 5, split bud; 6, exposed shoot). Some phases are missing because they were not observed during the weekly monitoring.

a.s.l.), bud phenology started 2 weeks later than in GP (227 m a.s.l.). ANCOVA model was highly significant ($P < 0.001$) for both altitudes, with $F = 435.77$ and 431.51 for 227 and 900 m a.s.l., respectively (Table 2). As expected, the bud development phases occurred on different DOY. The two species showed different phenological development behaviour and the years considered differed from each other. All the factors considered were statistically significant with $P < 0.001$.

Xylem phenology

In the analysed cross section, cambial cells were characterized by thin cell walls and small radial diameters. At the onset of cambial activity, the cambial zone began to widen rapidly as the number of cells increased, revealing that cell division had started. During cell enlargement, the tracheids were composed of protoplast still enclosed in the thin primary wall, but with radial diameter at least twice that of a cambial cell.

The start of xylem cells production was similar and within the same week in the two species (Figure 3). As expected, xylem growth and differentiation started first at lower elevation, at the beginning of May. At higher elevation, cambial resumption occurred between the end of May and beginning of June. The first mature cells were recorded 22 and 24 days after the start of enlargement in fir and spruce, respectively. The difference in days of mature cell production was greater in LA than in GP. In 2010 and 2013, too few microcores were available to determine the date of cell maturation with precision.

Comparing xylem and bud phenology

The correlations between the date of bud and xylem phenology were highly significant (Table 3). A strong association was detected between the open bud (Phase 1) and date of the first enlargement for both fir ($r = 0.91$) and spruce ($r = 0.90$) ($P < 0.001$). The first mature xylem cell and the stage of exposed shoot showed a significant correlation in fir ($r = 0.98$, $P < 0.01$),

Table 2. F -value and resulting probability of the ANCOVA model for the phases of bud phenology.

Altitude (m a.s.l.)	Model		Factor	Statistics	
	F -value	R^2		F -value	P
900	431.51	0.67	Year	53.25	<0.0001
			DOY	1281.49	<0.0001
			Species	93.64	<0.0001
227	435.77	0.66	Year	49.44	<0.0001
			DOY	1266.52	<0.0001
			Species	156.93	<0.0001

but not in spruce ($r = 0.57$, $P > 0.05$), meaning that the variation between both stages is not always synchronous for spruce.

The multiple comparison of phenological events, performed with ANOVA ($P < 0.001$), showed that the dates (DOY) of first cell enlargement and open bud phases were not statistically different in fir. In addition, DOY of the first mature cell and exposed shoot phases did not show statistical difference. In spruce, only the first phases of xylem differentiation and bud development at higher elevation showed no statistical differences. On the contrary, differences were observed at lower elevation between all the analysed phases (Figure 4).

Degree-days sum and thermal threshold

Open bud occurred in fir and spruce when 230 ± 44 (mean \pm standard deviation) and 284 ± 33 degree-days had accumulated, respectively. The occurrence of the first enlargement cell was observed at lower DDS than the open bud phase, with 217 ± 43 degree-days in spruce and 201 ± 37 degree-days in fir. Degree-days sum in the exposed shoot phase were higher than that in the first mature cell phase (Figure 5). The mean THR were estimated at 7.9 ± 1.2 °C for the first enlarging cell phase in spruce and 7.4 ± 1.5 °C in fir. The open bud phase had a THR of $\sim 9.1 \pm 1.2$ °C in spruce and 7.7 ± 1.8 °C in fir (Figure 5). The standard




Figure 3. Days of occurrence of the phases of xylem differentiation in spruce and fir observed during 2010–14 (FE, first enlarging cell; FWT, first wall-thickening cell; FM, first mature cell). Error bars indicate standard deviation between trees.

Table 3. Correlation matrix between dates of bud and cambium phenology during 2010–14 in spruce (white background) and fir (grey background). One, two and three asterisks correspond to a $P < 0.05$, < 0.01 and < 0.001 , respectively.

	Open bud	Exposed shoot	First enlarging xylem cell	First mature xylem cell
Open bud	–	0.82**	0.90***	0.81*
Exposed shoot	0.82**	–	0.77*	0.57
First enlarging xylem cell	0.91***	0.78**	–	0.88*
First mature xylem cell	0.93*	0.98**	0.89*	–

deviations of DDS and THR were similar between the first phases of xylem differentiation and the first phases of bud phenology, in both species. For spruce, the coefficient of variation showed higher values for the first enlarging cell phase than for the other phases, though the exposed shoot phase showed higher values than the open bud and first mature cell phases. The same trend was observed for the coefficient of variation of THR. In fir, the highest coefficient of variation was found in the open bud phase, which occurred for both DDS and THR. The coefficient of variation was smaller for DDS than for THR for the open bud phase in both species. Finally, the first enlarging cell phase showed higher coefficient of variation for DDS than for THR (Table 4).

Discussion

Synchronisms and correlations between meristems

In this study, we monitored all the phenological phases of bud development and xylem differentiation in black spruce and balsam fir during five growing seasons, investigating the possible relationships between bud development and xylem phenology. This is the first study, to our knowledge, that examines long-term chronologies of the two meristems in these boreal species. The hypothesis that the beginning of bud development in conifers occurs after xylem reactivation was only partially accepted. This result was related to the criteria defined a priori in this study.




Figure 4. Bud and cambium phenology observed during 2010–14 for spruce and fir (error bars indicate standard deviation between years). Bars with the same letters are not statistically different ($P > 0.05$).

Generally, bud break has been represented by the last phases of bud development, leaf unfolding (Rossi et al. 2009, Moser et al. 2010, Cuny et al. 2012, Takahashi and Koike 2014). However, the bud growth resumption begins 1 month before or even earlier, when the scales start to separate following bud swelling. Although the phases of bud phenology are well known and documented (Dhont et al. 2010, Sutinen et al. 2012, Rossi and Bousquet 2014), we are not aware of any studies that have




Figure 5. Degree-days sum and thermal threshold assessed for bud and cambium phenology (error bars indicate standard deviation between years).

Table 4. Mean \pm standard deviations and coefficient of variation (expressed as a percentage in brackets) for DDS and THR for bud and cambium phenology in spruce and fir.

	DDS	THR
Spruce		
First enlarging cell	217 \pm 43 (19.9)	7.9 \pm 1.2 (14.9)
Open bud	230 \pm 44 (11.7)	9.1 \pm 1.2 (12.9)
First mature cell	468 \pm 50 (10.8)	11.8 \pm 1.1 (9.3)
Exposed shoot	795 \pm 103 (13.0)	13 \pm 1.7 (13.1)
Fir		
First enlarging cell	201 \pm 37 (18.4)	7.4 \pm 1.5 (18.7)
Open bud	284 \pm 33 (19.2)	7.7 \pm 1.8 (22.8)
First mature cell	420 \pm 68 (16.3)	11.3 \pm 1.3 (11.2)
Exposed shoot	585 \pm 98 (16.8)	11.3 \pm 1.1 (9.4)

considered their relationships with xylem phenology in detail. Indeed, according to the general definition of bud swelling, apical phenology can occur before or after cambial resumption. However, from a biological point of view, the phase of bud swelling, occurring when the scales start to separate and a pale spot is visible at the tip (Rossi and Bousquet 2014), and not that of leaf unfolding, seems to be the most appropriate to represent the break of bud dormancy, as also suggested by Sutinen et al. (2012). As a consequence, we are confident that the beginning of xylem differentiation was synchronous with the beginning of bud development in both the studied species, although no causal relationship between the two meristems can be deduced by this monitoring. Temperature is known to control the phenological events in spring, thus the synchronisms observed between meristems could be the results of a response of the meristems to a common driving factor (e.g., high temperature and available liquid water in early spring may promote photosynthetic activity and carbohydrate storage).

Recent studies have demonstrated that localized heating anticipated cambium reactivation and earlywood vessel formation in seedlings of a broadleaved deciduous species, in comparison with non-heated seedlings, and occurred before bud break (Kudo

et al. 2014). Through artificially heated and cooled stems, several other authors were able to induce an earlier break of cambial dormancy and higher rates of cambial cell production (during heating) and a delay in cambial reactivation and lower rates of cambial cell production (during cooling) (Oribe and Kubo 1997, Oribe et al. 2001, 2003, Gričar et al. 2006, Begum et al. 2007, 2010, 2012, 2013). Nevertheless, our results showed a strong coupling between phases of bud development and those of xylem phenology in spring, under natural growing conditions in the boreal forest. This means that both phenological patterns varied in a similar manner along the time axis, i.e., when the phase of open bud occurred earlier, the first enlarging cell was also observed earlier. The synchronization between the last two phases, exposed shoot in bud phenology and mature cell in xylem differentiation, was unclear, because significant correlations between the last phases of the two meristems were observed only in fir. Variability in xylem and phloem phenology among years and between species might be determined by exogenous factors (contrasting temperatures prevailing at the start of the growing season and water availability) and intrinsic factors (gene expression and hormonal signals) (Swidrak et al. 2014). Correlation between primary and secondary growth processes over the growing season would indicate optimal mechanisms to simultaneously allocate photosynthetic products and stored non-structural carbon for the growth of the different organs in trees (Huang et al. 2014).

Synchronisms between species

Phenological differences were observed between the two species. Bud phenology began between mid-May and late June. Fir began earlier than spruce, which might be attributed to a greater sensitivity to spring temperatures, fir being a microthermal species in comparison with spruce. This behaviour could also be explained by genetic differences related to the species-specific capacity to perceive exogenous signals through hormones (Aloni 2001). Jones and Cregg (2006) studied the timings of bud

break in different types of North American firs, showing that the date varied with location, species and year. As expected, in all the years considered, open bud occurred first in trees at lower elevation. This was attributed to earlier spring warming at lower elevation (Moser et al. 2010). Several studies suggest that the accumulation of lipids in the cortex area of the buds would trigger a series of events connected to the onset of bud development (Alaudinova and Mironov 2010, Sutinen et al. 2012). However, studies on xylem phenology showed similar timing between species. Thibeault-Martel et al. (2008) showed no differences in onset of cell enlargement for *A. balsamea* and *P. mariana*, whereas the same cambium activity date was observed for fir and pine trees in temperate environments (Cuny et al. 2012, Gričar et al. 2014).

By comparing bud phenology, it was possible to disentangle the differential responses to climate patterns in these co-occurring species. While conservative responses (low sensitivity to warming temperature) may opportunistically minimize the risk of late (spring) or early (autumn) frost at little cost (Guy 2014), with greater climate warming, phenologically flexible species may advantageously occupy new early or late temporal niches within a short time (Wolkovich et al. 2013). Black spruce has a current competitive advantage over balsam fir in coniferous forests of the boreal shield due to a greater tolerance to cooler temperatures and soil waterlogging (Messaoud et al. 2014). However, warmer temperatures will potentially cause asynchrony in biological activities, disrupting ecological interactions (e.g., between these conifers and spruce budworm and/or fire). A longer period for primary and secondary tree growth with warming temperature could advance black spruce phenology, making this species a more suitable host for spruce budworm.

Temperature of growth

Accumulated temperature in spring is important for bud development and the onset of radial growth (Tadaki et al. 1994). Degree-days sum increased in the order of first cell enlargement, open bud, first mature cell and exposed shoot phases, as did the thermal threshold. In our study, we used the coefficient of variation for comparing the methods generally applied in agriculture and forestry to evaluate the influence of temperature on phenological phases. The coefficient of variation for the open bud phase was lower in DDS than in THR. The opposite occurred for the first cell enlargement phase. This coefficient indicated that DDS better explained the first phase of bud phenology, whereas THR were more suitable for the first phase of xylem differentiation. The degree-days, therefore, appeared to be the more appropriate monitoring technique for apical meristems, while the thermal threshold was more advisable for cambium phenology. Rossi et al. (2008) determined common critical temperature thresholds for xylogenesis in conifers at different latitudes and altitudes in Europe and Canada, with average daily temperature values of 8–9 °C. Analyses at a wider geographical

scale comparing several conifer species have confirmed convergence of the thermal threshold around specific critical values, demonstrating the existence of a precise thermal limit in radial growth and tree ring formation (Deslauriers et al. 2008, Rossi et al. 2008). Prislán et al. (2013) observed that the onset of cambial cell division in *Fagus sylvatica* L. was associated with an extended period of preconditioning that varied with site, rather than with a threshold temperature. Patterns of cambial activity and climate thresholds have been found to vary among sites in *Picea abies* (Gričar et al. 2014), suggesting that local adaptation may play a role in coupling processes of wood formation and foliage renewal within populations. Gričar et al. (2014) showed site-specific amounts of accumulated heat units at the onset of cambial cell production. In contrast, Schmitt et al. (2004) reported that growing degree-days for the onset of cambial activity in *Pinus sylvestris* were similar along a latitudinal gradient in the boreal forests of Finland. Instead, DDS have been used to predict the growth stages of major crops and trees, particularly in tracking bud development (Wang 1960, Førland et al. 2004, Jones and Cregg 2006). It should be pointed out that longer and warmer summers at higher elevations and northern latitudes might result in disproportional resource requirements for these species, namely water.

Conclusions

To our knowledge, this is the first study that examines the synchronisms between the phases of growth resumption in buds and xylem, using chronologies of 5 years of data collected from two species and two sites. Detailed phases of bud break and xylem cell differentiation showed a significant synchronism in the timings of activation between the two meristems. Considering the bud swelling as break of dormancy and not exposed shoot, we found that the beginning of bud development occurred in the same period as xylem differentiation in both species. This demonstrates the importance of considering all phases of bud break in order to correctly evaluate the relationships with xylem phenology. Data from the two meristems were strongly correlated to one another, although fir dormancy release was earlier than spruce. This study allowed the dynamics of spring phenology to be explored in detail, providing novel information about the potential physiological correlation between meristems and filling an important gap in the knowledge on the growth synchronisms in conifer species. A straightforward quantification of the dynamics of growth and response of phenology to climatic drivers is essential to link plant-level events of life cycles with the environmental factors occurring at a short time scale. The presence of synchronism in leaf and cambial activity in these conifers may be conveniently used to relate field (leaf phenology and wood production) and satellite (vegetation indices) data and explain seasonal variability in boreal forest productivity.

Acknowledgments

The authors thank C. Pelletier and A. Ferland-Roy from the Parc National des Monts-Valin, C. Soucy and G. Savard for their help with data collection and A. Garside for editing the English text. This work was encouraged by activities conducted within the COST FP1 106 (STReESS—Studying Tree Responses to extreme Events: a Synthesis) network.

Conflict of interest

None declared.

Funding

This project was funded by the Natural Sciences and Engineering Research Council of Canada (NSERC).

References

- Alaudinova EV, Mironov PV (2010) Lipids of the meristems of the main coniferous edificators from central Siberia under low-temperature adaptation: 2. Features of the fatty acid metabolism of phospholipids from winter meristems of *Larix sibirica* Ledeb., *Picea obovata* L., and *Pinus sylvestris* L. *Russ J Bioorg Chem* 36:872–876.
- Aloni R (2001) Foliar and axial aspects of vascular differentiation: hypotheses and evidence. *J Plant Growth Regul* 20:22–34.
- Aloni R (2015) Ecophysiological implications of vascular differentiation and plant evolution. *Trees* 29:1–16.
- Antonova G, Shebeko V (1981) Applying cresyl violet in studying wood formation. *Khim Drev* 4:102–105.
- Badeck FW, Bondeau A, Böttcher K, Doktor D, Lucht W, Schaber J, Sitch S (2004) Responses of spring phenology to climate change. *New Phytol* 162:295–309.
- Baskerville GL, Emin P (1969) Rapid estimation of heat accumulation from maximum and minimum temperatures. *Ecology* 50:514–517.
- Begum S, Nakaba S, Oribe Y, Kubo T, Funada R (2007) Induction of cambial reactivation by localized heating in a deciduous hardwood hybrid poplar (*Populus sieboldii* × *P. grandidentata*). *Ann Bot* 100:439–447.
- Begum S, Nakaba S, Oribe Y, Kubo T, Funada R (2010) Changes in the localization and levels of starch and lipids in cambium and phloem during cambial reactivation by artificial heating of main stems of *Cryptomeria japonica* trees. *Ann Bot* 106:885–895.
- Begum S, Nakaba S, Yamagishi Y, Yamane K, Islam MA, Oribe Y, Ko JH, Jin HO, Funada R (2012) A rapid decrease in temperature induces late-wood formation in artificially reactivated cambium of conifer stems. *Ann Bot* 110:875–885.
- Begum S, Nakaba S, Yamagishi Y, Oribe Y, Funada R (2013) Regulation of cambial activity in relation to environmental conditions: understanding the role of temperature in wood formation of trees. *Physiol Plant* 147:46–54.
- Boulouf-Lugo J, Deslauriers A, Rossi S (2012) Duration of xylogenesis in black spruce lengthened between 1950 and 2010. *Ann Bot* 110:1099–1108.
- Chen X, Xu L (2012) Temperature controls on the spatial pattern of tree phenology in China's temperate zone. *Agric For Meteorol* 154–155:195–202.
- Chuine I, Beaubien EG (2001) Phenology is a major determinant of tree species range. *Ecol Lett* 4:500–510.
- Čufar K, Prislán P, De Luis M, Gričar J (2008) Tree-ring variation, wood formation and phenology of beech (*Fagus sylvatica*) from a representative site in Slovenia, SE Central Europe. *Trees* 22:749–758.
- Cuny HE, Rathgeber CB, Lebourgeois F, Fortin M, Fournier M (2012) Life strategies in intra-annual dynamics of wood formation: example of three conifer species in a temperate forest in north-east France. *Tree Physiol* 32:612–625.
- Deslauriers A, Morin H, Begin Y (2003) Cellular phenology of annual ring formation of *Abies balsamea* in the Quebec boreal forest (Canada). *Can J For Res* 33:190–200.
- Deslauriers A, Rossi S, Anfodillo T, Saracino A (2008) Cambial phenology, wood formation and temperature thresholds in two contrasting years at high altitude in Southern Italy. *Tree Physiol* 28:863–871.
- Deslauriers A, Giovannelli A, Rossi S, Castro G, Fragnelli G, Traversi L (2009) Intra-annual cambial activity and carbon availability in stem of poplar. *Tree Physiol* 29:1223–1235.
- Dhont C, Sylvestre P, Gros-Louis M, Isabel N (2010) Field guide for identifying apical bud break and bud formation stages in white spruce. Natural Resources Canada, Quebec.
- Donnelly A, Jones M, Sweeney J (2004) A review of indicators of climate change for use in Ireland. *Int J Biometeorol* 49:1–12.
- Førland EJ, Skaugen TE, Benestad RE, Hanssen-Bauer I, Tveito OE (2004) Variations in thermal growing, heating, and freezing indices in the Nordic Arctic, 1900–2050. *Arct Antarct Alp Res* 36:347–356.
- Gričar J, Čufar K, Oven P, Schmitt U (2005) Differentiation of terminal late-wood tracheids in Silver Fir trees during autumn. *Ann Bot* 95:959–965.
- Gričar J, Zupančič M, Čufar K, Koch G, Schmitt U, Oven P (2006) Effect of local heating and cooling on cambial activity and cell differentiation in the stem of Norway spruce (*Picea abies*). *Ann Bot* 97:943–951.
- Gričar J, Prislán P, Gryc V, Vavrčik H, De Luis M, Čufar K (2014) Plastic and locally adapted phenology in cambial seasonality and production of xylem and phloem cells in *Picea abies* from temperate environments. *Tree Physiol* 34:869–881.
- Guy RD (2014) The early bud gets to warm. *New Phytol* 202:7–9.
- Häkkinen R, Linkosalo T, Hari P (1998) Effects of dormancy and environmental factors on timing of bud burst in *Betula pendula*. *Tree Physiol* 18:707–712.
- Heide OM (1993) Daylength and thermal time responses of budburst during dormancy release in some northern deciduous trees. *Physiol Plant* 88:531–540.
- Huang JG, Bergeron Y, Zhai L, Denneler B (2011) Variation in intra-annual radial growth (xylem formation) of *Picea mariana* (Pinaceae) along a latitudinal gradient in western Quebec, Canada. *Am J Bot* 98:792–800.
- Huang JG, Deslauriers A, Rossi S (2014) Xylem formation can be modeled statistically as a function of primary growth and cambium activity. *New Phytol* 203:831–841.
- Jones G, Clegg B (2006) Screening exotic firs for the midwestern United States: interspecific variation in adaptive traits. *HortScience* 41:323–328.
- Kudo K, Nabeshima E, Begum S, Yamagishi Y, Nakaba S, Oribe Y, Yasue K, Funada R (2014) The effects of localized heating and disbudding on cambial reactivation and formation of earlywood vessels in seedlings of the deciduous ring-porous hardwood, *Quercus serrata*. *Ann Bot* 113:1021–1027.
- Ladefoged K (1952) The periodicity of wood formation. *Det Kongelige Danske Vidensk Selsk Skrift Dan Biol* 7:1–98.
- Man R, Lu P (2010) Effects of thermal model and base temperature on estimates of thermal time to bud break in white spruce seedlings. *Can J For Res* 40:1815–1820.
- Menzel A, Sparks TH, Estrella N et al. (2006) European phenological response to climate change matches the warming pattern. *Glob Change Biol* 12:1969–1976.
- Messaoud Y, Asselin H, Bergeron Y, Grondin P (2014) Competitive advantage of black spruce over balsam fir in coniferous boreal forests of Eastern North America revealed by site index. *For Sci* 60:57–62.

- Moser L, Fonti P, Büntgen U, Esper J, Luterbacher J, Franzen J, Frank D (2010) Timing and duration of European larch growing season along altitudinal gradients in the Swiss Alps. *Tree Physiol* 30:225–233.
- O'Reilly C, Owens JN (1989) Shoot, needle, and cambial growth phenology and branch tracheid dimensions in provenances of lodgepole pine. *Can J For Res* 19:599–605.
- Oribe Y, Kubo T (1997) Effect of heat on cambial reactivation during winter dormancy in evergreen and deciduous conifers. *Tree Physiol* 17:81–87.
- Oribe Y, Funada R, Shibagaki M, Kubo T (2001) Cambial reactivation in locally heated stems of the evergreen conifer *Abies sachalinensis* (Schmidt) Masters. *Planta* 212:684–691.
- Oribe Y, Funada R, Kubo T (2003) Relationships between cambial activity, cell differentiation and the localization of starch in storage tissues around the cambium in locally heated stems of *Abies sachalinensis* (Schmidt) Masters. *Trees* 17:185–192.
- Partanen J, Leinonen I, Repo T (2001) Effect of accumulated duration of the light period on bud burst in Norway spruce (*Picea abies*) of varying ages. *Silva Fenn* 35:111–117.
- Prislan P, Gričar J, de Luis M, Smith KT, Čufar K (2013) Phenological variation in xylem and phloem formation in *Fagus sylvatica* from two contrasting sites. *Agric For Meteorol* 180:142–151.
- Rensing KH, Owens JN (1994) Bud and cambial zone phenology of lateral branches from Douglas-fir (*Pseudotsuga menziesii*) seedlings. *Can J For Res* 24:286–296.
- Rossi S (2015) Local adaptations and climate change: converging sensitivity of bud break in black spruce provenances. *Int J Biometeorol* 59:827–835.
- Rossi S, Bousquet J (2014) The bud break process and its variation among local populations of boreal black spruce. *Front Plant Sci* 5:574.
- Rossi S, Anfodillo T, Menardi R (2006a) Trephor: a new tool for sampling microcores from tree stems. *IAWA J* 27:89–97.
- Rossi S, Deslauriers A, Anfodillo T (2006b) Assessment of cambial activity and xylogenesis by microsampling tree species: an example at the Alpine timberline. *IAWA J* 27:383–394.
- Rossi S, Deslauriers A, Anfodillo T, Carraro V (2007) Evidence of threshold temperatures for xylogenesis in conifers at high altitudes. *Oecologia* 152:1–12.
- Rossi S, Deslauriers A, Gričar J et al. (2008) Critical temperatures for xylogenesis in conifers of cold climates. *Glob Ecol Biogeogr* 17:696–707.
- Rossi S, Rathgeber CBK, Deslauriers A (2009) Comparing needle and shoot phenology with xylem development on three conifer species in Italy. *Ann For Sci* 66:206.
- Rossi S, Anfodillo T, Čufar K et al. (2013) A meta-analysis of cambium phenology and growth: linear and non-linear patterns in conifers of the northern hemisphere. *Ann Bot* 112:1911–1920.
- Sarvas R (1972) Investigations on the annual cycle of development of forest trees. Active period. *Commun Inst For Fenn* 76:1–110.
- Schmitt U, Möller R, Eckstein D (2000) Seasonal wood formation dynamics of beech (*Fagus sylvatica* L.) and black locust (*Robinia pseudoacacia* L.) as determined by the pinning technique. *J Appl Bot* 74:10–16.
- Schmitt U, Jalkanen R, Eckstein D (2004) Cambium dynamics of *Pinus sylvestris* and *Betula* spp. in the northern boreal forest in Finland. *Silva Fenn* 38:167–178.
- Schwalm CR, Ek AR (2001) Climate change and site: relevant mechanisms and modeling techniques. *For Ecol Manag* 150:241–257.
- Sheridan SC (2002) The redevelopment of a weather-type classification scheme for North America. *Int J Climatol* 22:51–68.
- Sutinen S, Partanen J, Viherä-Aarnio A, Häkkinen R (2012) Development and growth of primordially shoots in Norway spruce buds before visible bud burst in relation to time and temperature in the field. *Tree Physiol* 32:987–997.
- Swidrak I, Gruber A, Oberhuber W (2014) Xylem and phloem phenology in co-occurring conifers exposed to drought. *Trees* 28:1161–1171.
- Tadaki Y, Kitamura H, Kanie K, Sano H, Shigematsu A, Ohtsu S (1994) Leaf opening and falling of Japanese larch at different altitudes. *Jpn J Ecol* 44:305–314.
- Takahashi K, Koike S (2014) Altitudinal differences in bud burst and onset and cessation of cambial activity of four subalpine tree species. *Landsc Ecol Eng* 10:349–354.
- Thibeault-Martel M, Krause C, Morin H, Rossi S (2008) Cambial activity and intra-annual xylem formation in roots and stems of *Abies balsamea* and *Picea mariana*. *Ann Bot* 102:667–674.
- Uggla C, Mellerowicz EJ, Sundberg B (1998) Indole-3-acetic acid controls cambial growth in Scots pine by positional signaling. *Plant Physiol* 117:113–121.
- Uggla C, Magel E, Moritz T, Sundberg B (2001) Function and dynamics of auxin and carbohydrates during earlywood/latewood transition in Scots pine. *Plant Physiol* 125:2029–2039.
- Wang JY (1960) A critique of the heat unit approach to plant response studies. *Ecology* 41:785–790.
- Wolkovich EM, Davies TJ, Schaefer H, Cleland EE, Cook BI, Travers SE, Willis CG, Davis CC (2013) Temperature-dependent shifts in phenology contribute to the success of exotic species with climate change. *Am J Bot* 100:1407–1421.