

1 This is the peer reviewed version of the following article

2 ***Panuccio M.R. Chaabani S., Roula R., and Muscolo A. 2018. Bio-priming mitigates***
3 ***detrimental effects of salinity on maize improving antioxidant defense and preserving***
4 ***photosynthetic efficiency. Volume 132, Pages 465-474***

5 which has been published in final doi <https://doi.org/10.1016/j.plaphy.2018.09.033>

6 The terms and conditions for the reuse of this version of the manuscript are specified in
7 the publishing policy. For all terms of use and more information see the publisher's
8 website

9

10

11 **Bio-priming mitigates detrimental effects of salinity on maize improving**
12 **antioxidant defense and preserving photosynthetic efficiency.**

13 **Bio-priming enhances salinity tolerance in maize seedlings.**

14

15 Panuccio M.R., ¹ Chaabani S²., Roula R²., and Muscolo A^{1*}

16 ¹ **Agriculture Department, Mediterranea University, Feo di Vito-89122 Reggio**
17 **Calabria Italy**

18 ² **High Agronomic Institute of chott Mariem University of Sousse**

19

20

21

22 *Corresponding author: Professor Adele Muscolo

23 Agriculture Department, Mediterranea University,

24 Feo di Vito-89122 Reggio Calabria Italy

25 Email: amuscolo@unirc.it

26 Tel: 003909661694364

27

28

29

30

31

32

33

34 **ABSTRACT**

35 Salinity is an abiotic stress which seriously affects crop production over the world,
36 particularly in arid and semi-arid regions, with harmful effects on germination, growth
37 and yield. Maize (*Zea mays* L.), cultivated in a wide spectrum of soil and climatic
38 conditions, is the third most important cereal crop after rice and wheat, moderately
39 sensitive to salt stress. A saline level more than 0.25 M NaCl damages maize plants,
40 causing severe wilting. In this study, the effects of hydro-priming (distilled water) and
41 bio-priming (*Rosmarinus officinalis* L. and *Artemisia* L. leaf extracts) on seed
42 germination and seedling growth of maize, under 100 mM NaCl salinity were
43 investigated. The factorial experiments were carried out in greenhouse under controlled
44 condition (25°C in 12/12 day/night) based on a completely randomized design with
45 three replicates. Results showed that both hydro- and bio-priming increased germination
46 percentage and germination indexes in maize seeds. *Rosmarinus* extract was the most
47 effective in inducing salt resistance in 30 days old seedlings, with beneficial effects in
48 the strengthening of the antioxidant system and in the maintenance of a higher
49 photosynthetic efficiency under salt stress condition.

50

51 **Keywords:** Antioxidants; *Artemisia*; Priming; *Rosmarinus*, Salinity, *Zea mays*.

52

53 **1. Introduction**

54 Salinity is one of the major abiotic stresses that negatively affects crop productivity.
55 More than 800 million hectares of land worldwide are affected by either salinity (397
56 million hectares) or sodicity (434 million hectares) with a decline, by more than 50
57 percent, in average yield of the major crop plants (FAO, 2011; Munns and Tester,
58 2008). Salt stress occurs in areas where soils are naturally high in salts and
59 precipitations are low and/or where irrigation, hydraulic lifting of salty underground
60 water, or invasion of sea water in coastal areas bring salt to the surface soil. NaCl is the
61 predominant salt causing salinization worldwide (Munns and Tester, 2008). The most
62 widely accepted effects of salinity are decrease in germination percentage, germination
63 rate and in growth and metabolism of seedlings by creating an external osmotic
64 potential that prevents water uptake, or by causing specific ion toxicity and ion
65 imbalance. Salinity affects also cellular metabolism including photosynthesis and
66 synthesis of compatible solutes called “osmolytes” like proline, sugars (Amirjani, 2011)
67 and proteins (Sen and Alikamanoglu, 2011). Maize (*Zea mays* L.) is the third most
68 important cereal crop after rice and wheat and it is grown under a wide spectrum of soil
69 and climatic conditions. It is an important C₄ plant from the Poaceae family, moderately
70 sensitive to salt stress (Farooq et al., 2015). A saline level containing more than 0.25 M
71 NaCl may damage maize plants and stunt growth causing severe wilting (Menezes-
72 Benavente et al., 2004). For this reason, the aim of this work was to find a method that
73 could alert and/or attenuate the negative impact of salts increasing the tolerance to
74 salinity in maize plants. Seed priming is an easy, low cost and low risk technique
75 recently used to overcome the salinity problem in agricultural lands (Ibrahim, 2016;
76 Chen and Arora, 2013) It is a pre-sowing treatment and the most important priming
77 treatments include halo-priming (soaking seeds in inorganic salt solutions), solid matrix
78 priming (treatment of seeds with solid matrices), osmo-priming (soaking seeds in

solutions of different organic osmotic) and bio-priming (using a priming mixture integrated with bioactive molecules or beneficial microorganisms) (Nouman et al., 2014). All the above listed priming are able to 1) stimulate metabolic processes involved in the early phases of germination, producing high germination rate and great germination percentage 2) induce uniformity and faster emergence of seedlings from primed seeds, bring vigorous growth in adverse conditions (Imran et al., 2013). Maher et al. (2013) and Tzortzakis (2009) indicated also that seed priming in fenugreek, endive and chicory increased final germination percentage, germination speed and radicle length over the non-primed treatments in saline conditions. Generally, farmers facing with saline problems, cannot reclaim soil, or use expensive plant hormones, antioxidants or nutrients for seed priming (Basra et al., 2011; Imran et al., 2013). So, there is the need to explore new plant growth enhancers, naturals and environmentally friendly which should be reliable and economically sustainable under prevailing salinity conditions. This study was planned to investigate the potential of *Rosmarinus officinalis* L. and *Artemisia* L. aqueous extracts as seed bio-priming agents. These species, mostly found in arid and semi-arid areas, are widely distributed in Mediterranean countries, and represent new sources of natural antioxidant and antimicrobial agents. Thus, we hypothesized that seed priming with these natural extracts may alleviate salt stress in germinating maize seeds and improve seedling establishment by modulating antioxidants, photosynthetic pigments, ionic homeostasis and photosynthesis.

2. Materials and methods

2.1 Extract preparation and chemical characterization

Rosmarinus (*Rosmarinus officinalis* L.) and *Artemisia* (*Artemisia erba alba* L.) fresh leaves were collected between February and March 2016 from the region of Chouachi-HadjebAyoune (Tunisia 35° 23' North 9° 32' Est), identified according to the flora of Tunisia (Pottier-Alapetite, 1981), were

105 dried and grounded . Aqueous extracts were prepared by soaking the dried leaves,
106 overnight in distilled water (1:10 w/v). The suspensions were filtered with Whatman's
107 paper. Extracts have been analyzed by HPLC analysis with a Knauer (Berlin, Germany)
108 apparatus interfaced to a DAD detector (model 2600). HPLC-DAD technique is
109 commonly used to detect antioxidants in many matrix (Giuffrè, 2013). For separation a
110 binary gradient was prepared: (A) bi-deionized water and (B) acetonitrile, both were
111 acidified at pH 3 by with formic acid. The applied gradient was: 0-20 min, 95% A and
112 5% B; 20-50 min, the eluent B increased from 5 to 40%; 50-60 min, eluent B increased
113 from 40% to 95%; 60-65 min, the eluent B decreased from 95% to 5%; 65-70 min 95%
114 A and 5% B in isocratic. The analysis was performed with a constant flow rate of 1
115 ml/min. A Knauer C18 Eurosphere II separation column (Berlin, Germany) was used
116 (250 mm length x 4.6 mm internal diameter x 5 µm particle size). All standard
117 components (purity \geq 97%) were purchased from Sigma Co. (St. Louis, MO). All
118 solvents and reagents (HPLC grade) were purchased from Panreac (Barcelona, Spain).
119 The identification of unknown components in the extracts were performed by
120 comparison the retention times of the detected compounds with those of appropriate
121 standards.

122 2.2 Seed priming

123 For each treatment, 30 health seeds were surface sterilized with 5% sodium
124 hypochlorite for 5 min and then rinsed with sterile bi-distilled water. Aqueous extracts
125 of Rosmarinus (RP) and Artemisia (AP) were used for seed priming (Durak et al., 2016)
126 . Maize seeds were soaked in extracts ratio 1:5 (w/v) for 24 hours in darkness at room
127 temperature. The Hydro-primed (HP), bio-primed (RP, AP) and the un-primed (CNP)
128 seeds were dried back to their original moisture contents at room temperature and were
129 used as control. The experiments were performed in triplicate.

130 *2.3 Assessment of seed germination and morphological, physiological and biochemical*
131 *responses of seedlings*

132 After priming, seed germination tests were carried out. 10 seeds for each treatment
133 (priming and control) were placed in plastic pots (26 cm diameter × 27 cm height) filled
134 with sand and equilibrated with water (control) or NaCl 100 mM. 70% of field
135 capacity was maintained with distilled water. Each treatment was replicated three times.
136 Experiments were carried out in climatic chamber at 25°C in a 12/12-h photoperiod for
137 30 d. Seeds were considered germinated when a visible coleoptile protrusion was
138 observed. The germinated seeds were counted daily and the germination percentage was
139 calculated at the 7 days. Germination index (GI), mean germination time (MGT), time
140 to reach 50% germination (T_{50}) and total germination percentage TG were calculated as
141 follow:

142
$$GI = \sum(G_t/T_t); \text{ MGT} = \sum(G_t \times T_t) / \sum G_t \text{ and } T_{50} = t_i + [(N/2 - n_i) (t_j - t_i) / (n_j - n_i)].$$

143 G_t is the number of germinated seeds on day t , T_t is the time corresponding to G_t in
144 days, N is the final number of germination. n_i and n_j are the cumulative number of
145 seeds germinated by counts adjacent at times when $n_i < N/2 < n_j$ (Zhang et al., 2007).

146 Root length and shoot height were measured manually with a ruler and dry weights
147 were determined after drying at 80 °C for 24 h.

148 *2.4 Photosynthetic pigments in seedlings*

149 Fresh leaves, 0.05g for each treatment, were mixed with 2.5 ml of pure ethanol and
150 incubated for 24 h at 4°C in the dark. After, the samples were centrifuged for 10
151 minutes at 7000 rpm. For chlorophyll and carotenoid analysis, the absorbance of
152 supernatants was recorded at 649 nm, 665 nm, and 470 nm and their concentrations
153 (mg/g fresh weight) were calculated using Lichtenthaler's equations. (Lichtenthaler,

154 1987). Anthocyanins were extracted incubating 20 mg of fresh leaf in 0.5 ml of
155 methanol: HCl (99:1). After 24 h incubation at 4°C, samples were centrifuged at 6000 g
156 for 10 min at 4°C. (Panuccio et al., 2016). The absorbance was read at 530 and 657 nm
157 and anthocyanin content (μg anthocyanin /g fresh weight) was calculated according to
158 the following equation: $[A_{530\text{nm}} - (0,025 \cdot A_{657\text{nm}}) \cdot \text{ml of extract}] / \text{g fresh weight}$

159 *2.5 Chlorophyll fluorescence imaging*

160 Photosynthetic efficiency of primed and un-primed seedlings in absence and in presence
161 of salinity was evaluated by using an Imaging PAM Fluorometer (Walz, Effeltrich,
162 Germany). The chlorophyll fluorescence parameters detected were: Maximum quantum
163 yield of PSII photochemistry (F_v/F_m); Effective quantum yield of PSII photochemistry
164 ($Y(II)$); Quantum yield of regulated energy dissipation at PSII ($Y(NPQ)$); Quantum yield
165 of non-regulated energy dissipation at PSII ($Y(NO)$); Non photochemical quenching
166 coefficient (NPQ) and Electron transport rate (ETR).

167 *2.6 Antioxidant enzyme assay*

168 Fresh leaves, 0.5 g, were extracted (1:3 w:v) using a chilled mortar (4°C) in a 0.1 M
169 phosphate buffer solution (pH 7.0), containing 100 mg polyvinylpolypyrrolidone
170 (PVPP) and 0.1 mM $\text{Na}_2\text{-EDTA}$. The homogenates were centrifuged at 14000 g at 4°C
171 for 15 min and the supernatants were used. All enzyme activities were measured at 25
172 °C by a UV–visible light spectrophotometer (UV-1800 CE, Shimadzu, Japan).

173 Catalase (CAT, EC 1.11.1.6) activity was assessed, evaluating the disappearance of
174 H_2O_2 at 240 nm. The extinction coefficient (ϵ) = $0.036 \text{ mM}^{-1} \text{ cm}^{-1}$ was used (Beaumont
175 et al.,1990).

176 Ascorbate peroxidase (APX, EC 1.11.1.11) activity was evaluated monitoring the
177 decrease in absorbance at 290 nm for the oxidation of ascorbate (Nakano and

178 Asada,1981). Enzyme activity was quantified using the molar extinction coefficient for
179 ascorbate ($2.8 \text{ mM}^{-1} \text{ cm}^{-1}$).

180 Superoxide dismutase (SOD, EC 1.15.1.1) activity was assayed by recording the
181 decrease in absorbance of formazan produced from NBT, at 560 nm (Gupta et al.,
182 1993). The reaction mixture (3mL) contained 0.1 ml of 200 mM methionine, 0.1 ml of
183 2.25 mM nitro-blue tetrazolium, 0.1 ml of 3 mM EDTA, 1.5 ml of 100 mM potassium
184 phosphate buffer, 1 ml of distilled water and 0.05 ml of enzyme extract. The reaction
185 was started by adding 0.1 mL of riboflavin ($60 \mu\text{M}$) and placing the tubes below a light
186 source of two 15 W florescent lamps for 15 min. After, tubes were covered with
187 aluminum paper and the light was switched off. Tubes without enzyme extract were
188 used as control and they developed maximum colour. A non-irradiated complete
189 reaction mixture was the blank. One unit of enzyme activity corresponded to the
190 quantity of enzyme reducing the absorbance (560) nm of samples of 50%, compared to
191 tubes lacking enzymes.

192 All enzymatic activities were expressed as enzyme units (U)/mg fresh weight. One unit
193 of enzyme was the amount of enzyme necessary to decompose 1 mmol of substrate/ min
194 at 25°C .

195

196 *2.7 Total phenols and reduced glutathione analysis*

197 Total phenol content was determined with the Folin–Ciocalteu reagent according to the
198 method of Julkenen-Titto (1985). The reduced glutathione (GSH) was assayed
199 according the method of Jollow et al. (1974). Fresh leaves were homogenized in 3%
200 CCl_3COOH at 4°C and centrifuged at 1000g for 10 min at 4°C . The absorbance was
201 measured at 412 nm and related to a calibration curve of GSH solutions.

202 *2.8 Measurement of root morphology*

Seedlings were harvested 30 days after treatments and root weight was measured. Roots were analyzed by using the Epson Expression/STD 1600 scanner and personal computer (Intel Pentium III/500 CPU, 128 MB RAM), Regent Instrument, Inc. Root morphology was detected with the Win-RHIZO image analysis system (Regent Instruments, Quebec, Canada) (Panuccio et al., 2014).

2.9 Ion Analysis

Cations (Na^+ , K^+ , Ca^{2+} , Mg^{2+}) and anion (Cl_2) were extracted from leaves and analysed by ion chromatography (DIONEX ICS-1100). For anions, 0.500 g of dried material was extracted using 50 ml of anion solution (Na_2CO_3 / NaHCO_3 3.5 mM) under stirring for 20 minutes, then the homogenate was filtered and the chromatographic analysis was carried out. For cations, 1 g of dry material was ashed at 550 ° C for 5-6 hours in a porcelain capsule. The ash was then mineralized for 30 minutes at 100 ° C using 1M HCl solution. Finally, the obtained solution was filtered and analyzed by ion chromatograph, eluent (Methansulfonic acid 20 mM).

2.10 Statistical analysis

The experiments were in randomized complete block design with three replicates. Statistical analyses were performed using one-way ANOVA and mean comparisons were made using Tukey's test ($p < 0.05$). All data were analysed using SYSTAT 13.0 software (SPSS Inc.) The germination percentage data were previously subjected to arcsine transformation but were reported in tables as untransformed values.

3. Results

3.1 Composition of *Artemisia* and *Rosmarinus* leaf extracts

HPLC analysis showed qualitative and quantitative differences in the composition of the leaf aqueous extracts of the two officinal plants (Table 1). *Rosmarinus* extract

228 contained rosmarinic acid (63%) rutin (21%) carnosinic acid (6%) and neochlorogenic
229 acid (2.6%) that were absent in the artemisia extract. In contrast, Artemisia extract
230 contained narirutin (8%), naringin (13%) and protocatechuic acid (2%) that were absent
231 in rosmarinus leaf extract. Chlorogenic (52%) and syringic (6%) acids were present in
232 greater quantity in Artemisia than in rosmarinus leaf extract.

233

234 *3.2 Seed germination*

235 Bio and hydro-priming, in absence of salinity (0 mM NaCl), increased significantly ($p >$
236 0.05) total germination percentage (TG %) and germination index (GI) in comparison to
237 untreated seeds (CNP) (Table 2), and decreased MGT and T_{50} . RP and HP were the
238 treatments with the greatest positive effects on maize TG % and GI (Table 2). Under
239 100 mM NaCl, RP and AP increased the germination (TG %) and GI of about 60-70%
240 compared to CNP, and the T_{50} and MGT values were lower. No significant differences
241 in T_{50} and MGT were observed between bio and hydro-priming treatments (Table 2).
242 Analysis of Variance showed that GI, T_{50} and TG% were mainly affected by priming
243 and salinity, while MGT only by type of priming treatment. The interaction between
244 salinity and priming didn't affect significantly all the germination indices.

245

246 **Table 1**
 247 Chemical composition of leaf aqueous extracts of *Rosmarinus officinalis L.* and
 248 *Artemisia L.* Values are the means of three replicates \pm Standard Deviation (SD)

Rosmarinus extract		
	mg/l	SD
Gallic Acid	3.864	0.018
Neoclorogenic Ac.	4.092	0.097
Clorogenic Acid	1.961	0.062
Siringic Acid	1.969	0.061
Rutin	32.623	0.060
Rosmarinic Acid	99.116	0.004
Apigenin	4.089	0.050
Carnosol	1.047	0.052
Carnosic Acid	9.183	0.052

249

Artemisia extract		
	mg/l	SD
Gallic Acid	1.655	0.005
Protocatechic Acid	3.093	0.028
Clorogenic Acid	32.395	0.134
Siringic Acid	6.506	0.108
Narirutin	5.043	0.060
Naringin	7.838	0.035
3,4-Di-O-caffeoylquinic ac.	1.907	0.013
Luteolin	0.518	0.003
Apigenin	2.175	0.010
Kaempferol	0.384	0.019

250

251

Table 2.
Effect of NaCl (100 mM) and seed priming on germination indexes of maize. Control
no Priming (CNP) Hydro-Priming (HP) Rosmarinus Priming (RP) Artemisia Priming
(AP). Values are the means of three experiments \pm SE. Different letters indicate
significant differences ($P \leq 0.05$) among different plant treatments at the same salt
concentration. ANOVA *** $P \leq 0.001$; ** $P \leq 0.01$; * $P \leq 0.05$ (ANOVA and mean
comparison with Tukey's test).

MGT					ANOVA summary	
NaCl	CNP	HP	RP	AP	Salt	0.02
0	4.07 \pm 0.19 a	2.96 \pm 0.26 b	3.07 \pm 0.06 b	3.14 \pm 0.06 b	Priming	16.22***
100	3.81 \pm 0.12 a	3.23 \pm 0.07 b	3.10 \pm 0.06 b	3.16 \pm 0.05 b	Salt x Prim.	1.09
GI						
NaCl	CNP	HP	RP	AP	Salt	5.97*
0	6.17 \pm 1.31 c	10.91 \pm 1.7 b	13.6 \pm 0.97 a	12.04 \pm 0.3ab	Priming	29.99***
100	5.03 \pm 1.16 c	9.38 \pm 1.41 b	12.18 \pm 2.3 a	10.52 \pm 1.1ab	Salt x Prim.	0.026
T ₅₀						
NaCl	CNP	HP	RP	AP	Salt	10.86**
0	4.96 \pm 0.80 a	3.11 \pm 0.76 b	2.21 \pm 0.34 c	3.09 \pm 0.16 b	Priming	27.43***
100	5.25 \pm 0.43 a	3.88 \pm 0.33 b	3.21 \pm 0.32 b	3.60 \pm 0.17 b	Salt x Prim.	0.66
TG %						
NaCl	CNP	HP	RP	AP	Salt	5.26*
0	60.1 \pm 3c	76.7 \pm 5b	96.7 \pm 5a	90.3 \pm 5a	Priming	22.60***
100	46.7 \pm 5d	73.3 \pm 1c	90.1 \pm 2a	82.3 \pm 2b	Salt x Prim.	0.35

Table 3. Shoot and root length of 30 days old maize seedlings derived from no-primed (CNP), hydro-primed (HP), Rosmarinus (RP) and Artemisia (AP) bio-primed seeds and grown in the absence (0) or in the presence of 100 mM NaCl. Values are the means of three experiments \pm SE. Different letters indicate significant differences ($P \leq 0.05$) among different plant treatments at the same salt concentration. * indicates significant differences ($P \leq 0.05$) between different salt concentrations of the same priming treatment.

Shoot Length		
	0 mM NaCl	100 mM NaCl
CNP	39.7 \pm 2.5 b*	30.0 \pm 2.6 c
HP	48.3 \pm 3.1 a	47.5 \pm 1.3 a
RP	48.0 \pm 2.3 a	48.0 \pm 2.3 a
AP	50.3 \pm 1.5 a*	41.0 \pm 1.0 b
Root Length		
	0 mM NaCl	100 mM NaCl
CNP	31.3 \pm 2.5c*	25.1 \pm 1.5c
HP	47.5 \pm 2.3b*	30.2 \pm 1.0b
RP	57.3 \pm 1.5 a*	45.3 \pm 1.3a
AP	57.0 \pm 2.0a*	47.5 \pm 2.0a

272

273 3.3 Seedling growth

274 Hydro- and bio-priming increased shoot and root length both in presence and in
 275 absence of salinity (Table 3). Bio-priming stimulated better their growth in
 276 comparison to HP and control. No significant differences between the two bio-
 277 priming treatments (RP and AP) were observed both in absence and in presence of
 278 salinity. In absence of salinity, total root length was increased mainly by HP,
 279 instead in presence of 100 mM NaCl, the greatest elongation was observed in
 280 presence of RP and AP (Table 4). Root volume increased in primed seedlings, in
 281 presence and in absence of salinity, and the greatest enhancement was observed
 282 with RP treatments.

283

284 **Table 4**

285 Root parameters of 30 days old maize seedlings derived from no-primed (CNP), hydro-primed
 286 (HP), Rosmarinus (RP) and Artemisia (AP) bio-primed seeds and grown in the absence (0) or in
 287 the presence of 100 mM NaCl. Values are the means of three replicates \pm SE. Different letters
 288 indicate significant differences ($P \leq 0.05$) among different plant treatments at the same salt
 289 concentration. ANOVA *** $P \leq 0.001$; ** $P \leq 0.01$; * $P \leq 0.05$ (ANOVA and mean comparison
 290 with Tukey's test).

Total root length (cm)					ANOVA summary	
NaCl	CNP	HP	RP	AP	Salt	
0	5454.7 $\pm 25.6b$	8113.6 $\pm 11.0a$	4947.0 $\pm 22.0c$	5728.7 $\pm 15.1b$	Priming	5753.5*** 3059.1***
100	2598.6 $\pm 11.1d$	3750.9 $\pm 13.5c$	5077.1 $\pm 43.02a$	4950.7 $\pm 15.2b$	Salt x Prim.	2651.1***
Root volume (cm ³)						
NaCl	CNP	HP	RP	AP	Salt	
0	1.83 $\pm 0.01d$	3.15 ± 0.004 c	4.04 $\pm 0.01a$	3.25 $\pm 0.01b$	Priming	1561.3**** 1341.8****
100	2.05 $\pm 0.02c$	2.39 $\pm 0.03b$	2.99 $\pm 0.01a$	2.07 $\pm 0.05c$	Salt x Prim.	328.9****
Surface root area (cm ²)						
NaCl	CNP	HP	RP	AP	Salt	
0	519.6 $\pm 7.7d$	1792.9 $\pm 7.0a$	591.0 $\pm 7.0c$	1658.9 $\pm 13.4b$	Priming	193.2**** 488.9****
100	1588.9 $\pm 45.5a$	1060.7 $\pm 19.0d$	1382.1 $\pm 58.5b$	1133.9 $\pm 23.1c$	Salt x Prim.	1870.2** *
Root DW (g plant ⁻¹)						
NaCl	CNP	HP	RP	AP	Salt	
0	0.08 $\pm 0.01d$	0.11 ± 0.001 c	0.15 $\pm 0.003a$	0.14 $\pm 0.001b$	Priming	721.4**** 238.4****
100	0.04 $\pm 0.01c$	0.10 ± 0.002 a	0.10 $\pm 0.004a$	0.08 $\pm 0.003b$	Salt x Prim.	62.0****

291

292

293

294 Root surface area increased in absence of salinity with HP and AP priming in respect to
295 untreated, conversely, under salinity, CNP roots showed the highest surface area. Root
296 dry weight (DW) was significantly the highest when seedlings were pretreated with
297 Rosmarinus extract (Table 4). ANOVA analysis underlined that Total root length, was
298 the parameters most affected by salinity, priming and their interaction (F-ratios) (Table
299 4). Salinity more than priming treatments influenced DW and root volume; conversely,
300 the interaction of the two variables mainly affected root surface area (Table 4). Under
301 salinity, the specific length (SRL) and the fineness (RF) of roots in the RP and AP
302 primed seedlings increased significantly (Fig. 1). These results indicated longer roots
303 per unit of root mass respect to control and HP seedlings in which a reduction of SRL
304 and RF was instead evident (Fig.1). The interaction between treatment and salinity had
305 a significant effect on both SRL and RF values (F-ratios). The ratio between root mass
306 and volume (RTD) decreased at 100mM NaCl, except for the root of the HP primed
307 seedlings. The salinity mainly affected RTD (F-ratio) and this can be explained by the
308 remarkable reduction of the radical dry weights caused by salt (Fig. 1).

309

310 **Table 5**

311 Phenols and reduced glutathione (GSH) in leaves of 30 days old maize seedlings
 312 derived from no- primed (CNP), hydro-primed (HP), Rosmarinus (RP) and Artemisia
 313 (AP) bio-primed seeds and grown at 0 and 100 mM NaCl. Values are the means of three
 314 replicates experiments \pm SE. Different letters in the same row denote significant
 315 differences among treatments ($P \leq 0.05$). ANOVA *** $P \leq 0.001$; ** $P \leq 0.01$; * $P \leq 0.05$
 316 (ANOVA and mean comparison with Tukey's test).

317 .

Phenols ($\mu\text{g GAE/g D.W.}$)					ANOVA summary	
NaCl	CNP	HP	RP	AP	Salt	
0	24.78 \pm 2.61a	22.23 \pm 1.0a	23.53 \pm 1.16a	22.12 \pm 0.77a	Priming	50.57*** 5.63**
100	8.32 \pm 0.64b	19.97 \pm 1.34a	20.11 \pm 0.04a	21.63 \pm 0.07a	Salt x Prim.	22.88***
GSH ($\mu\text{moles GSH/g F.W.}$)						
NaCl	CNP	HP	RP	AP	Salt	
0	9.54 \pm 0.03b	13.29 \pm 0.1a	13.33 \pm 0.04a	9.18 \pm 0.04b	Priming	20419*** 11484***
100	11.31 \pm 0.07d	17.44 \pm 0.03b	24.15 \pm 0.03a	13.14 \pm 0.03c	Salt x Prim.	2930***


318

319

320

321 **Figure1**

322 Specific root length (SRL = root length/root DW), root tissue density ((RTD = root DW/root
323 volume), root fineness (RF= root length/root volume) of 30 days old maize seedlings, derived
324 from no-primed (CNP), hydro-primed (HP), Rosmarinus (RP) and Artemisia (AP) bio-primed
325 seeds and grown at 0 and 100 mM NaCl. Values are the means of three replicates experiments \pm
326 SE. Bars with different letters are statistically different at ($P \leq 0.05$). ANOVA *** $P \leq 0.001$;
327 ** $P \leq 0.01$; * $P \leq 0.05$ (ANOVA and mean comparison with Tukey's test).




328

329

Figure 2

Antioxidant enzymatic activities in leaves of 30 days old maize seedlings: Catalase (CAT), Superoxide dismutase (SOD), Ascorbate peroxidase (APX). Seedlings were derived from no-primed (CNP), hydro-primed (HP), Rosmarinus (RP) and Artemisia (AP) bio-primed seeds and grown at 0 and 100 mM NaCl. Values are the means of three replicates experiments \pm SE. Bars with different letters are statistically different at ($P \leq 0.05$). ANOVA *** $P \leq 0.001$; ** $P \leq 0.01$; * $P \leq 0.05$ (ANOVA and mean comparison with Tukey's test).



339 *3.4 Enzyme activities, phenols and antioxidants*

340 Antioxidant enzyme activities of seedlings resulted differently influenced by salt and by
341 priming treatment (Fig.2). In untreated leaves (CNP), CAT activity significantly
342 increased (+ 25%) under salinity, while SOD and APX activities lowered (Fig. 2). The
343 same trend was shown for HP treated seedlings. In AP primed seedlings, SOD and CAT
344 activities were not significantly influenced by 100mM NaCl, and the highest APX
345 activity was detected (Fig. 2). The salinity in RP primed seedlings caused remarkable
346 increases in SOD and APX activities, while CAT activity was reduced. F-ratios values
347 evidenced that salt and priming, individually and in combination, significantly affected
348 the antioxidant enzymatic system of maize leaves (Fig. 2). CAT activity was mainly
349 influenced by the interaction of salt and treatment, whereas the salinity induced the
350 major changes in SOD and APX activities. In absence of salinity, in all seedlings,
351 regardless of the treatments, the phenol content was similar (Table 5). The salinity
352 caused significant reductions in phenols only in leaves of untreated seedlings. The
353 amount of reduced glutathione (GSH) increased at 100mM NaCl in all samples and the
354 highest concentrations were detected in leaves of HP and RP primed seedlings. The
355 salinity induced the most significant effects both on phenols and GSH contents (F-
356 ratios).


357 *3.5 Ion accumulation*

358 Under salinity, total ion content increased significantly (+ 40%) in leaves of un-primed
359 seedlings, with a remarkable decrease in cation and a simultaneous increase in anion
360 percentage (Fig. 3). Under salinity HP and AP primed seedlings had the highest content
361 of total ions and the major anion percentages compared to all other treatments (Fig. 3).

362 In RP primed seedlings, only a slight increase in the total ion content (+8%) was
 363 detected, without causing any significant change in the percentages of anions and
 364 cations (Fig. 3).

365 **Figure 3**


366 Total ion content and anion and cation percentages in leaves of 30 days old maize
 367 seedlings derived from no- primed (CNP), hydro-primed (HP), Rosmarinus (RP) and
 368 Artemisia (AP) bio-primed seeds and grown at 0 and 100 mM NaCl. Values are the
 369 means of three replicates experiments \pm SE. Bars with different letters are statistically



370 different at ($P \leq 0.05$).

371 **Figure 4.**

372 Na^+ K^+ and Cl^- content in roots and shoots of in leaves of 30 days old maize seedlings
 373 derived from no-primed (CNP), hydro-primed (HP), Rosmarinus (RP) and Artemisia
 374 (AP) bio-primed seeds and grown at 0 and 100 mM NaCl. Values are the means of
 375 three replicates experiments \pm SE. Bars with different letters are statistically different at
 376 ($P \leq 0.05$).



379

380 **Table 6**

381 Cation content against sodium and chloride, in leaves of 30 days old maize seedlings
 382 derived from no-primed (CNP), hydro-primed (HP),
 383 Rosmarinus (RP) and Artemisia (AP) bio-primed seeds and grown at 0 and 100 mM
 384 NaCl. Values are the means of three experiments \pm SE.

385 Different letters in the same column denote significant differences among treatments (P
 386 ≤ 0.05).

387

388

		$\text{Ca}^{2+}/\text{Na}^+$		Mg^{2+}/N	$\text{Mg}^{2+}/\text{Cl}^-$			
		K^+/Na^+	$^+$	a^+	K^+/Cl^-	$\text{Ca}^{2+}/\text{Cl}^-$	$^-$	Na^+/Cl^-
CN	0	8.75 \pm 0.7	1.37 \pm 0.0	1.32 \pm 0.0	2.64 \pm 0.0	0.41 \pm 0.0	0.40 \pm 0.02	0.32 \pm 0.0
	P	a	2 a	2 a	4 a	2 b	a	1 d
	10	0.49 \pm 0.0	0.14 \pm 0.0	0.09 \pm 0.0	0.47 \pm 0.0	0.14 \pm 0.0	0.10 \pm 0.1	0.97 \pm 0.0
	0	5 e	1 e	1 e	2 e	1 d	1 d	3 a
HP	0	2.09 \pm 0.1	0.73 \pm 0.0	0.74 \pm 0.0	1.19 \pm 0.0	0.39 \pm 0.0	0.42 \pm 0.0	0.57 \pm 0.0
	P	0 b	5 b	3 b	5 b	2 b	2 a	2 c
	10	1.06 \pm 0.1	0.37 \pm 0.0	0.44 \pm 0.0	0.61 \pm 0.0	0.22 \pm 0.0	0.25 \pm 0.0	0.56 \pm 0.0
	0	1 d	2 d	4 d	3 d	2 c	2 b	2 c
RP	0	1.64 \pm 0.0	0.66 \pm 0.0	0.55 \pm 0.0	1.05 \pm 0.0	0.42 \pm 0.0	0.37 \pm 0.0	0.64 \pm 0.0
	P	5 c	4 b	4 c	3 c	3 b	3 a	3 b
	10	1.51 \pm 0.0	0.57 \pm 0.0	0.39 \pm 0.0	1.03 \pm 0.0	0.39 \pm 0.0	0.27 \pm 0.0	0.69 \pm 0.0
	0	5 c	3 b	3 d	1 c	3 b	3 b	4 b
AP	0	1.02 \pm 0.0	0.49 \pm 0.0	0.44 \pm 0.0	1.03 \pm 0.0	0.49 \pm 0.0	0.44 \pm 0.0	1.01 \pm 0.0
	P	6 d	2 c	3 d	4 c	2 a	2 a	3 a
	10	0.14 \pm 0.0	0.42 \pm 0.0	0.39 \pm 0.0	0.21 \pm 0.0	0.14 \pm 0.0	0.14 \pm 0.0	0.34 \pm 0.0
	0	2 e	1 d	2 d	2 f	2 d	2 c	1 d

389

390

391 **Table 7.**

392 Photosynthetic pigments in leaves of 30 days old maize seedlings derived from no-
 393 primed (CNP), hydro-primed (HP), Rosmarinus (RP) and Artemisia (AP) bio-primed
 394 seeds and grown at 0 and 100 mM NaCl. Values are the means of three replicates
 395 experiments \pm SE. Different letters in the same row denote significant differences
 396 among treatments ($P \leq 0.05$).

	CNP	HP	RP	AP
Total Chlorophyll (mg/g F.W.)				
0	10.93 \pm 0.11d	13.30 \pm 0.08c	17.92 \pm 0.13a	16.65 \pm 0.18b
100	19.81 \pm 0.02c	20.75 \pm 0.04c	27.64 \pm 0.07a	22.61 \pm 0.10b
Chlorophyll a (mg/g F.W.)				
0	7.92 \pm 0.08d	9.56 \pm 0.06c	11.76 \pm 0.09a	13.40 \pm 0.09a
100	11.42 \pm 0.01c	11.67 \pm 0.02c	14.89 \pm 0.03b	16.04 \pm 0.07a
Chlorophyll b (mg/g F.W.)				
0	3.03 \pm 0.03d	3.74 \pm 0.02b	6.16 \pm 0.04a	3.25 \pm 0.04c
100	8.39 \pm 0.01c	9.09 \pm 0.02b	12.75 \pm 0.04a	6.57 \pm 0.03d
Carotenoids (mg/g F.W.)				
0	2.12 \pm 0.02c	2.49 \pm 0.02b	3.30 \pm 0.02a	1.10 \pm 0.01d
100	0.63 \pm 0.03c	0.69 \pm 0.02c	1.00 \pm 0.01b	4.63 \pm 0.02a
Anthocyanins (μ g/g F.W.)				
0	7.08 \pm 0.07c	7.45 \pm 0.05b	6.44 \pm 0.04d	15.08 \pm 0.16a
100	12.52 \pm 0.13a	10.27 \pm 0.02c	11.56 \pm 0.03b	12.40 \pm 0.05a

397

398

399 **Table 8**

400 Effects of salinity and priming treatments on chlorophyll fluorescence parameters in
 401 leaves of 30 days old maize seedlings derived from no priming (CNP), hydro-priming
 402 (HP), Rosmarinus (RP) and Artemisia (AP) bio-priming seeds and grown at 0 and 100
 403 mM NaCl. Values are the means of three replicates experiments \pm SE. Different letters
 404 in the same row denote significant differences among treatments ($P \leq 0.05$).

	CNP	HP	RP	AP
Fv/Fm				
0	0.54 \pm 0.02c	0.66 \pm 0.02a	0.62 \pm 0.02b	0.59 \pm 0.01b
100	0.43 \pm 0.02c	0.51 \pm 0.01b	0.54 \pm 0.01b	0.58 \pm 0.02a
Y(II)				
0	0.35 \pm 0.01c	0.40 \pm 0.01b	0.43 \pm 0.01a	0.45 \pm 0.01a
100	0.19 \pm 0.02c	0.21 \pm 0.02c	0.30 \pm 0.01b	0.35 \pm 0.01a
Y (NPQ)				
0	0.29 \pm 0.02a	0.27 \pm 0.01a	0.20 \pm 0.02b	0.23 \pm 0.01b
100	0.33 \pm 0.03b	0.41 \pm 0.03a	0.23 \pm 0.01c	0.25 \pm 0.01c
Y(NO)				
0	0.23 \pm 0.01a	0.21 \pm 0.01a	0.14 \pm 0.01b	0.14 \pm 0.03b
100	0.30 \pm 0.03a	0.20 \pm 0.02b	0.13 \pm 0.02c	0.14 \pm 0.02c
NPQ				
0	0.43 \pm 0.02a	0.33 \pm 0.03b	0.35 \pm 0.02b	0.40 \pm 0.01a
100	0.58 \pm 0.02a	0.50 \pm 0.03b	0.39 \pm 0.01c	0.39 \pm 0.03c
ETR				
0	35.02 \pm 0.02b	46.50 \pm 1.02a	45.25 \pm 0.21a	45.04 \pm 0.31a
100	19.02 \pm 0.23d	30.56 \pm 0.14c	33.76 \pm 0.30b	40.99 \pm 1.21a

405 The salt condition significantly increased the Na^+ and Cl^- concentrations in all seedlings
406 (Fig. 4). Na^+ was generally higher in roots of primed seedling. Conversely, CNP had the
407 highest sodium and chloride content, mainly accumulated in shoots (Fig. 4). In CNP and
408 AP primed seedlings a significant decrease in K^+ content was observed. Under salinity,
409 HP and RP primed seedlings showed a similar trend regarding content, uptake and
410 distribution of Na^+ and Cl^- ions. In RP seedlings, at 100mMNaCl, the highest K^+
411 content was detected, mainly accumulated in shoots. (Fig. 4). In shoot of CNP plants
412 grown in absence of salinity, the K^+/Na^+ , $\text{Ca}^{2+}/\text{Na}^+$, $\text{Mg}^{2+}/\text{Na}^+$ ratios were the highest
413 (Table 6). Conversely, under salinity the above mentioned ratios were significantly
414 lower in comparison with all primed plants. In AP seedlings, due to the highest Cl^- ion
415 accumulation, at 100mM NaCl, the lowest ratios of cations against chloride were
416 detected .

417 *3.6 Chlorophyll, carotenoid and anthocyanin content*

418 Total chlorophyll content increased in all seedlings under salinity (Table 7). At 100 mM
419 NaCl, the highest amount of total chlorophyll was detected in RP leaves. Salinity caused
420 a decrease in carotenoids and a simultaneous increase in anthocyanin content in all
421 samples, except for AP seedlings which showed an opposite behaviour (Table 7)

422 Bio-primed seedlings had the higher Y(II), ETR, and Fv/Fm values in respect to un-
423 primed seedlings both in presence and in absence of salinity (Tab.8). In CNP and HP
424 primed seedlings, a decline in effective quantum yield of photochemical energy
425 conversion in PSII, Y(II), was accompanied by a significant increase in the quantum
426 yield of regulated energy dissipation in PSII (YNPQ). The quantum yield of non-
427 regulated energy dissipation at PSII, Y(NO), significantly increased under salinity only
428 in CNP leaves, differently a weak decrease in primed seedlings was observed. Salinity
429 caused an increase in non-photochemical quenching (NPQ) of all samples to prevent
430 photoinhibition in leaves.

431

432 **4. Discussion**

433 Salinity represents nowadays a serious problem for the Mediterranean countries, and in
434 particular for North African regions where changing climatic conditions are further
435 exacerbating water scarcity and soil salinity. It is known that soil salinity, with high
436 percentages of chloride and sodium, affects plant growth by modifying their
437 morphological, anatomical biochemical and physiological traits (Muscolo et al., 2015;
438 Panuccio et al., 2003) in particular way at seed and seedling stage (Ibrahim, 2016). Our
439 results showed that the detrimental effects of salinity on germination were less severe
440 when maize seeds were pretreated with RP and AP extracts rather than water or
441 untreated. This ameliorative effect on germination under salinity could be related to the
442 phytochemical content of these two extracts. Rosmarinic extract contained great amount
443 of rosmarinic acid that is known to have numerous beneficial and protective effects at
444 cellular levels with antioxidant potential, and it is considered as an effective scavenger
445 compound against Na^+ toxicity, combating cellular damage under stress (Adomako-
446 Bonsu et al., 2017). Conversely, the effectiveness of AP extract could be prevalently
447 related to its content of chlorogenic and syringic acids, referred as potent cellular
448 antioxidants with scavenging properties (Xu et al., 2017). Generally, salinity causes
449 oxidative stress in plants, by disruption of the balance between ROS production and
450 elimination, leading to an increase in antioxidant compounds such as phenols, reduced
451 glutathione, anthocyanins, carotenoids, and with a shift from primary to secondary
452 metabolism (Munns & Tester, 2008). In maize, the negative effects of salt stress depend
453 on the stress degree and plant growth stages (Imran et al., 2013). Our results showed,
454 under salinity, a significant increase in reduced glutathione (GSH) content of all
455 seedlings, confirming its essential role in keeping ROS under control. GSH can be
456 involved as an antioxidant in direct reactions with free radicals, or in cooperation with
457 ascorbate in the ascorbate-glutathione cycle which plays a central role in integration of

458 redox signals. According to the most significant increase in GSH content, APX activity
459 increased in AP and RP primed seedlings under salinity. Among antioxidant enzymes,
460 CAT activity is generally low under normal growth conditions and it increases only at
461 relatively high H₂O₂ concentrations or under stress conditions, to support APX, SOD,
462 and other enzymes primarily involved in ROS homeostasis (Papalia et al., 2017). At
463 100mM NaCl, the lowest activities of CAT in RP and AP primed seedlings suggested
464 that priming treatment was able to contrast the salt sensitivity of maize seedlings. In
465 numerous cases, the beneficial impact of priming on plant growth is more evident under
466 non-optimal than under optimal conditions because of an increase in stress resistance
467 (Ibraim, 2016). As reported by some authors, several components of the ROS-mediated
468 signaling pathways are accumulated and activated during the first hydration phase and
469 the final degree of stress resistance of seedlings can be linked to the persistence, even
470 after germination, of all the antioxidant mechanisms activated in seed (Paparella et al.,
471 2015). Then the priming treatment with AP and RP extracts, containing molecules with
472 recognized antioxidant properties, may have contributed to amplify and strengthen the
473 antioxidant response leading to an increase in salt resistance of maize seedlings.

474 Maize seedling grown in presence of salinity, showed an accumulation of sodium and
475 chloride. The ability of primed seedlings to limit Na⁺ transport into shoot, compared to
476 untreated plants, was important for the maintenance of growth rates and protection of
477 the metabolic process from the toxic effect of Na⁺. This results perfectly agree with
478 findings reported by Jamil et al. (2012) in rice, by Akram et al. (2011) in sunflower, by
479 Perveen et al. (2012) in wheat, demonstrating that the plant growth reduction under
480 saline condition was dependent mainly by the accumulation of sodium at leaf level. The
481 increase of sodium content caused a decrease in K⁺/Na⁺ ratios of all seedlings. In maize
482 plants, salt toxicity is mainly due to antagonistic effect of Na⁺ on K⁺ uptake and to a
483 strong interference of sodium and chloride ions with other essential mineral elements,

484 leading to severe nutritional imbalances (Farooq et al., 2015). In HP and RP primed
485 seedling exposed to salt stress, Na^+ and Cl^- concentrations were significantly lower
486 compared to all other samples, and potassium content did not changed. These results
487 suggest that HP and RP primed plants were able to avoid an excessive sodium entry and
488 maintain an efficient K^+ uptake as adaptive strategy under salt stress. Increased sodium
489 accumulation also disturbs calcium nutrition (Shahzad et al., 2012). The Ca^{2+} content, at
490 100mM NaCl, was significantly higher in primed seedlings, and the maintenance of
491 adequate levels of Ca^{2+} , under stress condition, is important by considering the
492 fundamental role of this ion in stabilizing the cell wall and membrane and also as a
493 signal in induction of the antioxidant enzymes. As reported by many authors, the
494 decrease in growth observed in many plants subjected to salinity stress is often
495 associated with a decline in their photosynthetic capacity (Akram et al., 2011). This
496 decrease aggravates the amount of excess excitation energy which should be
497 appropriately dissipated to avoid that uncontrolled production of ROS and
498 photosynthetic apparatus damage occur. The synthesis of anthocyanins is induced in
499 many plants for their protective role against light and other stress conditions. Eryilmaz
500 (2006) demonstrated an increase in anthocyanins in response to salt stress in tomato and
501 cabbage seedlings, evidencing a positive correlation between anthocyanins and NaCl.
502 Our results, confirmed the involvement of anthocyanins in maize response to salinity
503 and the highest content was in leaves of CNP seedlings, indicating an higher stress
504 condition respect to that of primed plants. Salt stress decreases photosynthetic
505 performance in plants, leading to a decline of CO_2 fixation and an enhancement of the
506 oxigenase activity of RUBPco (Kangasjärvi et al., 2012). Our results evidenced that
507 priming treatments, with RP and AP extracts, were able to preserve photosynthetic
508 apparatus and photosynthetic efficiency of maize seedlings against salt stress. Salinity
509 lowered the potential photochemical activity of PSII, expressed by Fv/Fm ratios in

maize, as reported by Han et al. (2010), Akram et al. (2011). The lowest Fv/Fm values were detected in CNP and HP primed seedlings, indicating that the conversion efficiency of primary light energy and the potential activity of PSII were mainly affected. In addition, in CNP plants, the significant increase in Y(NO), the index of non regulated energy dissipation at PSII, confirmed a damage to photosynthetic apparatus due to the stress condition. Photodamage steps are mediated by reactive oxygen species and the main target are the photosynthetic reaction centers, primarily of photosystem II (PSII), and likely also PSI (Ruban et al., 2012). The general NPQ increase in all samples may reflect heat dissipation of light energy in the antenna system. NPQ, often referred to as “feedback de-excitation”, is considered the most important short-term reversible photoprotective process in higher plants (Lambrev et al., 2012).

The priming treatment influenced also the growth and morphology of root apparatus. The root system possesses a certain phenotypic plasticity and under stress condition this variability represents a major survival strategy allowing plants to concentrate their resources where nutrients and water are more easily available (Panuccio et al., 2014). Salinity reduced total root length in CNP and HP primed seedlings indicating a decrease in carbon skeleton supply from the shoot. This reduced root growth and elongation was also confirmed by the lowest values of SRL and RF, structural root traits associated with the nutrient acquisition capacities of plants, that respond rapidly to stresses and environmental changes. A different behavior was observed in root system of AP and RP primed seedlings, where the greatest SRL and RF ratios suggested that the plants maximized the effectiveness of roots in water and nutrient uptake (Fitter and Stickland, 1991). Both the extracts are effective in inducing salt resistance in maize, however RP treated seedlings showed a better response to salinity and in AP treated seedlings some stress signals were detected. The root and shoot grew less, the content of anthocyanins and carotenoids were higher while total chlorophyll was lower than rosmarinus treated

536 ones. In AP primed seedlings, under salinity, was also observed a significant increase
537 in total ion content and a different ion compartmentalization of cations and anions
538 between root and shoot, leading to ion imbalance with consequent possible toxicity.

539

540 **5. Conclusion**

541 In short, although bio-primed maize seedlings didn't maintain equal growth under salt
542 stress in comparison with the same plants grown under normal conditions, they showed
543 a better growth performance under salinity as compared to untreated and hydro-primed
544 seedlings. Beneficial effects of bio-priming treatment are mainly expressed by the
545 improvement of nutritional and nutraceutical qualities due to the increasing of
546 antioxidant compounds with relevant beneficial effect on human health. The use of
547 natural extracts as bio-priming agents could be attractive because it reduces the risk of
548 negative environmental impact in respect to the use of chemical agents, representing an
549 up and coming ecofriendly technique to overcome agricultural problems in degraded
550 land. The results obtained evidence a new potential application of these two species
551 already widely used in the pharmaceutical and cosmetic field. Bio-priming represents a
552 non-expensive and a value added practice that greatly can increase yield of salt sensitive
553 crops in saline lands, improving the economic returns to farmers in Tunisia

554

555

556

557 **Conflicts of interest**

558

559 The authors have declared that no competing interests exist.

560

561 **Author contributions**

562 All authors discussed the results and commented on the manuscript.

563 Muscolo Adele designed the project.

564 Panuccio Maria Rosaria analyzed the data.

565 Chaabani Saber, Roula Rabia worked in the laboratory

566 Muscolo Adele and Panuccio Maria Rosaria wrote the manuscript.

567

568 **References**

569
570 Adomako-Bonsu, A.G., Chan S.L., Pratten, M., Fry, J.R., 2017. Antioxidant activity of
571 rosmarinic acid and its principal metabolites in chemical and cellular systems:
572 Importance of physico-chemical characteristics. *Toxicol In Vitro*, 40, 248-255.
573 <https://doi.org/10.1016/j.tiv.2017.01.016>.

574 Akram , M., Ashraf, M.Y., Jamil, M., Iqbal, R.M., Nafees, M., Khan, M.A., 2011.
575 Nitrogen Application Improves Gas Exchange Characteristics and Chlorophyll
576 Fluorescence in Maize Hybrids under Salinity Conditions. *Russ J Plant Physiol*.
577 58, 394–401. <https://doi.org/10.1134/S1021443711030022>

578 Amirjani, M.R., 2011. Effect of salinity stress on growth, sugar content, pigments and
579 enzyme activity of rice. *Int. J. Botany*. 7, 73-81. [https://doi.org/](https://doi.org/10.3923/ijb.2011.73.81)
580 [10.3923/ijb.2011.73.81](https://doi.org/10.3923/ijb.2011.73.81)

581 Basra, S.M.A., Iftikhar, M.N., Afzal, I., 2011. Potential of moringa (*Moringa oleifera*)
582 leaf extract as priming agent for hybrid maize seeds. *Int J Agric Biol*. 13,
583 1006–1010.

584 Beaumont, F., Jouve, H.M., Gagnon, J., Gaillard, J., Pelmont, J., 1990. Purification
585 and properties of a catalase from potato tubers (*Solanum tuberosum*). *Plant Sci*.
586 72, 19–26.

587 Chen, K., Arora, R., 2013. Priming memory invokes seed stress-tolerance. *Environ*
588 *Exp Bot*. 94, 33–45. <http://dx.doi.org/10.1016/j.envexpbot.2012.03.005>.

589 Eryılmaz, F., 2006. The relationships between salt stress and anthocyanin content in
590 higher plants. *Biotechnol Equip*. 20, 47–52.
591 <https://doi.org/10.1080/13102818.2006.10817303>

592 FAO, 2011. The state of the world’s land and water resources for food and agriculture
593 (SOLAW) – Managing systems at risk. Food and Agriculture Organization of the
594 United Nations, Rome and Earthscan, London.

595 Farooq, M., Hussain, M., Wakeel, A., Siddique, K.H.M., 2015. Salt stress in maize:
596 Effects, resistance mechanisms, and management. A review. *Agron Sustain Dev*.
597 35, 461–481. <https://doi.org/10.1007/s13593-015-0287-0>

598 Fitter, A.H., Stickland, T.R., 1991. Architectural analysis of plant root systems .
599 Influence of nutrient supply on architecture in contrasting plant species *New*
600 *Phytol*. 118, 383-389.

601 Giuffrè, A., 2013. HPLC-DAD detection of changes in phenol content of red berry
602 skins during grape ripening. *Eur Food Res Technol*. 237, 555–564. [https://doi.org/](https://doi.org/10.1007/s00217-013-2033-7)
603 [10.1007/s00217-013-2033-7](https://doi.org/10.1007/s00217-013-2033-7)

604 Gupta, S.A., Webb, R.P., Holaday, A.S., Allen, R.D., 1993. Overexpression of
 605 superoxide dismutase protects plants from oxidative stress. *Plant*
 606 *Physiol.* 103, 1067–1073.

607 Han, H., Gao, S., Li, B., Dong, X.C., Feng, H.L., Meng, Q.W., 2010. Overexpression
 608 of violaxanthin de-epoxidase gene alleviates photoinhibition of PSII and PSI in
 609 tomato during high light and chilling stress. *J. Plant Physiol.* 167, 176–183.
 610 <https://doi.org/10.1016/j.jplph.2009.08.009>

611 Ibrahim, E.A., 2016. Seed priming to alleviate salinity stress in germinating seeds. *J.*
 612 *Plant Physiol.* 192, 38–46. <http://dx.doi.org/10.1016/j.jplph.2015.12.011>

613 Imran, S., Afzal, I., Basra S.M.A., Saqib, M., 2013. Integrated seed priming with
 614 growth promoting substances enhances germination and seedling vigour of spring
 615 maize at low temperature. *Int J Agric Biol.* 15, 1251–1257.

616 Jamil M., Bashir S., Anwar S., Bibi S., Bangash A., Ullah F., Rha S. (2012) Effect of
 617 salinity on physiological and biochemical characteristics of different varieties of
 618 rice Pak. *J. Bot.* 44, 7-13.

619 Jollow, D.J., Mitchell, J.R., Zampaglione, N., Gillette, J.R., 1974. Bromobenzene
 620 Induced Liver Necrosis. Protective Role of Glutathione and Evidence for 3,4-
 621 bromobenzene Oxide as the Hepatotoxic Metabolites. *Pharmacol.* 11, 151-169.
 622 <http://dx.doi.org/10.1159/000136485>

623 Julkunen-Tiitto, R., 1985. Phenolic constituents in the leaves of northern willows:
 624 methods for the analysis of certain phenolics. *J Agric Food Chem.* 33 (2), 213–
 625 217. <https://doi.org/10.1021/jf00062a013>

626 Kangasjärvi, S., Neukermans, J., Li, S., Aro, E.M., Noctor G., 2012. Photosynthesis,
 627 photorespiration, and light signalling in defence responses. *J. Exp. Bot.* 63, 1619-
 628 36. <https://doi.org/10.1093/jxb/err402>

629 Lambrev P.H., Miloslavina Y., Jahns P., Holzwarth, A.R., 2012. On the relationship
 630 between non-photochemical quenching and photoprotection of Photosystem II
 631 *Biochim Biophys Acta.* 1817(5), 760-9. doi: 10.1016/j.bbabi.2012.02.002

632 Lichtenthaler, H. K., 1987. Chlorophylls and carotenoids: Pigments of photosynthetic
 633 biomembranes. *Methods Enzymol.* 148, 350-382. Maher, S., Fraj, H., Cherif, H.,
 634 2013. Effect of NaCl priming on seed germination of Tunisian fenugreek
 635 (*Trigonella foenum-graecum* L.) under salinity conditions *J Stress Physiol*
 636 *Biochem.* 9, 86-96.

637 Menezes-Benavente, L., Kernodle, S.P., Margis-Pinheiro, M., Scandalios, J.G., 2004.
638 Salt-induced antioxidant metabolism defenses in maize (*Zea mays* L.) seedlings.
639 Redox Rep. 9, 29–36. <https://doi.org/10.1179/135100004225003888>.

640 Munns, R., Tester, M., 2008. Mechanisms of salinity tolerance. *Annu Rev Plant Biol.*
641 59, 651–681 <http://dx.doi.org/10.1146/annurev.arplant.59.032607.092911>.

642 Muscolo, A., Junker, A., Klukas, C., Weigelt-Fischer, K., Riewe, D., Altmann, T.,
643 2015. Phenotypic and metabolic responses to drought and salinity of four
644 contrasting lentil accessions. *J. Exp. Bot.* 66, 5467–548.

645 Nakano, Y., Asada, K., 1981. Hydrogen peroxide is scavenged by ascorbate-specific
646 peroxidase in spinach chloroplasts. *Plant Cell Physiol.* 22, 867–880.

647 Nouman, W., Basra, S.M.A., Yasmeen, A., Gull T., Hussain S.B., Zubair, M., Gul, R.,
648 2014. Seed priming improves the emergence potential, growth and antioxidant
649 system of *Moringa oleifera* under saline conditions. *Plant Growth Regul.* 73, 267-
650 278. <https://doi.org/10.1007/s10725-014-9887-y>

651 Panuccio M.R., Sidari M., Muscolo A., 2003. Tolerance of Kikuyu grass to long term
652 salt stress is associated with the induction of antioxidant defences. *Plant Growth*
653 *Regul.* 41, 57-62.

654 Panuccio, M.R., Fazio, A., Papalia, T. Barreca, D., 2016. Antioxidant Properties
655 and Flavonoid Profile in Leaves of Calabrian *Lavandula multifida* L., an
656 Autochthon Plant of Mediterranean Southern Regions. *Chem Biodivers.* 13,
657 416 – 421. <https://doi.org/10.1002/cbdv.201500115>

658 Panuccio, M.R., Jacobsen, S.E., Akhtar, S.S., Muscolo, A., 2014. Effect of saline
659 water on seed germination and early seedling growth of the halophyte quinoa AoB
660 PLANTS 6, plu047. <https://doi.org/10.1093/aobpla/plu047>.

661 Papalia, T., Barreca, D., Panuccio M.R., 2017. Assessment of Antioxidant and
662 Cytoprotective Potential of *Jatropha* (*Jatropha curcas*) grown in Southern Italy. *Int*
663 *J Mol Sci.* 18(3), 660. <https://doi.org/10.3390/ijms18030660>

664 Paparella, S., Araujo, S., Rossi, G., Wijayasinghe, M., Carbonera, O., Balestrazzi, A.,
665 2015. Seed priming: State of the art and new perspectives. *Plant Cell Rep.* 34,
666 1281–1293. <https://doi.org/10.1007/s00299-015-1784-y>

667 Perveen, S., Shahbaz, M., Ashraf, M., 2012. Changes in mineral composition, uptake
668 and use efficiency of salt stressed wheat (*Triticum aestivum* L.) plants raised from
669 seed treated with triacontanol. *Pak J Bot.* 44, 27-35.

670 Ruban, A.V. , Johnson, M.P., Duffy, C.D.P., 2012. The photoprotective molecular
671 switch in the photosystem II antenna, *Biochim. Biophys. Acta* 1817, 167–181

672 Sen, A., Alikamanoglu, S., 2011. Effect of salt stress on growth parameters and
673 antioxidant enzymes of different wheat (*Triticum Aestivum* L.) varieties on in
674 vitro tissue culture. *Fresenius Envir. Bul.* 20(2), 489- 495.

675 Shahzad, M., Witzel, K, Zorb, C., Muhling, K.H., 2012. Growth-related changes in
676 subcellular ion patterns in maize leaves (*Zea mays* L.) under salt stress. *J Agron*
677 *Crop Sci.* 198, 46–56. <https://doi.org/10.1111/j.1439-037X.2011.00487.x>.

678 Tzortzakis, N.G., 2009. Effect of pre-sowing treatment on seed germination and
679 seedling vigour in endive and chicory. *Hortic Sci.* 36, 117-125.
680 <https://doi.org/10.17221/28/2008-HORTSCI>

681 Xu, L., Wang, P., Ali, B., Yang, N., Chen, Y., Wu, F., Xu, X., 2017. Changes of the
682 phenolic compounds and antioxidant activities in germinated adlay seeds. *J Sci*
683 *Food Agric.* 97, 4227–4234 <https://doi.org/10.1002/jsfa.8298>.

684 Zhang, S., Hu, J., Zhang, Y., Xie, X.J., Allen, K., 2007. Seed priming with
685 brassinolide improves lucerne (*Medicago sativa* L.) seed germination and seedling
686 growth in relation to physiological changes under salinity stress. *Aust J Agric Res.*
687 58(8), 811–815. <https://doi.org/10.1071/AR06253>.

688