



Università degli Studi Mediterranea di Reggio Calabria
Archivio Istituzionale dei prodotti della ricerca

In vitro effects of Italian *Lavandula multifida* L. leaf extracts on gilthead seabream (*Sparus aurata*) leucocytes and SAF-1 cells

This is the peer reviewed version of the following article:

Original

In vitro effects of Italian *Lavandula multifida* L. leaf extracts on gilthead seabream (*Sparus aurata*) leucocytes and SAF-1 cells / Fazio, A; Cerezuela, R; Panuccio, Maria Rosaria; Cuesta, A; Esteban, Má. - In: FISH & SHELLFISH IMMUNOLOGY. - ISSN 1095-9947. - 66:(2017), pp. 334-344. [10.1016/j.fsi.2017.05.033]

Availability:

This version is available at: <https://hdl.handle.net/20.500.12318/3393> since: 2020-11-25T18:11:14Z

Published

DOI: <http://doi.org/10.1016/j.fsi.2017.05.033>

The final published version is available online at: <https://www.sciencedirect>.

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website

Publisher copyright

This item was downloaded from IRIS Università Mediterranea di Reggio Calabria (<https://iris.unirc.it/>) When citing, please refer to the published version.

(Article begins on next page)

1 This is the peer reviewed version of the following article 2017. Angela Fazio,
2 Rebeca Cerezuela, Maria Rosaria Panuccio, Alberto Cuesta, Maria Ángeles
3 Esteban *In vitro* effects of Italian *Lavandula multifida* L. leaf extracts on gilthead
4 seabream (*Sparus aurata*) leucocytes and SAF-1 cells

5 which has been published in final doi <http://dx.doi.org/10.1016/j.fsi.2017.05.033>
6 (<https://www.sciencedirect.com/science/article/pii/S1050464817302784>)

7
8 The terms and conditions for the reuse of this version of the manuscript are specified in
9 the publishing policy. For all terms of use and more information see the publisher's
10 website

***In vitro* effects of Italian *Lavandula multifida* L. leaf extracts on gilthead seabream
(*Sparus aurata*) leucocytes and SAF-1 cells**

Angela Fazio^{1,2}, Rebeca Cerezuela², Maria Rosaria Panuccio³, Alberto Cuesta²,
Maria Ángeles Esteban^{2*}

¹Department of Innovation for Biological, Agrofood and Forestry Systems (DIBAF),
University of Tuscia, 01100, Viterbo, Italy

² Department of Cell Biology and Histology, Faculty of Biology, *Campus Regional de
Excelencia Internacional “Campus Mare Nostrum”*, University of Murcia. 30100.
Murcia, Spain

³ Department of Agriculture, University Mediterranea, 89122, Reggio Calabria, Italy

Corresponding Author:

*Maria Ángeles Esteban, Department of Cell Biology and Histology, Faculty of
Biology, *Campus Regional de Excelencia Internacional “Campus Mare Nostrum”*,
University of Murcia. 30100 Murcia, Spain.

E-mail address: aesteban@um.es

Abstract

Lavandula multifida is very appreciated by pharmaceutical and cosmetic industries. In Italy is only found in Calabria and Sicily and, at present, urge its valorization due to its high extinction and genetic erosion risks. Possible applications of *L. multifida* extracts on fish were assayed by using gilthead seabream (*Sparus aurata*) as a marine fish model, due to its importance in fish aquaculture. The *in vitro* effects of both aqueous and ethanolic leaf extracts obtained from two Italian populations of *L. multifida* on head kidney leucocyte activities (viability, phagocytosis, respiratory burst and peroxidase content) were compared with those originated from similar extracts obtained from a widely-distributed species of *L. angustifolia*. Furthermore, the possible cytotoxic effects of all extracts on SAF-1 cells and bactericidal effects for three pathogenic bacteria for fish (*Vibrio harveyi*, *Vibrio anguillarum*, *Aeromonas salmonicida*) were also evaluated.

Keywords: *Lavandula multifida* L.; Leaf extracts; Innate immune response; *Sparus aurata*; SAF-1 cell line.

1. Introduction

Lavandula (lavender, belongs to the family *Labiatae*, Lamiaceae) is a genus of 39 species, which has been largely used (either dried or as an essential oil) in folk and traditional medicine (Cavanagh and Wilkinson, 2002; Denner, 2009). Lavender has been used for centuries. Several studies revealed that *Lavandula* species are rich in a wide range of secondary metabolites such as phenolic compounds (Areias et al, 2000; Upson et al, 2000), monoterpenes, diterpenes (Politi et al, 2002), triterpenes (Topcu et al, 2001), sesquiterpenes (Ulubelen et al, 1988), and coumarins (Shimizu et al, 1990) which are supposed to be responsible for their antimicrobial, antifungal and antioxidant properties. Because of the high content of these secondary metabolites, Lamiaceae family accounts for beneficial and therapeutical properties which allow a wide spectrum of applications in food industry, cosmetic, perfumery and pharmaceutical preparations with high industrial and commercial value (Wilkinson et al, 2003).

Among *Lavandula* species, *Lavandula multifida* is a short-living plant 30 to 100 cm high, with a diploid genetic pool, equipped with triangular pinnatisect leaves, and blue or white purple flowers which give off a strong smell. It spontaneously grows along the Mediterranean coast of Egypt, Tunisia, Morocco, Algeria, Portugal and Spain, while in Italy it has been found only in Calabria and in Sicily (Galesi et al, 2005). The great interest for this plant is due to its essential oils and metabolites (whose antimicrobial, antifungal and antioxidant properties are well-documented) and to the wide possibilities of application in pharmaceutical and cosmetic industries (Ramchoun et al, 2009; Sosa et al, 2005; Zuzarte et al, 2012). Unfortunaltely, the human impact on its natural habitat caused the reduction and fragmentation of the populations of this plant in Southern Italy, and therefore, due to the rarity and the threats to this species, it was included in the “Regional Red Lists of Italian Plants” under the status IUCN of “critically endangered”

in Calabria region and “endangered” in Sicily (Conti et al, 1997). For this important reason, the valorization of *L. multifida* is strictly necessary due to its high extinction and genetic erosion risks.

Currently, there is a growing interest in screening medicinal plants extracts for their bactericidal, fungicidal and even immunostimulant properties, in order to exploit new biocompounds of natural origin which could be employed in the prevention and/or control of fish diseases in aquaculture (Reverter et al, 2014). The importance of using immunostimulants in aquaculture is known since many years (Sakai, 1999) and, at present, it is a topic of great interest in research. In the intensive aquaculture system, application of antibiotics and chemotherapeutics as prophylactic measures has been widely criticized for their negative impacts on the environment and also on fish, such as immunodepression or residue accumulation in tissues, and besides this leads to the development of drug resistant pathogens (Rijkers et al, 1980; Harikrishnan et al, 2009; FAO, 2003; Smith et al, 1994). The use of plant extracts as immunostimulants has recently received increasing attention not only because they combine lower costs with the low impact on the environment, but also because they may have additional physiological effects since they contain many nutrients, micronutrients as well as other immunostimulant substances (Cuesta et al, 2005). Taking in considerations the needed of looking for new applications of *L. multifida* extracts, and taking into account the importance of gilthead seabream (*Sparus aurata*) in Mediterranean and marine fish aquaculture, we developed the present study. The *in vitro* effects of leaf extracts obtained from two Italian populations of *L. multifida* on head kidney leucocytes activities (viability, phagocytosis, respiratory burst and peroxidase activity) were tested, making a comparison with similar extracts obtained from the widely-distributed species *L. angustifolia*. Furthermore, the cytotoxic activity on SAF-1 cells (a cell line obtained

from gilthead seabream) and the bactericidal activity of the extracts on *Vibrio harveyi*,
Vibrio anguillarum and *Aeromonas salmonicida* were also checked.

2. Materials and Methods

2.1. Plant extracts

Leaves from Sicilian *L. multifida* L. (collected from Brucoli, Siracusa, indicated in the
figures as *L. multifida* S), from Calabrian *L. multifida* L. (collected from Capo
dell'Armi, indicated in the figures as *L. multifida* C), and from *L. angustifolia* Miller
were collected and air-dried at 50 °C until the weight was stable.

For preparation of aqueous extracts, air-dry leaves were macerated and shaken with
boiling water for 4 h at 25°C, then the mixture was filtered twice using nylon net filter
with a 100µm pore size, and evaporated in a rotary evaporator until dryness. Prior to be
used in the assays, extracts were filtered using sterile filters of 0.22 µm of diameter. For
preparation of ethanol extracts, air-dry leaves were macerated and shaken with pure
ethanol (1:40, 48 h, 25 °C). The resulting mixture was then filtered-twice as described
above, and concentrated by vaporizing using a rotary evaporator.

2.2. Animals

Thirty specimens (40.51 ± 1.47 gr weight) of the seawater teleost gilthead seabream (*S.*
aurata L.), obtained from a local farm (Murcia Spain), were kept in re-circulating
seawater aquaria (250 L) in the Marine Fish Facility at the University of Murcia. The
water temperature was maintained at 20 ± 2 °C with a flow rate of 900 L h⁻¹ and 28‰
salinity. The photoperiod was 12 h light: 12 h dark. Fish were allowed to acclimatise for
15 days before the start of the trial, where they were fed with a commercial pellet diet
(Skretting, Spain) at a rate of 2% body weight day⁻¹. The fish were killed after starving

for 24 h by using an overdose of MS-222 (Sandoz, 100 mg ml⁻¹ water). All experimental protocols were approved by the Ethical Committee of the University of Murcia (Permit Number A13150104).

2.3. Head-kidney leucocyte isolation and incubation with extracts

Before the dissection of the head-kidney (HK), the specimens were bled. Blood was collected from the caudal vein and afterwards fish were dissected to obtain HK fragments. For isolation of leucocytes to carry out the assays, HK fragments were transferred to 8 ml of sRPMI [RPMI-1640 culture medium (Gibco) supplemented with 0.35% sodium chloride (to adjust the medium's osmolarity to gilthead seabream plasma osmolarity of 353.33 mOs), 3% fetal calf serum (FCS, Gibco), 100 i.u. ml⁻¹ penicillin (Flow) and 100 mg ml⁻¹ streptomycin (Flow)] (Esteban et al., 1998). Cell suspensions were obtained by forcing fragments of the organ through a nylon mesh (mesh size 100 µm), washed twice (400 g, 10 min), counted (Z2 Coulter Particle Counter) and adjusted to 2 x 10⁷ cells ml⁻¹ in sRPMI. Cell viability was higher than 98%, as determined by the trypan blue exclusion test.

To study the possible effects of water and ethanolic extracts on leucocyte activities, aliquots of 50 µl of the obtained leucocytes suspension containing 2 x 10⁷ cells ml⁻¹ were dispensed into flat-bottomed 96-well microtitre plates (Nunc). Then aliquots of 50 µl well⁻¹ of water or ethanolic extracts ranging from 10, 100, 500, to 1000 µg ml⁻¹ prepared in sRPMI were added. The extract aliquot was replaced by sRPMI for control samples for assays with water extracts, or by 0,1% dymethyl sulfoxide (DMSO, Sigma) in RPMI in case of control samples for assays with ethanolic extracts. Cells were incubated in presence of the extracts for 24 h at 20 °C in an incubator with 5% CO₂ and 85%

humidity. After incubation, leucocyte viability and phagocytic, respiratory burst and peroxidase activities were determined as described below.

2.4. *Leucocyte viability*

Aliquots of 100 μ l of leucocytes previously incubated for 24h without (control) or with the plant extracts were placed in 5 ml glass tubes (Falcon, Becton–Dickinson) and 40 μ l of propidium iodide (PI) (400 μ g ml⁻¹, Sigma) were added to each sample. The tubes were gently mixed before analysis in a FACScan (Becton–Dickinson, Madrid, Spain) flow cytometer with an argon-ion laser adjusted to 488 nm. Analyses were performed on 5000 cells, which were acquired at a rate of 300 cells s⁻¹. Data were collected in the form of two-parameter side scatter (granularity, SSC) and forward scatter (size, FSC), and green fluorescence (FL1) and red fluorescence (FL2) dot plots or histograms were made on a computerised system. Dead cells were estimated as the percentage of cells with propidium iodide (red-PI fluorescent cells).

2.5. *Phagocytic activity*

The phagocytic activity of gilthead seabream HK leucocytes was studied by flow cytometry according to Esteban et al. (1998). Heat killed (30 min, 60 °C) lyophilised *Saccharomyces cerevisiae*, strain S288C, were washed twice, counted and adjusted to 108 yeast cells/ml in sRPMI-1640. To label yeast cells with fluorescein isothiocyanate (FITC, Sigma) yeast cells were incubated with 5 μ g ml⁻¹ FITC at 22 °C with constant stirring (40 cycles min⁻¹) and in darkness for 15 min (Rodriguez et al, 2003). After labeling, free FITC was removed by washing twice in phosphate buffer saline (PBS) and the yeast cells were resuspended in sRPMI-1640. FITC-labeled yeast cells were acquired for flow cytometric study. The staining uniformity was examined and then the yeast cell suspensions were aliquoted and stored at 4 °C.

Samples of 125 µl of medium containing FITC-labeled-yeast cells and 100 µl of HK leucocytes in sRPMI were mixed, centrifuged (400 g, 5 min, 25°C), resuspended and incubated at 25 °C for 60 min in dark conditions. At the end of the incubation time, the samples were placed on ice to stop phagocytosis and 400 µl of cold PBS was added to each sample. The fluorescence of the extracellular yeast cells (i.e. free yeast cells and yeast cells adhered to phagocytes but not ingested) was quenched by adding 50 µl of cold Trypan Blue (0.4% in PBS) per sample. Standard samples of FITC-labelled *S. cerevisiae* yeast cells or HK leucocytes were included in each phagocytosis assay. Immediately, the samples were mixed gently, acquired, and analysed in a FACScan. Data were collected in the form of two-parameter side scatter (SSC) and forward scatter (FSC), and green fluorescence (FL1) dot plots or histograms were made on a computerized system. Fluorescence histograms represented relative fluorescence on a logarithmic scale. Flow cytometer was set to analyze the phagocytic cells gated from all the leucocytes because of their higher SSC and FSC values. Phagocytic ability was defined as the percentage of cells with one or more ingested yeast cells (green-FITC fluorescent cells) within the phagocytic cell population. The relative number of ingested yeast cells per cell (phagocytic capacity) was assessed in arbitrary units from the mean fluorescence intensity of the phagocytic cells.

2.6. Respiratory burst activity

The respiratory burst activity of seabream HK leucocytes was studied by a chemiluminescence method (Bayne and Levy, 1991). Samples of 100 µl of a PMA/luminol solution [1 ng ml⁻¹ phorbol myristate acetate (PMA, Sigma) and 10⁻⁴M luminol (Sigma) in HBSS with Ca²⁺ and Mg²⁺] were added to the HK leucocytes (previously incubated as described above). The plate was shaken and immediately read in a chemiluminometer (BMG, Fluoro Star Galaxy). Measurements were performed in 30

cycles of 2 minutes each. The kinetics of the reactions were analyzed and the maximum slope of each curve calculated. Control samples containing leucocytes that had not been incubated with the extracts were also analyzed.

2.7. Peroxidase content

The total peroxidase content of HK leucocytes was measured according to Quade and Roth (1997). To do this, 5 µl of HK leucocytes (previously incubated as described above) were incubated for 10 min with 0.02% cetyltrimethylammonium bromide (CTAB, Sigma) at 60 rpm. Afterwards, 100 µl of 10mM 3,3',5,5'-tetramethylbenzidine hydrochloride (TMB, Sigma) and 5mM H₂O₂ (both substrates prepared daily) were added and after 2 min, 50 µl of 2M sulfuric acid was also added to stop the reaction. The absorbance of the samples was measured at 450 nm in a microplate reader (BMG Fluostar Omega, USA). Control samples containing leucocytes that had not been incubated with extracts were also analyzed.

2.8. SAF-1 cell culture

The established cell line SAF-1 (ECACC n 00122301) was seeded in 25 cm² plastic tissue culture flasks (Nunc, Germany) cultured in L-15 Leibowitz medium (Life Technologies, UK), supplemented with 10% fetal bovine serum (FBS, Life Technologies), 2mM L-glutamine (Life Technologies), 100 i.u. ml⁻¹ penicillin (Life Technologies) and 100 µg ml⁻¹ streptomycin (Life Technologies). Cells were grown at 25 °C and with 85% humidity. Exponentially growing cells were detached from culture flasks by brief exposure to 0.25% of trypsin in PBS, pH 7.2-7.4, according to the standard trypsinization methods. The detached cells were collected by centrifugation (1000 x g, 5 min, 25 °C) and the cell viability was determined by the trypan blue exclusion test.

2.9. Cytotoxicity assay on SAF-1 cell line

Cytotoxicity assay was performed in quadruplicates. When SAF-1 cell lines were approximately 80% confluent, they were detached from flasks culture with trypsin (as described before), and aliquots of 100 μL containing 50000 cells well^{-1} were dispensed in 96-well tissue culture plates and incubated (24 h, 25 °C). This cell concentration was previously determined in order to obtain satisfactory absorbance values in the cytotoxic assay and avoided cell overgrowth. After that, the culture medium was replaced by 100 $\mu\text{L well}^{-1}$ of the extracts to be tested at the appropriate dilution. Tested concentrations of water and ethanol extracts ranged from 1 to 1000 $\mu\text{g mL}^{-1}$ (1, 10, 100, 1000). Cells were then incubated for 24 h. Control samples received the same volume of culture medium (for water extracts) or of DMSO 0,1% (for ethanolic extracts). Cells were incubated for 24h at 25°C and then their viability determined using the MTT assay.

The MTT assay is based on the reduction of the yellow soluble tetrazolium salt (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) (MTT, Sigma) into a blue, insoluble formazan product by the mitochondrial succinate dehydrogenase (Berridge and Tan, 1993; Denizot and Lang, 1986). After incubation with the leaf-extracts, SAF-1 cells with phosphate buffer saline solution (PBS) and 200 μLwell^{-1} of MTT (1 mg mL^{-1}) were added. After 4 h of incubation, the cells were washed and the formazan crystals solubilized with 100 $\mu\text{L well}^{-1}$ of DMSO. Plates were shaken for 5 minutes in dark conditions and the absorbance at 570 nm and 690 nm determined in a microplate reader.

2.10. Bacteria

Three pathogenic bacteria for fish (*V. harveyi*, *V. anguillarum* and *A. salmonicida*) and *Escherichia coli*, as control, were used in the bactericidal assays. All bacterial strains were grown from 1 mL of stock culture that had been previously frozen at -80°C. *V. harveyi*, *V. anguillarum* and *A. salmonicida* were cultured for 48 h at 25 °C in Tryptic Soy Agar (TSA, Difco Laboratories), and then inoculated in Tryptic Soy Broth (TSB,

Difco Laboratories), both supplemented with NaCl to a final concentration of 1% (w/v). Bacteria in TSB medium were then cultured at the same temperature, with continuous shaking (100 rpm) during 24 h. *E. coli* was cultured in Luria Bertani Agar (LB Agar, Difco) for 48 h at 37 °C and then inoculated in Luria Bertani Broth (LB Broth, Difco). *E. coli* bacteria in LB broth medium were then cultured for 24 h at 37 °C with continuous shaking (100 rpm). Exponentially growing bacteria were resuspended in sterile Hank's balanced salt solution (HBSS) and adjusted to 1×10^8 colony forming units (cfu) ml⁻¹.

2.11. Bactericidal assay

Bactericidal activity was determined following the method of Stevens et al (1991) with some modifications. Samples of 20 µl of water or ethanolic leaf-extracts previously adjusted to 20, 100, 200, 500, 1000 µg ml⁻¹ were added in quadruplicate wells of a U-shaped 96-well plate (Nunc). Hank's balanced solution was added to some wells instead of the extracts and served as positive control. Aliquots of 20 µl of the bacteria previously cultured were added and the plates were incubated for 2.5 h at 25°C (in case of *V. harveyi*, *V. anguillarum* and *A. salmonicida*), or at 37°C (in case of *E. coli*). After that, 25 µl of MTT (1mg ml⁻¹) were added to each well and the plates were newly incubated for 2 h (at the appropriate temperature taken into account the assayed bacteria) to allow the formation of formazan. Plates were then centrifuged (2000 x g, 10 min), being the precipitates dissolved in 200 µl of DMSO and transferred to a flat-bottom 96 well-plate. The absorbance of the dissolved formazan was measured at 560 nm. Bactericidal activity was expressed as percentage of no viable bacteria, calculated as the difference between absorbance of bacteria surviving compared to the absorbance of bacteria from positive controls (100%).

2.12. Statistical analysis

Results showed are representative of at least three independent experiments and are expressed as Mean \pm Standard Error. All assays related to leucocyte activities were performed in duplicate and results were expressed as Mean \pm Standard Error for each group (three fish per group). Data were analyzed by one-way analysis of variance (ANOVA), and Tukey post-hoc test was performed in order to make a multiple comparison between experimental groups. Differences were considered statistically significant when $P < 0.05$.

3. Results

3.1. Effects of leaves extracts on gilthead seabream head kidney leucocyte

Head kidney leucocyte viability was tested after 24 h of incubation with water or ethanolic extracts of the three tested plants in the range of concentrations from 10 to 1000 $\mu\text{g ml}^{-1}$ (10, 50, 100, 250, 500, 750, 1000 $\mu\text{g ml}^{-1}$). Results demonstrated that any significant effect was observed on HK leucocyte viability after incubation with the cited extracts, respect to the values recorded for control HK leucocytes (Fig. 1).

Regarding phagocytosis, incubation of HK leucocytes with water or ethanolic extracts of *L. multifida* (from both Calabrian and Sicilian populations) significantly enhanced their phagocytic ability when using the range of extract concentrations from 10 to 500 $\mu\text{g ml}^{-1}$ and 10 to 100 $\mu\text{g ml}^{-1}$, respectively (Fig. 2). However, after incubation of HK leucocytes with *L. angustifolia* water extracts significant enhancements of the phagocytic ability were only recorded when using highest concentrations (from 100 to 1000 $\mu\text{g ml}^{-1}$) (Fig. 2a). On the other hand, incubation of leucocytes with ethanolic extracts from *L. angustifolia* did not affect their phagocytic ability.

On the contrary, significant decreases were observed in the phagocytic ability of HK leucocytes after being incubated with 500 and 1000 $\mu\text{g ml}^{-1}$ of ethanolic extracts from all plant tested, except for 500 $\mu\text{g ml}^{-1}$ ethanolic extract of *L. multifida* S which increased their phagocytic ability (Fig. 2b).

A similar trend was observed in the phagocytic capacity because HK leucocytes incubated with 10 $\mu\text{g ml}^{-1}$ and 10 to 100 $\mu\text{g ml}^{-1}$ water and ethanolic extracts from *L. multifida* S, respectively, showed an increased phagocytic capacity (Fig. 3). Instead, water and ethanolic extracts from *L. angustifolia* did not affect significantly the phagocytic capacity of gilthead seabream HK leucocytes, respect to control samples.

Regarding respiratory burst, incubation of HK leucocytes with water extracts from all populations of *Lavandula* tested increased this activity at all concentrations tested in a significant manner, in comparison with the control group, except for *L. multifida* S water extract which decreased significantly this activity when using at 1000 $\mu\text{g ml}^{-1}$ of concentration (Fig. 4). Also, ethanolic extracts from both populations of *L. multifida* used at 10 and 100 $\mu\text{g ml}^{-1}$ increased the HK leucocyte respiratory burst activity, while the ethanolic extracts from all plants tested at 1000 $\mu\text{g ml}^{-1}$ significantly decreased the respiratory burst activity of HK leucocytes compared to the control samples (Fig. 4). The observed enhancements of the respiratory burst activity of leucocytes was always higher after incubation with water extracts than with ethanolic extracts.

No significant variations in the peroxidase content of HK leucocytes were observed after being incubated neither with aqueous nor with ethanolic extracts from the three plants tested in the present work, respect to the values recorded for control samples (Fig. 5).

3.2. Effects of leaves extracts on SAF-1 cell line

The effects of water and ethanolic extracts of *L. multifida* on cytotoxicity of SAF-1 cells were also evaluated. Results from the cytotoxicity test showed that water extracts from all plants tested did not alter significantly the cell viability, respect to values for control samples (Fig. 6). Interestingly, incubation of SAF-1 cells with *L. multifida* S water extract increased the cell viability in a significant manner at the higher concentration tested (1000 $\mu\text{g ml}^{-1}$) (Fig. 6a).

On the contrary, incubation of SAF-1 cells with ethanolic extracts significantly affect their viability, indeed showing high levels of cytotoxicity at the higher concentration tested (1000 $\mu\text{g ml}^{-1}$) (Fig. 6b). Particularly, ethanolic extracts from the two populations of *L. multifida* had a significantly higher level of cytotoxicity on SAF-1 cells if compared to the cytotoxicity provoked on these cells after being incubated with *L. angustifolia* ethanolic extracts (Fig. 6).

3.3. Bactericidal activity of leaves extracts

In this study *L. multifida* plant extracts were also tested (in comparison with the extracts from the commercial species *L. angustifolia*) for their bactericidal activity against some fish pathogens relevant in fish aquaculture: *V. harveyi*, *V. anguillarum* and *A. salmonicida*, being *E. coli* used as control bacteria. Results from bactericidal assays revealed that any of the assayed plant extracts neither water nor ethanolic affect bacteria viability in a significant manner (Figs. 7-10). Only a slight difference in bacteria viability (80% cell viability) was observed when ethanolic extracts from *L. multifida* C were used to incubate *V. harveyi* bacteria cells (Fig. 7).

4. Discussion

The massive use of antimicrobials for disease control of farmed animals is considered, at present, a real hazard. Indeed food-producing animals are one of several potential sources of antibiotic-resistant bacteria which may spread from animals to man via the food chain (Harikrishnan et al, 2011). Furthermore, pathogens may also transfer their antibiotic-resistance genes into human pathogenic bacteria thus posing a threat to human health and considered as one of the main environmental problems (Alderman and Hastings, 1998; Cabello, 2006; Abutbul et al, 2004; Smith et al, 1994; MacMillan, 2001).




Fig. 1. Percentage of viability of *S. aurata* HK leucocytes after 24 h of incubation with water (a) or ethanolic extracts (b) obtained from leaves of Calabrian and Sicilian *Lavandula multifida*. *L. angustifolia* was considered as control plant. Results of cytotoxicity are given as the percentage of viability, related to the control of untreated cells (the mean optical density of untreated cells was set to 100% viability). Results showed are representative of at least three independent experiments and are expressed as Mean \pm Standard Error.

Regarding farmed fish, while some immunostimulants cannot be used due to their high cost and/or limited effectiveness (Ringo et al, 2012), treatment with medicinal plants is gaining attention due to the fact that they may be a potentially beneficial alternative to antibiotics. Firstly, these plants mitigate many of the side effects associated with the use of chemicals and antibiotics (Madhuri et al, 2012). Furthermore, treatment of bacterial diseases with different herbs has been safely used in organic agriculture, veterinary and human medicine (Madhuri et al, 2012; Bairwa et al, 2012).




Fig. 2. Percentage of phagocytic cells of *S. aurata* HK leucocytes after 24 h of incubation with water (a) and ethanolic extracts (b) obtained from leaves of Calabrian and Sicilian *Lavandula multifida*. *L. angustifolia* was considered as control plant. Small letters indicate significant differences between different concentrations of the same plant extract. Capital letters indicate significant differences between different plant extracts at the same treatment concentration. Differences were considered significant when $P < 0.05$. Results showed are representative of at least three independent experiments and are expressed as Mean \pm Standard Error.




Fig. 3. Phagocytic capacity of *S. aurata* HK leucocytes after 24 h of incubation with water (a) and ethanolic extracts (b) obtained from leaves of Calabrian and Sicilian *Lavandula multifida*. *L. angustifolia* was considered as control plant. Small letters indicate significant differences between different concentrations of the same plant extract. Capital letters indicate significant differences between different plant extracts at the same treatment concentration. Differences were considered significant when $P < 0.05$. Results showed are representative of at least three independent experiments and are expressed as Mean \pm Standard Error.






Fig. 4. Respiratory burst activity (expressed as slope/minute) of *S. aurata* HK leucocytes after 24 h of incubation with water (a) and ethanolic extracts (b) obtained from leaves of Calabrian and Sicilian *Lavandula multifida*. *L. angustifolia* was considered as control plant. Small letters

390 indicate significant differences between different concentrations of the same plant extract.
 391 Capital letters indicate significant differences between different plant extracts at the same
 392 treatment concentration. Differences were considered significant when $P < 0.05$. Results
 393 showed are representative of at least three independent experiments and are expressed as Mean
 394 \pm Standard Error



395 Fig. 5. Peroxidase activity (expressed as unit enzyme) of *S. aurata* HK leucocytes after 24 h of
 396 incubation with water (a) and ethanolic extracts (b) obtained from leaves of Calabrian and
 397 Sicilian *Lavandula multifida*. *L. angustifolia* was considered as control plant. Small letters
 398 indicate significant differences between different concentrations of the same plant extract.
 399 Capital letters indicate significant differences between different plant extracts at the same
 400 treatment concentration. Differences were considered significant when $P < 0.05$.
 401 Results showed are representative of at least three independent experiments and are expressed
 402 as Mean \pm Standard Error.



405 Fig. 6. Viability (expressed as percentage) of SAF-1 cell line after 24 h of incubation with water
 406 (a) and ethanolic extracts (b) obtained from leaves of Calabrian and Sicilian *Lavandula*
 407 *multifida*. *L. angustifolia* was considered as control plant. Small letters indicate significant
 408 differences between different concentrations of the same plant extract. Capital letters indicate
 409 significant differences between different plant extracts at the same treatment concentration.
 410 Differences were considered significant when $P < 0.05$. Results showed are representative of at
 411 least three independent experiments and are expressed as Mean \pm Standard Error.




Fig. 7. Bactericidal activity of water (fig. 7a) and ethanolic extracts (fig. 7b) from Calabrian and Sicilian *L. multifida* (indicated in the figure as C and S respectively) and from *L. angustifolia* on *Vibrio harveyi*. Small letters indicate significant differences between different concentrations of the same plant extract. Differences were considered significant when $P < 0.05$. Results showed are representative of at least three independent experiments and are expressed as Mean \pm Standard Error.




Fig. 8. Bactericidal activity of water (fig. 8a) and ethanolic extracts (fig. 8b) from Calabrian and Sicilian *L. multifida* (indicated in the figure as C and S respectively) and from *L. angustifolia* on *Vibrio anguillarum*. Small letters indicate significant differences between different concentrations of the same plant extract. Differences were considered significant when $P < 0.05$. Results showed are representative of at least three independent experiments and are expressed as Mean \pm Standard Error.




Fig. 9. Bactericidal activity of water (fig. 9a) and ethanolic extracts (fig. 9b) from Calabrian and Sicilian *L. multifida* (indicated in the figure as C and S respectively) and from *L. angustifolia* on *Aeromonas salmonicida*. Small letters indicate significant differences between different concentrations of the same plant extract. Differences were considered significant when $P < 0.05$. Results showed are representative of at least three independent experiments and are expressed as Mean \pm Standard Error.




Fig. 10. Bactericidal activity of water (fig. 10a) and ethanolic extracts (fig. 10b) from Calabrian and Sicilian *L. multifida* (indicated in the figure as C and S respectively) and from *L. angustifolia* on *Escherichia coli*. Small letters indicate significant differences between different concentrations of the same plant extract. Differences were considered significant when $P < 0.05$. Results showed are representative of at least three independent experiments and are expressed as Mean \pm Standard Error.

Lately, the use of new and non-purified immunostimulants has received many attentions, and an essential prerequisite is that any new potential immunostimulant should always be respectful with the environment (Sakai et al, 1999). Currently increased consumer demand of quality in farmed fish and shellfish, news efforts in order to avoid the possible pollutants, use of antibiotics and/or carcinogens during the production process urged. In this scenario, plants or their byproducts are preferred since they content numerous compounds like phenols, polyphenols, alkaloids, quinones, terpenoids, lectines and polypeptides many of which have been shown to be very effective alternatives to antibiotics, chemicals, vaccines, and other synthetic compounds (Harikrishnan et al, 2011). Since herbs have little side effects and are easily degradable, various studies have investigated the effect of plant products on innate and adaptive immune response and to prevent and control fish and shellfish diseases (reviewed by Valladão et al, 2015). Fruits, leaves, seeds, flowers, roots and barks extracts from many plants have been reported to have immunomodulator porperties (NagaPreethi and Rajeshwari, 2014). Concomitantly, natural plant products promote different activities in animals like growth, appetite stimulation, immunostimulation, aphrodisiac, antistress and antimicrobial properties due to the already mentioned numerous active principles

present in them (Reverter et al, 2014). The necessity of looking for new natural substances with potential application in fish aquaculture due to their immunostimulant properties or biocides activities against potential pathogenic microbes move us to carry out the present work.

Among plants belonging to *Lavandula* genus, *L. multifida* represents a remarkable species because of its wide utilization in traditional medicine since ancient times. Most of the studies performed to date focused on the antibacterial, antifungal and antioxidant properties of its essential oils, as well as on their biochemical constituents.

It is well known that biological and therapeutic properties of medicinal plants are closely related to their chemical compounds; due to the differences in solubility in water or in alcohol, each class of these chemical compounds has a preferred effective extraction method which facilitate getting out the chemicals out of the plant and into the herbal remedy that is being prepared, and this is the reason why water and ethanolic extracts are obtained in order to analyze the different biological and therapeutic properties (Hashemi et al, 2008). Many plant species are known to synthesize many bioactive secondary metabolites like terpenoids, alkaloids, glycoside, phenolic compounds, polysaccharides, flavonols and tannins which possess various biological properties. Furthermore, it is well-known that phytomedicines are inexpensive, eco-friendly and easy to be prepared (Hashemi et al, 2008).

Although actually there are no available data concerning the effects of lavender extracts on fish immunity and their possibility of use as immunostimulant in aquaculture, some scientific evidences exist about the effect of extracts from other plants belonging to Lamiaceae family on innate immune response. Extracts from *Thymus guyonii*, *Salvia verbenaca* and *Stachys circinata* (Lamiaceae family) have been tested for their immunostimulant potential on the phagocytic activity in adult male mice (*Mus*

musculus) through the carbon clearance rate test, resulting in an appreciable increasing in phagocytic index in animals when administered plant extracts by intraperitoneal injection (Nassar et al, 2015). Also, leaf extracts from *Ocimum sanctum* (Lamiaceae family) have been tested for their effect on the innate and specific immune responses and disease resistance against *Aeromonas hydrophila* in *Oreochromis mossambicus*, resulting in an important immunostimulatory effect when leaf extracts were administered orally and intraperitoneally (Logambal et al, 2000). Similarly, dietary administration of peppermint (*Menta piperita*, Lamiaceae family) promoted growth performance and increased the main hematological and immune humoral parameters in fry Caspian white fish (*Rutilus frisii kutum*) (Adel et al, 2015). Furthermore *Zataria multiflora* (Lamiaceae) exerted immunostimulatory effects on some immunological factors such as antibody titers, total white blood cells and serum bactericidal activity when its essential oils were dietary administered in *Cyprinus carpio* (Soltani et el, 2010). Present results corroborate, firstly that both aqueous and ethanolic extracts of *L. multifida* have any negative impact on HK leucocyte viability at the concentrations and incubation times tested in the present study. Furthermore, and more interesting, our results demonstrate that *L. multifida* L. extracts increased HK leucocytes phagocytosis and respiratory burst activities, thus supporting the possibility of use of *L. multifida* L. extracts as immunostimulant in the production of this important farmed marine fish species. Also, respiratory burst activity showed a significant increase when HK leucocytes were treated with water and ethanolic extracts from all populations of Lavandula, although a negative effect was observed when using ethanolic extracts at the maximum concentration tested (1000 mg mL⁻¹), being this similarly observed for the phagocytic ability. Present results show that *L. multifida* L. extracts positively affect phagocytosis and respiratory burst activity of gilthead seabream HK leucocytes,

without any significant effect neither on HK leucocytes viability nor on HK leucocytes peroxidase, thus supporting the possibility of use of *L. multifida* L. extracts, in a well-defined range of concentrations, as immunostimulant in the production of this important farmed marine fish species. Immunostimulants are capable of promoting a greater and more effective sustained immune response to those infectious agents (viruses, bacteria, fungi, and parasites), producing subclinical disease without risks of toxicity, carcinogenicity or tissue residues (Muthusamy et al, 2013). Future in vivo studies could demonstrate if similar immunostimulant effects are induced after dietary administration of such extracts to gilthead seabream specimens.

Fish cell lines are increasingly important research tools as an alternative to the experimental animals. The long-term SAF-1 cell line was established in 1996 and it is a fibroblast-like culture derived from gilthead seabream (Béjar et al., 1997). Since then, the SAF-1 cell line has proved useful in many applications and varied studies related to virology (Pérez-Prieto et al., 1999; Tafalla et al., 2004; Alonso et al., 2005; Bandín et al., 2006; García Rosado et al., 2008; Cano et al., 2016), bacteriology (Acosta et al., 2009; El Aamri et al., 2012), immunology (Ray et al., 2002; Pelegrín et al., 2004), and more recently in nanoparticles interiorization (Trapani et al., 2015), metal toxicity (Morcillo et al., 2016) or freshness and freeze-thawing of seabream fillets (Diop et al., 2016) studies. Curiously, to the best of our knowledge, only one previous paper focused on the immune modulatory effects of a plant extract (*Aloe arborescens*) on SAF-1 cells (Picchietti et al., 2013). In the study, SAF-1 cells were treated with Aloe extract at different concentrations ($1.2\text{--}4.8\text{ mg ml}^{-1}$) and times (24-72 h). The Aloe extract (1.2 mg ml^{-1}) induced a synergic effect when cells were also stimulated with LPS- or poly I:C which was detected by studying some immune-related gene expression (Picchietti et al., 2013).

Present results from the cytotoxic assay developed on SAF-1 cell line after incubation with *L. multifida* extracts demonstrate no significant effects on cell viability in a limited range of concentrations tested. However, the maximum doses used in our experiments (1000 mg mL⁻¹) affected their viability in a significant manner, causing an increase or a decrease in this parameter when SAF-1 cells were incubated with water or ethanolic extracts, respectively. These data agreed with results obtained from the determination of phagocytosis and respiratory burst activities: indeed ethanolic extracts from *L. multifida* C and S provoked a significant decrease in the number of yeast cells ingested after 24 h of incubation at 1000 mg mL⁻¹ of concentration, furthermore also the percentage of phagocytic cells showed a significant decrease when HK leukocytes were incubated with ethanolic extracts from all plants tested during 24 h at 1000 mg mL⁻¹ of concentration, and a similar trend was observed also for the respiratory burst activity. Although the main components of the leaves extract have not been studied yet, the analysis of the chemical composition is one of our future next step. These could allow to know if differences in the biochemical components between water and ethanolic extracts exist, which may explain the different properties of the extracts.

Finally, present results demonstrated that *L. multifida* L. extracts had no bactericidal activity against the fish pathogens tested. There are no many available data about the antibacterial activity of *L. multifida* L. extracts until now, and the majority of antibacterial studies have been performed on its essential oils. As reported by Khadir and colleagues (2016), *L. multifida* essential oils were tested against methicillin-resistant *Staphylococcus aureus* (MRSA) using disc diffusion method, revealing a good anti-MRSA activity, whereas the ethanolic extract was less active, thus suggesting that different biological effects between leaf extracts and essential oils could be due to differences in their chemical composition. In fact the chemical composition of *L.*

multifida essential oils studied by GC and GC-MS (Khadir et al, 2016) revealed that carvacrol is the main component (from 27.5% to 57%), being carvacrol a monoterpenic phenol responsible for many biological activities such as antimicrobial, antitumor, antimutagenic, antigenotoxic, analgesic, antiinflammatory, antiparasitic, antiplatelet (Baser, 2008). Instead, in a recent work ethanolic leaf extract of *L. multifida* L. from Southern Italy was screened by UV spectroscopic analysis (Fazio et al, 2014), revealing the presence of vitexin and flavones glucoside derivatives of hypolaetin, scutellarein, luteolin, isoscutellarein, apigenin and chrysoeriol, which were reported to possess mainly antiinflammatory, anticancer and antioxidant properties (Alcaraz and Hoult, 1985; Lin et al, 2008; Saeed et al, 2015; Choi et al, 2005). On the other hand, opposite results were obtained with ethanolic extracts from *Lavandula officinalis*, *Melissa officinalis*, *Ocimum basilicum*, *Origanum vulgare*, *Rosmarinus officinalis* and *Salvia officinalis*, all belonging to Lamiaceae family, which were tested for antimicrobial activity, exhibiting a broad spectrum of inhibitory effects on some fish pathogens like *L. anguillarum*, *Y. ruckeri*, *P. damsela*, *L. garvieae* (Bulfon et al, 2014).

To conclude, *L. multifida* L. water and ethanolic leaf extracts up to 100 mg mL⁻¹ of concentration increased innate immune activities of *S. aurata* HK leucocytes, more concretely, their phagocytic and respiratory burst activities, but did not exert any bactericidal activity on the bacterial strains tested in the present study, and did not affect significantly SAF-1 cells viability up to 100 mg mL⁻¹ of concentration. Although the extracts did not exert any bactericidal activity directly on the assayed bacteria strains, present results suggest the possibility of use such extracts in in vivo studies in order to corroborate if could be possible using those extracts in aquaculture in order to achieve protection against pathogenic infections through enhancement of the innate immunity of fish.

583

584 **References**

- 585 (1) Denner SS (2009) "Lavandula angustifolia Miller: English Lavender", *Holistic Nursing*
586 *Practice*, 23(1): 57-64.
- 587 (2) Cavanagh HMA, Wilkinson JM (2002) "Biological activities of Lavender essential oil",
588 *Phytoterapy Research*, 16: 301-308.
- 589 (3) Areias FM, Valentao P, Andrade PB, Moreira MM, Amaral J, Seabra RM (2000)
590 "HPLC/DAD analysis of phenolic compounds from lavender and its application to quality
591 control", *Journal of Liquid Chromatography and Related Technologies*, vol. 23, no. 16, pp.
592 2563-2572.
- 593 (4) Upson TM, Grayer RJ, Greenham JR, Williams CA, Al-Ghamdi F, Chen FH (2000)
594 "Leaf flavonoids as systematic characters in the genera Lavandula and Sabaudia", *Biochemical*
595 *Systematics and Ecology*, vol. 28, no. 10, pp 991-1007.
- 596 (5) Politi M, De Tommasi N, Pescitelli G, Di Bari L, Morelli I, Braca A (2002) "Structure
597 and absolute configuration of new diterpenes from Lavandula multifida", *Journal of Natural*
598 *Products*, vol. 65, no. 11, pp 1742-1745.
- 599 (6) Topcu G, Ayral MN, Aydin A, Goren AC, Chai HB, Pezzuto JM (2001) "Triterpenoids
600 of the roots of Lavandula stoechas ssp. Stoechas", *Pharmazie*, vol. 56, no. 11, pp 892-895.
- 601 (7) Ulubelen A, Goren N, Olcay Y (1988) "Longipinene derivatives from Lavandula
602 stoechas subsp. Stoechas", *Phytochemistry*, vol. 27, no. 12, pp. 3966-3967.
- 603 (8) Shimizu M, Shogawa H, Matsuzawa T, Yonezawa S, Hayashi T, Arisawa M, Suzuki S,
604 Yoshizaki M, Morita N, Ferro E, Basualdo I, Berzanga LH (1990) "Anti-inflammatory
605 constituents of topically applied crude drugs. IV. Constituents and anti-inflammatory effect of
606 Paraguayan crude drug "Alhucema" (Lavandula latifolia Vill.) " *Chemical and pharmaceutical*
607 *bulletin*, vol 38, no. 8, pp. 2283-2284.

- 608 (9) Wilkinson JM , Hipwell M , Ryan T , Cavanagh HMA (2003) “Bioactivity of
609 Backhousia citriodora: Antibacterial and Antifungal Activity”, *J. Agric. Food Chem.*, 51 (1), pp
610 76–81.
- 611 (10) Galesi R, Giardina G, Rossello F (2005) “New data on the Sicilian Flora”, *Informatore*
612 *Botanico Italiano*, 37(2), 1161.
- 613 (11) Ramchoun M, Harnafi H, Alem C, Benlyas M, Elrhaffari L, Amrani S (2009) “Study
614 on antioxidant and hypolipidemic effects of polyphenol-rich extracts from *Thymus vulgaris* and
615 *Lavandula multifida*”, *Pharmacognosy Research*, 1(3), 106.
- 616 (12) Sosa S, Altinier G, Politi M, Braca A, Morelli I, Della Loggia R. (2005) “Extracts and
617 constituents of *Lavandula multifida* with topical anti-inflammatory activity”, *Phytomedicine*,
618 12(4), 271-277.
- 619 (13) Zuzarte M, Vale-Silva L, Gonçalves M J, Cavaleiro C, Vaz S, Canhoto J, Salgueiro L
620 (2012) “Antifungal activity of phenolic-rich *Lavandula multifida* L. essential oil”, *European*
621 *journal of clinical microbiology & infectious diseases*, 31(7), 1359-1366.
- 622 (14) Conti F, Manzi A, Pedrotti F (1997) “Liste Rosse regionali delle Piante d'Italia”. WWF
623 Italia, Società Botanica Italiana, Camerino.
- 624 (15) Renaud ENC, Charles DJ, Simon JE (2001) “Essential oil quantity and composition
625 from 10 cultivars of organically grown lavender and lavandin”. *Journal of Essential Oil*
626 *Research* 13, 269–273.
- 627 (16) Reverter M, Bontemps N, Lecchini D, Sasal P (2014) “Use of plant extracts in fish
628 aquaculture as an alternative to chemotherapy: Current status and future perspectives”.
629 *Aquaculture*, 433: 50-61.

- 630 (17) Hashemi SR, Zulkifli I, Zunita Z, Somchit MN (2008) "The effect of selected
631 sterilization methods on antibacterial activity of aqueous extract of herbal plants". *Journal of*
632 *Biological Sciences* 8: 1072-1076.
- 633 (18) Sakai M (1999) "Current research status of fish immunostimulants". *Aquaculture*;
634 172:63-92.
- 635 (19) Magnadottir B (2010) "Immunological control of fish diseases". *Marine Biotechnology*
636 *(New York)*, 12(4): 361-379.
- 637 (20) Bairwa MK, Jakhar JK, Satyanarayana Y, Reddy AD (2012) "Animal and plant
638 originated immunostimulants used in aquaculture". *Journal of natural products and plant*
639 *resources*, 2(3): 397-400.
- 640 (21) Secombes C J (1994) "Enhancement of fish phagocytic activity". *Fish and Shellfish*
641 *Immunology*. 4: 421-436.
- 642 (22) Muthusamy G, Joardar SN, Samanta I, Isore DP, Roy B and Maiti KT (2013) "β-Glucan
643 from edible mushroom (*pleurotus florida*) enhances mucosal immunity in poultry". *Advances in*
644 *Animal and Veterinary Sciences*. 1 (4): 116 – 119.
- 645 (23) Ringo E, Olsen RE, Vecino JLG, Wadsworth S, Song SK (2012) "Use of
646 immunostimulants and nucleotides in aquaculture: a review". *J Marine Sci Res Development*
647 1:104.
- 648 (24) Harikrishnan R, Balasundaram C, Heo MS (2011) "Impact of plant products on innate
649 and adaptive immune system of cultured finfish and shellfish". *Aquaculture* 317: 1-15.
- 650 (25) Cuesta A, Rodriguez A, Esteban MA, Meseguer J (2005) "In vivo effects of propolis, a
651 honeybee product, on gilthead seabream innate immune responses". *Fish & Shellfish*
652 *Immunology* 18: 71e80.

653 (26) Naga Preethi N, Rajeshwari P (2014) "An overview on immunomodulators".
654 *International Journal of Current Pharmaceutical & Clinical Research* 4 (2): 108-114.

655 (27) Bulfon C, Volpatti D, Galeotti M (2014) "In Vitro Antibacterial Activity of Plant
656 Ethanolic Extracts against Fish Pathogens". *Journal of the world aquaculture society* 45 (5):
657 545-557.

658 (28) Berridge, M.V. and Tan, A.S. (1993) 'Characterization of the Cellular Reduction of 3-
659 (4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium Bromide (MTT): Subcellular Localization,
660 Substrate Dependence, and Involvement of Mitochondrial Electron Transport in MTT
661 Reduction'. *Archives of Biochemistry and Biophysics* 303: 474-482.

662 (29) Denizot F and Lang R (1986) 'Rapid colorimetric assay for cell growth and survival.
663 Modifications to the tetrazolium dye procedure giving improved sensitivity and reliability'. *J*
664 *Immunol Methods* 89(2):271-7.

665 (30) Stevens MG, Kehrli ME, Canning PC (1991) "A colorimetric assay for quantitating
666 bovine neutrophil bactericidal activity". *Veterinary Immunology and Immunopathology*, 28: 45-
667 56.

668 (31) Esteban MA, Mulero V, Munoz J, Meseguer J (1998) "Methodological aspects of
669 assessing phagocytosis of *Vibrio anguillarum* by leucocytes of gilthead seabream (*Sparus aurata*
670 L.) by flow cytometry and electron microscopy". *Cell Tissue Res.* 293: 133–141.

671 (32) Rodriguez A, Esteban MA, Meseguer J (2003) "A mannose- receptor is possibly
672 involved in the phagocytosis of *Saccharomyces cerevisiae* by seabream (*Sparus aurata* L.)
673 leucocytes". *Fish Shellfish Immunol.* 14: 375–388.

674 (33) Bayne CJ, Levy S (1991) "Modulation of the oxidative burst in trout myeloid cells by
675 adrenocorticotrophic hormone and catecholamines: mechanisms of action". *J. Leukoc. Biol.* 50:
676 554–560.

677 (34) Quade MJ, Roth JA (1997) "A rapid, direct assay to measure degranulation of bovine
678 neutrophil primary granules". *Vet. Immunol. Immunopathol.* 58: 239–248.

679 Nassar M, Zerizer S, Kabouche Z, Kabouche A, Bechkri S (2015) "Antioxidant and the
680 immunomodulatory activities exhibited by three plants from Lamiaceae family". *International*
681 *Journal of Pharmacy and Pharmaceutical Sciences.* 7 (9): 331-334.

682 Logambal SM, Venkatalakshmi S, Dinakaran Michael R (2000) "Immunostimulatory effect of
683 leaf extracts of *Ocimum sanctum* Linn. in *Oreochromis mossambicus* (Peters)". *HydroBiologia.*
684 430: 113-120.

685 Adel M, Amiri AA, Zorriehzahra J, Nematolahi A, Esteban MA (2015) "Effects of dietary
686 peppermint (*Mentha piperita*) on growth performance, chemical body composition and
687 hematological and immune parameters of fry Caspian white fish (*Rutilus frisii kutum*). *Fish and*
688 *Shellfish Immunology.* 45: 841-847.

689 Fent K (2001) "Fish cell lines as versatile tools in ecotoxicology: assessment of cytotoxicity,
690 cytochrome P4501A induction potential and estrogenic activity of chemicals and environmental
691 samples". *Toxicology in Vitro.* 15: 477-488.

692 Baser KH (2008) "Biological and pharmacological activities of carvacrol and carvacrol bearing
693 essential oils". *Current Pharmaceutical Design.* 14(29): 3106-3119.

694 Khadir A, Bendahou M, Benbelaid F, Bellahcene C, Zenati F, Abdelouahid D, Muselli A, Costa
695 J (2014) "Chemical composition and anti-MRSA activity of essential oils and ethanol extract of
696 *Lavandula multifida* L. from Algeria". *Journal of Essential oil Bearing Plants.* 19(3): 712-718.

697 Fazio A, Barreca D, Bonaccorsi PM, Panuccio MR (2014) "Antioxidant properties and
698 polyphenolic composition of leaves of calabrian *Lavandula multifida*". *Journal of Biological*
699 *Research.* Bollettino della Societa' Italiana di Biologia Sperimentale (SIBS) 87(1): 4.

700 Alcaraz MJ, Hoult JRS (1985) "Action of flavonoids and the novel-antiinflammatory flavone,
701 hypolaetin-8-glucoside, on prostaglandin biosynthesis and inactivation". 34 (14): 2477-82.

702 Lin Y, Shi R, Wang X, Shen HM (2008) "Luteolin, a flavonoid with potentials for cancer
703 prevention and therapy". *Current cancer drug targets*. 8(7):634-646.

704 Saeed M, Kadioglu O, Khalid H, Sugimoto Y, Efferth T (2015) "Activity of the dietary
705 flavonoid, apigenin, against multidrug-resistant tumor cells as determined by
706 pharmacogenomics and molecular docking". *The Journal of Nutritional Biochemistry*. 26(1):44-
707 56.

708 Choi DY, Lee JY, Kim MR, Woo ER, Kim YG, Kang KW (2005) "Chrysoeriol potently inhibits
709 the induction of nitric oxide synthase by blocking AP-1 activation.*Journal of biomedical*
710 *science*. 12(6): 949-959.

711 Madhuri S, Mandloi AH, Pandey G, Sanhi YP (2012) "Antimicrobial activity of some medicinal
712 plants against fish pathogens". *International Research Journal of Pharmacy*. 3(4): 28-30.

713 Soltani M, Sheikhzadeh N, Ebrahimzadeh-Mousavi HA, Zargar A (2010) "Effects of Zataria
714 multiflora essential oil on innate immune responses of Common Carp (*Cyprinus carpio*)".
715 *Journal of Fisheries and Aquatic Science*. 5(3): 191-199.

716 Alderman DJ, Hasting TS (1998) "Antibiotic use in aquaculture: development of antibiotic
717 resistance-potential for consumer health risks". *International Journal of Food and Science*
718 *Technology*. 33, 139–155.

719 Cabello FC (2006) "Heavy use of prophylactic antibiotics in aquaculture: a growing problem for
720 human and animal health and for the environment". *Environmental Microbiology*. 8, 1137–1144.

721 Abutbul S, Golan-Goldhirsh A, Brazani O, Zilberg D (2004) "Use of *Rosmarinus officinalis* as a
722 treatment against *Streptococcus iniae* in tilapia (*Oreochromis sp.*)" *Aquaculture*. 238, 97–105.

723 Smith P, Hiney MP, Samuelesen OB (1994) “Bacterial resistance to antimicrobial agent used in
 724 fish farming: a critical evaluation of method and meaning”. *Annual Review of Fish Diseases* 4,
 725 273–313.

726 MacMillan JR (2001) “Aquaculture and antibiotic resistance: a negligible public health risk?”
 727 *World Aquaculture* 32, 49–68.

728 Rijkers GT, Teunissen AG, Van Oosterom R, Van Muiswinkel WB (1980) “The immune
 729 system of cyprinid fish: the immunosuppressive effect of the antibiotic oxytetracycline in carp
 730 (*Cyprinus carpio* L.)” *Aquaculture* 19, 177–189.

731 Harikrishnan R, Balasundaram C, Dharaneedharan S, Moon YG, Kim MC, Kim JS, Heo MS
 732 (2009) “Effect of plant active compounds on immune response and disease resistance in
 733 *Cirrhina mrigala* infected with fungal fish pathogen, *Aphanomyces invadans*”. *Aquaculture*
 734 *Research* 40, 1170–1181.

735 FAO (2003) “Paraquat. Pesticide Residues in Food, Report 2003: Evaluations”. Geneva,
 736 Switzerland. <http://www.fao.org/docrep/005/Y4544E/y4544e00.htm>.

737 Valladão GM, Gallani SU, Pilarski F. 2015 Phytotherapy as an alternative for treating fish
 738 disease. *J Vet Pharmacol Ther.* Oct;38(5):417-28.

739 Tafalla C, Aranguren R, Secombes CJ, Figueras A, Novoa B (2004) Cloning and analysis of
 740 expression of a gilthead seabream (*Sparus aurata*) Mx cDNA. *Fish Shellfish Immunol* 16, 11–24

741 Pelegrín EP, Chaves-Pozo E, Mulero V, Meseguer J (2004) Production and mechanism of
 742 secretion of interleukin-1beta from the marine fish gilthead seabream. *Dev Comp Immunol* 28,
 743 229–237.

744 Diop M, Watier D, Masson PY, Diouf A, Amara R, Grard T, Lencel P. Assessment of freshness
 745 and freeze-thawing of sea bream fillets (*Sparus aurata*) by a cytosolic enzyme: Lactate
 746 dehydrogenase. *Food Chem.* 2016 Nov 1;210:428-34.

747 Cano I, Collet B, Pereira C, Paley R, van Aerle R, Stone D, Taylor NG. In vivo virulence of
 748 viral haemorrhagic septicaemia virus (VHSV) in rainbow trout *Oncorhynchus mykiss* correlates
 749 inversely with in vitro Mx gene expression. *Vet Microbiol.* 2016 May 1;187:31-40.

750 Morcillo P, Esteban MÁ, Cuesta A. Heavy metals produce toxicity, oxidative stress and
 751 apoptosis in the marine teleost fish SAF-1 cell line. *Chemosphere.* 2016 Feb;144:225-33.

752 Trapani A, Mandracchia D, Di Franco C, Cordero H, Morcillo P, Comparelli R, Cuesta A,
 753 Esteban MA. In vitro characterization of 6-Coumarin loaded solid lipid nanoparticles and their
 754 uptake by immunocompetent fish cells. *Colloids Surf B Biointerfaces.* 2015 Mar 1;127:79-88.

755 Picchietti S, Bernini C, Belardinelli MC, Ovidi E, Taddei AR, Guerra L, Abelli L, Fausto AM.
 756 Immune modulatory effects of *Aloe arborescens* extract on the piscine SAF-1 cell line. *Fish*
 757 *Shellfish Immunol.* 2013 May;34(5):1335-44.

758 El Aamri F, Real F, Acosta F, Acosta B, Valdivia J, Ramos-Vivas J, Padilla D. In vitro study of
 759 adherence, invasion, and persistence of *Streptococcus iniae* in fibroblastic-like fish cell line
 760 SAF-1. *J Aquat Anim Health.* 2012 Sep;24(3):165-70.

761 Acosta F, Vivas J, Padilla D, Vega J, Bravo J, Grasso V, Real F. Invasion and survival of
 762 *Photobacterium damsela* subsp. *piscicida* in non-phagocytic cells of gilthead sea bream, *Sparus*
 763 *aurata* L. *J Fish Dis.* 2009 Jun;32(6):535-41.

764 García-Rosado E, Alonso MC, Béjar J, Manchado M, Cano I, Borrego JJ. Expression analysis
 765 of Mx protein and evaluation of its antiviral activity against sole aquabirnavirus in SAF-1 and
 766 TV-1 cell lines. *Vet Immunol Immunopathol.* 2008 Jan 15;121(1-2):123-9.

767 Cano I, Ferro P, Alonso MC, Bergmann SM, Römer-Oberdörfer A, Garcia-Rosado E, Castro D,
 768 Borrego JJ. Development of molecular techniques for detection of lymphocystis disease virus in
 769 different marine fish species. *J Appl Microbiol.* 2007 Jan;102(1):32-40.

770 Bandín I, Oliveira JG, Borrego JJ, Dopazo CP, Barja JL. Susceptibility of the fish cell line SAF-
 771 1 to betanodavirus. *J Fish Dis.* 2006 Oct;29(10):633-6.

772 Alonso MC, Cano I, Garcia-Rosado E, Castro D, Lamas J, Barja JL, Borrego JJ, Bergmann
 773 SM. Isolation of lymphocystis disease virus from sole, *Solea senegalensis* Kaup, and blackspot
 774 sea bream, *Pagellus bogaraveo* (Brunnich). *J Fish Dis.* 2005 Apr;28(4):221-8.

775 Pelegrín P, Chaves-Pozo E, Mulero V, Meseguer J. Production and mechanism of secretion of
 776 interleukin-1beta from the marine fish gilthead seabream. *Dev Comp Immunol.* 2004
 777 Mar;28(3):229-37.

778 Tafalla C, Aranguren R, Secombes CJ, Figueras A, Novoa B. Cloning and analysis of
779 expression of a gilthead sea bream (*Sparus aurata*) Mx cDNA. *Fish Shellfish Immunol.* 2004
780 Jan;16(1):11-24.

781 Ray A, Yu GY, Ray BK. Cytokine-responsive induction of SAF-1 activity is mediated by a
782 mitogen-activated protein kinase signaling pathway. *Mol Cell Biol.* 2002 Feb;22(4):1027-35.

783 Ray A, Kumar D, Ray BK. Promoter-binding activity of inflammation-responsive transcription
784 factor SAF is regulated by cyclic AMP signaling pathway. *DNA Cell Biol.* 2002 Jan;21(1):31-
785 40.

786 Perez-Prieto SI, Rodriguez-Saint-Jean S, Garcia-Rosado E, Castro D, Alvarez MC, Borrego JJ.
787 Virus susceptibility of the fish cell line SAF-1 derived from gilt-head seabream. *Dis Aquat*
788 *Organ.* 1999 Jan 29;35(2):149-53.

789

790

791