

*This is the peer reviewed version of the following article:*

**Synowiec, A., Mozdzeń, K., Krajewska, A., Landi, M., & Araniti, F. (2019). *Carum carvi* L. essential oil: A promising candidate for botanical herbicide against *Echinochloa crus-galli* (L.) P. Beauv. in maize cultivation. *Industrial Crops and Products*, 140, 111652.**

*which has been published in final doi:* <https://doi.org/10.1016/j.indcrop.2019.111652>

<https://www.sciencedirect.com/science/article/pii/S0926669019306624>

*The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website*

1 ***Carum carvi* essential oil: a promising candidate for botanical**  
2 **herbicide against *Echinochloa crus-galli* in maize cultivation**

3  
4 Synowiec A.<sup>a,\*</sup>, Możdżeń K.<sup>b</sup>, Krajewska A.<sup>c</sup>, Marco Landi<sup>d</sup>, Fabrizio Araniti<sup>e,\*</sup>

5  
6 <sup>a</sup>Department of Agrotechnology and Agricultural Ecology, University of Agriculture, Poland

7 <sup>b</sup>Department of Plant Physiology, Pedagogical University of Krakow, Poland

8 <sup>c</sup>Department of Biotechnology and Food Science, Lodz University of Technology, Poland

9 <sup>d</sup>Department of Agriculture, Food and Environment, University of Pisa, Pisa, Italy

10 <sup>e</sup>Dipartimento AGRARIA, Università Mediterranea di Reggio Calabria, – Località Feo di Vito,  
11 SNC I-89124 Reggio Calabria RC, Italy

12

13 **Corresponding authors:**

14 Agnieszka Synowiec: a.synowiec@urk.edu.pl

15 Fabrizio Araniti: fabrizio.araniti@unirc.it

16

17 **Highlights (3 to 5 bullets, max 85 characters per point including spaces)**

18

19

20 **Abstract**

21 In this study we tested the possibility that foliar-applied caraway or peppermint essential oils  
22 (EOs) can selectively inhibit the growth of *Echinochloa crus-galli* (a typical maize weed) but  
23 not that of maize plants, attempting to develop an eco-friendly botanical herbicide.

24 We tested the phytotoxic potential of oil-in-water emulsions of each EO with addition of  
25 commercial adjuvant mainly composed of fatty acids methyl esters, studying their effect on

26 visible plants injuries, biomass accumulation, chlorophyll *a* fluorescence and changes to  
27 biochemical patterns of both the main crop (maize) and the weed (*E. crus-galli*) via an  
28 untargeted metabolomic approach. We found that oil-in-water emulsion containing 2.5% of  
29 adjuvant and of caraway EO did not affect significantly the growth of maize plants, did not  
30 induce foliar symptoms and did not alter the status of the photosynthetic apparatus, as revealed  
31 by chlorophyll *a* fluorescence. On the contrary, this emulsion exerted significantly negative  
32 effects against *E. crus-galli* growth, inducing foliar injuries and reducing the photosynthetic  
33 efficiency of photosystem II. We also found that the studied emulsions caused a series of  
34 biochemical changes in the plant tissues, with caraway emulsion being more phytotoxic, as  
35 compared to the peppermint EO-emulsion. We conclude that oil-in-water emulsion containing  
36 2.5% of caraway EO could be used in future as a foliar-applied botanical herbicide against *E.*  
37 *crus-galli* in maize cultivation.

38

39 **Keywords:** Bio-herbicide; biochemical process; chlorophyll *a* fluorescence; leaf injury;  
40 metabolomics; phytotoxicity

## 41 **1. Introduction**

42 Numerous studies shown that essential oils (EOs), especially those for which the main  
43 compounds are monoterpene alcohols or oxygenated monoterpenes, are promising substances  
44 for production of botanical herbicides, since they can cause significant inhibition of weed  
45 germination and growth (Benvenuti et al., 2017; Synowiec et al., 2017, Rolli 2014, Vokou et  
46 al., 2003), as well as they provoke, severe leaf burns in foliar applications (Bainard et al., 2006,  
47 Stokłosa et al., 2012). Considering the chemical composition of their EOs, peppermint (*Mentha*  
48 *x piperita* L.) and caraway (*Carum carvi* L.) can be promising candidates for the production of  
49 bioherbicides in a temperate European climate (Synowiec et al., 2017). Both species are widely  
50 cultivated in Europe (Oroian et al., 2017; Seidler-Łożykowska and Bocianowski 2012) and are  
51 characterized by high EO yields. In particular, in plants cultivated in Poland, EOs extraction  
52 from peppermint leaves and caraway seeds could give a yield around 2.3% and 3.4-4.8%,  
53 respectively (Pisulewska et al., 2010; Seidler-Łożykowska et al., 2013). Both EOs are rich in  
54 oxygenated monoterpenes ( > 80 % for peppermint and > 60 % for caraway EO) (Seidler-  
55 Łożykowska et al., 2013; Fejér et al., 2017). The main chemical compounds of these EOs are  
56 menthol and menthone in peppermint (Guidi and Landi, 2014), and carvone and limonene in  
57 caraway EOs (Chemat et al., 2017).

58 However, physical and chemical properties of EOs, such as high volatility or poor water  
59 solubility make difficult a wider use of them as natural herbicides. These disadvantages can be  
60 overcome by creating appropriate emulsions. As shown by Hazrati et al. (2017), garden savory  
61 (*Satureja hortensis*) EOs applied as oil-in-water (o/w) nanoemulsion, with 2% (v/v) Tween 80,  
62 displayed adequate physical properties and posed a strong phytotoxic potential on germination  
63 and early growth of *Amaranthus retroflexus* and *Chenopodium album*. In turn, Synowiec et al.  
64 (2017) showed satisfactory effectiveness of o/w emulsion of peppermint EO (2.5%) applied  
65 with the addition of oilseed rape fatty acid methyl esters (1.5 L ha<sup>-1</sup>) against *E. crus-galli*.

66 Foliar-applied EOs display a contact action and induce visible injuries caused as early as  
67 few hours following their application (Hazrati et al., 2017; Synowiec et al., 2015). In general,  
68 foliar-applied EOs mixture leads to a general impairment of plant metabolism due to multi-  
69 spectrum targets (Synowiec et al., 2015). Conversely, application of a single or a few  
70 compounds isolated from EO may act selectively via inhibition of a specific metabolic pathway  
71 (Araniti et al., 2017a; Araniti et al., 2016; Graña et al., 2013), but this approach is often more  
72 important for obtaining a total herbicide rather than a selective herbicide. Many experiments  
73 showed that one of the main effects of EOs is the inhibition of photosynthesis, resulting from a  
74 decrease in the chlorophyll content (Hazrati et al., 2017) and alterations of the light phase of  
75 photosynthesis (Synowiec et al., 2015). In some cases, EOs lead to the production of  
76 uncontrolled level of reactive oxygen species, thereby promoting oxidative stress and oxidative  
77 burst (Ahuja et al., 2015) as well as loss of the efficiency of cellular respiration (Kaur et al.,  
78 2010). Recently, Araniti et al. (2018), through a physiological and metabolomic approach,  
79 described in detail the physiological response of *Arabidopsis thaliana* seedlings to the EO of  
80 oregano. The authors observed a reduction of plant growth and leaf chlorosis of *A. thaliana*  
81 seedlings as a result of series of metabolic alterations, including principally the inability to  
82 incorporate assimilated nitrogen into amino acids, especially the nitrogen devoted to the  
83 biosynthesis of one of the first precursors of other amino acids, namely glutamine.

84 The metabolomic approach allows to analyze simultaneously hundreds of metabolites in a  
85 given biological sample (Nicholson and Lindon, 2008), yielding a comprehensive picture of  
86 changes in the metabolism of plants under different types of stresses (Mosa et al., 2017, Ghatak  
87 et al., 2018). Therefore, untargeted metabolomics could consent to characterize the phytotoxic  
88 effects of foliar application of EOs emulsion on key metabolic pathways, in order to understand  
89 the main biochemical/physiological processes altered in the plant. For this reason, this research  
90 aimed at: i) assessing the phytotoxic potential of foliar-sprayed peppermint or caraway EOs,

91 applied as o/w emulsions with addition of a commercial adjuvant in maize (*Zea mays* L.) and  
92 barnyardgrass [*Echinochloa crus-galli* (L.) P.Beauv.], and ii) employ the imaging of  
93 chlorophyll *a* fluorescence and an untargeted metabolomic approach to dissect plant responses  
94 to foliar application of EOs emulsions.

95

## 96 **2. Materials and Methods**

### 97 *2.1. Characteristic of essential oils and adjuvant*

98 The EOs isolated from caraway (*Carum carvi* L.) seeds was purchased from the Avicenna-  
99 Oil company (Wrocław, Poland), whereas the essential oil of peppermint (*Mentha ×piperita*  
100 L.) was steam-distilled for 2 h in the laboratory conditions using Deryng-type apparatus (Baj et  
101 al., 2015) from the air-dry mass of herbs collected from the production fields in Michałowice,  
102 Poland (50°37'45''N, 20°48'03''E), in summer 2015.

103 Commercial multifunctional adjuvant ATPOLAN BIO 80 EC (Producer: AGROMIX  
104 Niepołomice, Poland) was chosen as adjuvant and emulsifier of EOs. This adjuvant is mainly  
105 composed of fatty acid methyl esters of oilseed rape oil (80%), surfactants and a pH buffer  
106 (according to the product label provided by the producer). It was chosen as in previous  
107 experiments this adjuvant displayed good herbicidal potential as an emulsifier of peppermint or  
108 caraway EOs (Synowiec and Drozdek, 2016).

109

### 110 *2.2. Chemical analysis of essential oils*

111 The chemical composition of the EOs was analyzed by gas chromatography coupled with  
112 mass spectrometry (GC-FID-MS). After dilution in diethyl ether (10 µL in 1 mL), the EOs were  
113 analyzed using a Trace GC Ultra gas chromatograph coupled with DSQ II mass spectrometer  
114 (Thermo Electron Corporation). The operating conditions were as follows: non-polar capillary  
115 column Rtx-1ms (60 m x 0.25 mm, 0.25 mm film thickness), programmed temperature: 50 (3

116 min)-300°C, 4°C/min, injector (SSL) temperature 280°C, flame ionization detector temperature  
117 300°C, transfer line temperature 250°C, carrier gas - helium, flow with constant pressure 200  
118 kPa, split ratio 1:20. The mass spectrometer parameters: ion source temperature 200°C,  
119 ionization energy 70 eV (EI), scan mode: full scan, mass range 33-420. The percentages of  
120 constituents were computed from the GC peak area without using a correction factor.

121 Identification of EO components was based on a comparison of their mass spectra and linear  
122 retention indices (RI, non-polar column), determined with reference to a series of n-alkanes C8-  
123 C26, by comparing with those reported by Adams (2007) as well as in computer libraries: NIST  
124 2011, and MassFinder 4.1. Percentages were obtained from the FID response without the use  
125 of correction factors.

126

### 127 *2.3. Preparation of o/w emulsions*

128 The emulsion (250 mL) based on 2.5% or 5% of each of EO was prepared at room  
129 temperature. For the emulsion, 6.25 g or 12.5 g of selected essential oil, respectively were  
130 weighted into a vial. Then, 6.25 g (for 2.5% solution) or 12.5 g (for 5% solution) of ATPOLAN  
131 BIO 80 EC was added. This mixture was stirred vigorously on magnetic stirrer (300 rpm) using  
132 3 cm stir bar. Afterwards, while constant mixing 237.4 mL or 225 mL of distilled water was  
133 added in small portions. Then the stirring was increased to 500 rpm for 5 min. Prepared  
134 emulsion was homogenized using handheld homogenizer (Ingenieurbüro CAT M. Zipperer  
135 GmbH, Unidrive D, rotation speed 5000 rpm). The emulsions were stored at room temperature  
136 until use.

137

### 138 *2.4. Pot experiment*

139 A pot experiment was settled up in the foil tunnel (16 m long, 6 m wide and 3 m high) in  
140 Krakow-Mydlniki, south of Poland (N 50° 08'54'', E 19° 85'21''), with daily temperature

141 monitoring. The experiment was established in the period of 6<sup>th</sup> April – 8<sup>th</sup> June 2017. There  
142 were ten replications (plants) per each species and emulsion treatment.

143 Seven seedling palettes (0.154 m<sup>2</sup> with 24 pots in 4 rows with a single pot size: 46 x 46 x 70  
144 mm per each palette), were filled up with a sieved layer (0-15 cm) of a sandy brown soil (pH  
145 6.3; P<sub>2</sub>O<sub>5</sub> 18.2; K<sub>2</sub>O 7.5; MgO 6.9 [mg 100 g<sup>-1</sup> of soil]). Two seeds of maize (cv. ‘Wilga’) or a  
146 few seeds of barnyard grass (*Echinochloa crus-galli* (L.) Beauv) were sowed into the 10 of pots  
147 per palette per species and after emergence the number of plants was thinned to one per pot.  
148 The four middle pots per palette were left empty to avoid shading between plants of maize and  
149 barnyard grass and to optimize the spraying process. Then the palettes were positioned  
150 alternately. During growth the plants were watered according to their needs.

151 When maize reached the growth stage of 4-6 leaves and barnyard grass the stage of 3-4 leaves,  
152 the plants were hand-sprayed with one of the following emulsions:

- 153 i) Water only (control; W);
- 154 ii) Water + 2.5 % adjuvant (Control; WA2.5);
- 155 iii) Water + 5.0 % adjuvant (Control; WA5.0);
- 156 iv) Water + 2.5 % caraway EOs + 2.5 % of adjuvant (WAC2.5);
- 157 v) Water + 5.0 % caraway EOs + 5.0 % of adjuvant (WAC5.0);
- 158 vi) Water + 2.5 % peppermint EOs + 2.5 % adjuvant (WAP2.5);
- 159 vii) Water + 5.0 % peppermint EOs + 5.0 % of adjuvant (WAP5.0).

160 Each palette was sprayed with 10 cm<sup>3</sup> of one of the emulsions (i-vii), using a 1 L volume hand  
161 pressure sprayer Kwazar Venus Super 360 PRO+ (Producer: Kwazar Corporation Sp. z o.o.,  
162 Poland). The calculated amounts of the EOs in the spraying solutions were equal to 1.5 g m<sup>-1</sup>  
163 (solutions iv and vi) and 3.0 g m<sup>-1</sup> (solutions v and vii).

164 Seven days after spraying the plants were visually assessed (by one person) for the percentage  
165 of aboveground injuries (0-100%) caused by the foliar-application of emulsions (i-vii). Next,



166 the plants were cut at the ground level. For each plant, the aboveground parts were placed in  
167 envelopes and dried in the temperature of 50 °C in a lab oven for 3 days. After that, their dry  
168 mass was recorded.

169

## 170 2.5. Chlorophyll a fluorescence imaging

171 The analyses were carried out on 5 plants (replications) of each species and for selected  
172 treatments:

- 173 i) Water (Control; W). The two other control treatments, namely: water plus 2.5% of  
174 adjuvant and water plus 5% of adjuvant, showed results similar to water, so for this  
175 analysis only water as a control is presented;
- 176 ii) Water + 2.5 % caraway EOs + 2.5 % of adjuvant (WAC2.5);
- 177 iii) Water + 5.0 % caraway EOs + 5.0 % of adjuvant (WAC5.0);
- 178 iv) Water + 2.5 % peppermint EOs + 2.5 % adjuvant (WAP2.5);
- 179 v) Water + 5.0 % peppermint EOs + 5.0 % of adjuvant (WAP5.0).

180 Plants of maize and barnyard grass were grown and sprayed with the emulsions similarly as in  
181 the pot experiment. A second leaf was cut 48 hours after spraying and placed flat on filter paper  
182 moistened with distilled water, and immediately placed in a lightproof measurement chamber  
183 FluorCam FC 800C (Photon Systems Instruments, Czech Republic) for 20 min of dark. The  
184 measurement was taken right after a pulse of saturation actinic light ( $4,000 \mu\text{mol m}^{-2}\text{s}^{-1}$  PAR,  
185 800 ms), according to Lichtenthaler et al. (2005). The following parameters were analyzed: 1)  
186  $F_v/F_m$  (*aka* QYmax) – maximum quantum yield of photosystem II photochemistry, 2) NPQ-  
187 non-photochemical quenching and 3) Rfd-fluorescence decrease ratio (Kalaji et al., 2014). All  
188 of the fluorescence parameters were graphically imaged using FluorCam 7, ver 1.0.20.4  
189 software ([www.psi.cz/downloads/](http://www.psi.cz/downloads/)). The color scale presents conventional values of the studied  
190 parameters of leaves subjected to treatments with the emulsions. The numeric comparisons

191 between the treatments were performed based on a calculation of a “mean gray value”, which  
192 is the sum of the gray values of all the pixels in the selection, by converting each RGB pixel to  
193 grayscale in ImageJ software ver. 1.52a (<http://imagej.nih.gov/ij>).

194

## 195 *2.6. The identification and quantification of primary metabolites in the aboveground parts of* 196 *plants*

197 The selected three plants (replications) sprayed with the higher doses of emulsions for both  
198 EOs were selected for these analyses, as they displayed significant increases in injuries as well  
199 as significant reductions in plants’ biomass, following their foliar applications.

200 i) Water + 5.0 % adjuvant (control; WA5.0);

201 ii) Water + 5.0 % caraway EOs + 5.0 % adjuvant (WAC5.0);

202 iii) Water + 5.0 % peppermint essential oil + 5.0 % adjuvant (WAP5.0).

203 Fourty eight hours after spraying the aboveground parts of plants for each species, samples  
204 sprayed with the emulsions i), ii) or iii), were collected and freeze-dried. Each plant was frozen  
205 separately in liquid nitrogen, homogenized using a laboratory mortar and immediately  
206 lyophilized using the Freeze Dry System (Freezone 4.5, Labconco, USA).

207 The metabolome extraction and derivatization, as well as metabolite identification and relative  
208 quantification of maize and *E. crus-galli* plants treated with caraway and peppermint EOs, were  
209 carried out as previously described by Araniti et al. (2017c). Derivatized samples were injected  
210 into a gas chromatograph apparatus (Thermo Fisher G-Trace 1310), equipped with a capillary  
211 column TG-5MS (30 m×0.25 mm×0.25 µm), coupled to a single quadrupole mass spectrometer  
212 (ISQ LT). Helium at high purity (6.0) was used as a carrier.

213 Injector and source were settled at the temperature of 250°C and 260°C, respectively. Samples  
214 (1 µL) were injected in splitless mode with a helium flow of 1 mL min<sup>-1</sup> and chemical separation  
215 was achieved using the following programmed temperature: isothermal 5 min at 70 °C, from

216 70° to 330°C with a ramp of 5°C min<sup>-1</sup>, isothermal at 330°C for 5 min. Mass spectra were  
217 recorded in electronic impact (EI) mode, scanning at 45–500 amu.

218 The identification of the metabolites was carried out comparing the unknown mass spectra with  
219 reference spectra of several commercial libraries (NIST 2005, Wiley 7.0, Fiehn library etc.).

220 Metabolites relative quantification was based on a pre-added internal standard (adonitol at 0.02  
221 mg mL<sup>-1</sup>), which was added during the extraction process.

222

### 223 *2.7. Experimental design and statistical analysis*

224 The experiments were carried out in a completely randomized design with different number  
225 of replications depending on the parameter evaluated. In particular, 10 replications for leaf  
226 injuries and dry biomass, 5 replications for chlorophyll *a* fluorescence and 3 replications for  
227 metabolomic experiments were used.

228 Estimation of plant injuries, biomass production and chlorophyll fluorescence parameters were  
229 analyzed with R software (R Core Team, 2014), using ‘dplyr’ package. The percentage values  
230 were Bliss transformed prior analyses. All data were tested for their homogeneity of variance  
231 (Levene test) and the normality of distribution (Kolmogorov-Smirnoff test). Data were then  
232 analyzed through one-way ANOVA using the Tukey’s HSD test as post-hoc ( $P \leq 0.05$ )  
233 (‘multcomp’ package).

234 A completely-randomized sampling, was applied for metabolomic analyses, for which  
235 three independent replicates were analyzed. Metabolite concentrations were checked for  
236 integrity and missing values were replaced by a small positive value (the half of the minimum  
237 positive number detected in the data). Data were successively normalized by a reference sample  
238 (adonitol), transformed through “Log normalization” and scaled through Pareto-Scaling. Data  
239 were then classified through Principal Component Analysis (PCA) and metabolite variations

240 were presented as heatmap. Significant differences among the treatments were highlighted  
241 through ANOVA using LSD test as post-hoc ( $P \leq 0.05$ ).

242 The analysis of the pathways perturbed by the treatments was carried out using MetPA, a  
243 web-based tool that combines the results from pathway enrichment analysis with the pathway  
244 topology analysis. Pathway analysis was carried out using the pathway library built on the  
245 metabolome of *Oryza sativa japonica* since the two plants studied (*Z. mays* and *E. crus-galli*)  
246 are monocots. All the metabolomic analysis were carried out using the software Metaboanalyst  
247 3.0 (Xia et al., 2015).

248

### 249 **3. Results and discussion**

#### 250 *3.1. The chemical composition of essential oils*

251 In Table 1 the chemical composition of caraway and peppermint EOs is reported. In  
252 caraway EO the 99.7% of the total ion chromatogram was identified. In particular, the main  
253 compounds were limonene (32.5%) and carvone (66.4%), whereas in peppermint EO the main  
254 components were represented by menthol (42.7%) and menthone (25.5%) (Table 1). The  
255 process of emulsification did not change the chemical composition of EOs (data not shown).  
256 These results are in agreement with Synowiec et al. (2017) who observed a similar chemical  
257 profile of EOs isolated from both species.

258 The phytotoxicity of the major molecules identified is largely known in literature. Menthone  
259 and carvone strongly affected germination and growth of several crops and weeds, including  
260 monocotyledonous *Triticum aestivum* and *Zea mays*, *Lolium multiflorum* and *Digitaria*  
261 *sanguinalis* and also dicotyledonous *Lactuca sativa* (Vaughn et al., 1993; Sunohara et al.,  
262 2015). Among monoterpenes, limonene is one of the most phytotoxic. In fact, recent studies  
263 reported that this molecule strongly affected carrot and cabbage growth and development  
264 causing leaf injuries and affecting the photosynthetic apparatus (Ibrahim et al., 2004). Similarly,

265 Vaid et al. (2011) observed an inhibition of germination, root growth, pigment content and  
266 respiration on *Amaranthus viridis*. Finally, Schults et al. (2007) observed that menthol was  
267 affecting *Arabidopsis* growth, dewaxed the leaf cuticular layer and altered stomatal anatomy  
268 and function. Moreover, it has been demonstrated that also minor compounds in the EO blend  
269 could act synergistically improving the phytotoxicity of the major chemicals (Araniti et al.,  
270 2013).

271

### 272 3.2. Effect of the emulsions on maize and *E. crus-galli* plant injuries and biomass

273 Thermal conditions in the foil tunnel during the course of experiment, both before and after  
274 spraying the plants with emulsions (April-June) were similar and in a range of 19-23 °C.

275 Spraying maize with emulsions containing only water and adjuvant did not cause any injuries  
276 of maize leaves (Fig. 1A). On the other hand, addition of EOs to the emulsions caused injuries  
277 of maize leaves (in the form of necroses) by 8-40 %, as compared to control (W), with  
278 significant injuries caused by the emulsions containing 5.0 % of caraway or peppermint oil (Fig.  
279 1A). The leaf-spraying of emulsions containing EOs caused a significant reduction of maize  
280 dry mass, especially with treatments containing 5% of caraway (57% reduction) or peppermint  
281 EO (41% reduction) (Fig. 1B). Should be noted that treatment WAC2.5 did not affect both leaf  
282 integrity and plant biomass.

283 *E. crus-galli*, plants sprayed with emulsions, which contained only water and adjuvant  
284 (both WA2.5 and WA5.0), did not cause any visible injury (Fig. 2A). Addition of caraway EO  
285 to the emulsion caused a significant increase of leaf injuries, as compared to W and WA2.5  
286 controls, ranging from 20 to 42% for WAC2.5 and WAC5.0, respectively. Both WAC2.5 and  
287 WAC5.0 induced a decrease of plant biomass with respect to W controls and WA2.5 (Fig. 2b).  
288 Noteworthy, the use of the adjuvant alone at the higher dose, WA5.0, promoted a reduction of  
289 *E. crus-gallis* biomass with a similar extent to those observed with WAC2.5 and WAC5.0,

290 suggesting a negative effect of the adjuvant alone. However, the adjuvant alone did not cause  
291 visible injuries at any concentration(Fig. 2A).. On the contrary, WAP2.5 and WAP5.0  
292 treatments similarly affected leaf integrity causing leaf injuries on the 30% of the leaf surface  
293 (no statistical differences were observed between the two treatments) (Fig. 2). Among  
294 treatments, the emulsion containing 5% of caraway oil was the most harmful for *E. crus-galli*  
295 leaves (Fig. 2A). Concerning plant biomass, the most significant decrease was observed on  
296 plant sprayed with both emulsions containing peppermint oil, which caused a 50% reduction in  
297 dry biomass compared to control (W) (Fig. 2B).

298 Therefore, biometric analyses revealed that water emulsions of EOs and a commercial  
299 adjuvant composed of FAME of oilseed rape caused a significant reduction of biomass of maize  
300 and *E. crus-galli*, with emulsions containing EO of peppermint being more toxic than those of  
301 caraway. Notably, a significant reduction of biomass was observed in maize for the dose of EO  
302 in emulsions as high as 5% by using caraway oil, whereas for *E. crus-galli* the negative effects  
303 occurred even when the lower dose was applied (WAC2.5). For the sack of the truth, reduction  
304 of plant biomass could be partially attributable to the effect of the adjuvant alone (see WA5.0  
305 biomass reduction) but that WAC5.0 did not cause any visible injury and plant damage (whilst  
306 WAC2.5 and WAC 5.0 did) is a valuable result of the effectiveness of caraway EO. Our dataset  
307 cannot explain the physiological reasons behind the side effect of adjuvant on *E. crus-gallis*  
308 biomass reduction and further research is needed to clarify this point. In any case, visible  
309 injuries were only detected in WAC2.5 plants of *E. crus-galli* and not in maize plants, which is  
310 a promising result in the attempt to develop a selective botanical herbicide. One should also  
311 consider that the selective 20% damage over *E. crus-galli* leaves (which is seems not a  
312 negligible result for an ecofriendly botanical herbicide bioassayed at such low concentrations)  
313 can strongly reduce the competition between the crop and the pest in the field in an early  
314 developmental stage, thus allowing maize to be more competitive and to grow faster. After the

315 first stage, *E. crus-galli* would suffer for the fast-growing maize developing, this resulting in a  
316 further reduction of the possibility to compete with maize plants. In addition, our experiment  
317 describes the use of a single treatment with WAC2.5 EO, but repetitive treatments could further  
318 increase the effectiveness of this EO at 2.5 concentration. Finally, addition of other co-  
319 formulants could also increase the effect of WAC EO. For all these reasons, we proposed that  
320 the use of WAC 2.5 lead to interesting results which could be exploited proficiently for  
321 developing an eco-friendly herbicide. About possible mechanism of action, below we propose  
322 the physiological and biochemical reasons on the base of our data. For example, it has been  
323 demonstrated that essential oils might act as a desiccant herbicide which alters the leaf cuticular  
324 wax layer causing alterations in leaf membrane integrity, dehydration and death (Bainard et al.,  
325 2006). Moreover, considering that some species are affected more than others by the same EOs,  
326 varying the concentration of a given EOs might be useful to increase/reduce its selectivity  
327 allowing weed control without damaging crop growth and production. Recent studies  
328 demonstrated that the impairment of photosynthetic process is one of the main cascade effects  
329 of multi-target EOs (Araniti et al., 2018). Therefore, in principle we decided to investigate the  
330 effects of peppermint and caraway EOs in relation to changes induced in chlorophyll  
331 fluorescence parameters.

332

### 333 *3.3. The effect of emulsions on chlorophyll a fluorescence*

334 In Figures 3-6 the detailed images of chlorophyll *a* fluorescence parameters monitored on  
335 both maize and *E. crus-galli* leaves after spraying are displayed.

336 In particular, three specific fluorescence parameters were monitored: i) Fv/Fm – maximum  
337 quantum yield of photosystem II, that is the most common fluorescence parameter to measure  
338 response of plants to different kinds of stress (Kalaji et al., 2014) ii) NPQ – non-photochemical  
339 quenching, a parameter which is principally associated with the dissipation of the excess of

340 excitation energy in the form of heat (Müller et al., 2001), and iii) Rfd – fluorescence decrease  
341 ratio, which can be considered as a measure of the photosynthetic activity of a whole leaf  
342 (Lichtenthaller et al., 2005).

343 False color images of chlorophyll fluorescence suggest a decline of all these parameters in  
344 leaves of maize when treated with peppermint EO at 5% and with caraway EO at both 2.5 and  
345 5%, as revealed by the reduced surface of the leaves which still emits a fluorescence signal (Fig.  
346 7). It is conceivable that this is related to the occurrence of symptoms over the leaf and to the  
347 pre-symptomatic reduction of photosynthetic efficiency in areas of the leaf laminae where  
348 symptoms will consequently appear. Similar results can be observed in *E. crus-galli* leaves,  
349 which differently from maize leaves, had a lower chlorophyll fluorescence signal, even when  
350 sprayed with 2.5 % of peppermint EO (Fig. 8). A conversion of these false color-RGB images  
351 into the “mean grey value” enabled a statistical comparison between treatments (Figs 7-8) that  
352 included both the area of a living leaf-tissue and the intensity of color. The statistical analysis  
353 in “mean grey values” confirms our previous observation made by false color images revealing  
354 that photosystem II performances, a key indicator of photosynthetic efficiency, was less  
355 affected by the treatment with caraway as compared to peppermint EO. Moreover, it confirms  
356 that the lower dose of caraway emulsion (WAC2.5) did not alter two out of the three  
357 fluorescence parameters in maize leaves (Fig. 7). On the contrary, Fv/Fm and Rfd declined in  
358 a dose-dependent manner in *E. crus-galli*, whose photosynthetic apparatus seems much more  
359 susceptible than that of maize to peppermint EOs emulsions (Fig. 8). A reduction in Fv/Fm as  
360 well as alteration of other chlorophyll fluorescence parameters have been observed by several  
361 authors on plants treated with both EOs or their pure constituents. Araniti et al. (2017c, 2018)  
362 reported that oregano essential oils as well as *D. viscosa* volatiles strongly affected the  
363 photosynthetic machinery of Arabidopsis and lettuce, principally reducing the efficiency of  
364 both dark and light adapted PSII and NPQ. Similar results were observed by Graña et al. (2013)



365 and Araniti et al. (2017b) on *Arabidopsis* plants treated with the terpenoids citral and *trans*-  
366 caryophyllene, respectively. Finally, Synowiec et al. (2015) highlighted that leaf-application of  
367 clove oil and its main constituents caused a significant alteration of fluorescence parameters.  
368 Therefore, the results of both experiments showed that fluorescence parameters are not only  
369 indicative of the early reaction of the photosynthetic apparatus to the stress caused by the foliar  
370 application of EOs, but also allow to display differences in the sensitivity of plant species to the  
371 individual EOs.

372 As a next step, a more detailed metabolomic analyses revealed differences in key  
373 biochemical pathways altered by the foliar-applied emulsions of caraway or peppermint EOs.

374

#### 375 3.4. Differential effects of essential oils on plant metabolism

376 The GC-MS analysis was performed to identify differentially produced metabolites,  
377 following leaf spraying with EO emulsions and adjuvant. In particular, we screened the effect  
378 of the highest concentration of both EOs (5%), which caused significant increase of leaf injury,  
379 and in consequence the reduction of biomass, in order to inspect the main compounds as well  
380 as the pathway differentially affected by EOs treatments in both maize and *E. crus-galli* (Tables  
381 2-5 and Figures 9-10).

382 In order to assess the influence of the treatments on overall metabolites, raw data were  
383 analyzed through principal component analysis (PCA) and successively significant features  
384 were identified through the univariate analysis ANOVA (analysis of variance). Finally, to get  
385 more insights into the metabolic pathways affected by the treatments, data were analyzed  
386 through the “pathway analysis” (Tables 2-5 and Figures 9-10).

387 GC-MS analysis led to the identification of 51 and 52 compounds in *Z. mays* and *E. crus-*  
388 *galli*, respectively (Table 2, 4 and Figures 9, 10). In particular (out of the parenthesis are  
389 reported the number of metabolites annotated in *Z. mays*, whereas in the parenthesis those in *E.*

390 *crus-galli*), 10 (13) amino acids, 13 (12) organic acids, 12 sugars, 3 (1) sugar acid, 5 (4) sugar  
391 alcohols, 3 (4) amines, 2 (4) fatty acids, 1 glycan, 1 glycoside and 1 lactone for maize and 1  
392 inorganic acid for *E. crus-galli* have been annotated (Tables 2, 4 and Figures 9, 10).

393

#### 394 3.4.1. Metabolic characterization of treated and non-treated maize

395 Concerning the results obtained from *Z. mays*, the PCA analysis pointed out a clear  
396 separation among all treatments and the combination of the two principal components PC1  
397 (55.2%) vs PC2 (26.3%) explained a total variance of 81.5% (Fig. 9A).

398 In *Z. mays* experiments the PCA loading plot highlighted that sample separation was  
399 mainly due to lactose, erythritol, ribono-1,4-lactone, glyceryl-glycoside, 2-oxoglutaric acid,  
400 silanamine, sedoheptulose, tagatose and fructose for the PC1, whereas in PC2 it was due to  
401 malonic acid, galactinol, acotinic acid, tagatose, maltose, valine, aspartic acid and arabitol (Fig.  
402 9B). Both *Z. mays* PCA and heatmap visualization of metabolomic data showed distinct  
403 segregation between control and treated seedlings (Fig. 9C). At a higher level the metabolome  
404 of WAC5.0-treated seedlings and the peppermint treatment (WAP5.0) clustered together,  
405 suggesting that the treatments completely changed the metabolic profile of treated plants  
406 compared to control plants (Fig. 9C). The univariate analysis of variance (ANOVA), carried  
407 out on *Z. mays* seedlings treated with both caraway (WAC5.0) and peppermint (WAP5.0) EOs,  
408 pointed out several statistical and contrasting differences among control and treatments (Table  
409 2). In particular, in seedlings treated with WAC5.0 three amino acids were significantly  
410 stimulated (glutamate, serine and L-alanine), whereas aspartic acid and norvaline were  
411 significantly reduced (Table 2). On the contrary, in WAP5.0 treated plants only the aspartic  
412 acid and valine were significantly reduced, whereas serine was accumulated (Table 2). A  
413 similar trend was also observed in organic acids and sugars contents which were differentially

414 affected by the treatments (Table 2), suggesting that the two EOs were able to interfere with  
415 different metabolic pathways.

416 This hypothesis was confirmed by the pathway analysis, which highlighted that caraway  
417 EOs significantly affected more metabolic pathways than peppermint EO (11 vs 7) (Table 3).  
418 In particular, both treatments significantly interfered with the amino acid metabolism (e.g.  
419 alanine, aspartate and glutamate metabolism as well as glycine, serine and threonine  
420 metabolism). On the contrary, the citrate cycle, the inositol phosphate metabolism, the starch  
421 and sucrose metabolism as well as the glycerolipid metabolism were only affected by caraway  
422 treatment (Table 3).

423 The highest accumulation of metabolites observed in maize plants treated with caraway  
424 EOs suggests that plants were less affected by this treatment as compared to those treated with  
425 peppermint EOs, which is in agreement with chlorophyll fluorescence data and the biomass  
426 changes observed in this study. In fact, the increase in glutamic acid, serine and alanine as well  
427 as the increments in sugars has been reported as an adaptation strategy adopted by resistant  
428 plants to cope with abiotic stress since they act as osmoprotectants (Kovács et al., 2012; Good  
429 and Zaplachinski, 1994; Rhodes et al., 1986). Moreover, the accumulation of maltose might  
430 hint at an enhanced potential for starch mobilization in plants when exposed to EO stress. It has  
431 been suggested that in plants  $\beta$ -amylase induction during biotic stress could lead to starch-  
432 dependent maltose accumulation, and that maltose might contribute in protecting proteins and  
433 the electron transport chain in the chloroplast stroma during acute stress (Kaplan and Guy,  
434 2005).

435 The high accumulation in maltose only observed in maize plants treated with caraway EOs  
436 suggests that this species has a higher ability to cope with the stress induced by this EOs-  
437 formulation. This hypothesis is strongly supported also by the weaker effects (in terms of leaf

438 injuries, reduction in biomass, impact to PSII efficiency etc.) induced by WAC treatment when  
439 assayed at the lower concentration (WAC2.5).

440

#### 441 3.4.2. Metabolic characterization of treated and non-treated *E. crus-galli*

442 The score plot of the unsupervised PCA (Fig. 10a) highlights a clear separation among  
443 control and treatments. In the experiments carried on *E. crus-galli* (Fig. 10a) the separation was  
444 achieved using the principal components (PCs) PC1 vs PC2, which explained a total variance  
445 of 80.2%. In particular, PC1 explained the highest variance (54.3 %) while PC2 explained  
446 25.9% of the total variance. In Fig. 10b is reported the PCA loading plot which highlighted that  
447 the PC1 was dominated by maltose, asparagine, glucose, mannose, fructose, tagatose, arabinose  
448 and tyrosine, whereas PC2 was dominated by turanose, galactinol, lactic acid, malonic acid,  
449 quinic acid and glutamine.

450 In the heatmap reported in Fig. 10c visualization of metabolomic data showed distinct  
451 segregation and a peculiar clusterization among control and treatments WAC5.0 and WAP5.0.  
452 Agglomerative hierarchical clustering begins with each sample as separate cluster and then  
453 proceeds to combine them until all samples belong to one cluster. At a higher level the  
454 metabolome of control seedlings (WA) and the treatment WAC5.0 clustered together.

455 As for *Z. mays*, the ANOVA pointed out a high number of statistically affected metabolites in  
456 *E. crus galli* WAP5.0-treated plants. Interestingly, in plants treated with caraway EO a general  
457 reduction in both amino acids and sugars content was observed, whereas in plants treated with  
458 caraway EO an opposite behavior was observed, which was also characterized by a general  
459 reduction of both classes of compounds (Table 4). The strong downregulation of different  
460 metabolic pathways leading to the biosynthesis of amino acid and sugars are likely on the bases  
461 of the strong effectiveness of the caraway EO treatment, confirming again the multitarget nature  
462 of this EO. Concerning the organic acids, in both treatments it was observed a reduction in

463 acotinic acid, malate and malonate as well as an accumulation of citric acid and glycolic acid  
464 (Table 4). In addition, in WAP5.0-treated plants it was also observed an increment in galactinol  
465 and glycerol content, while myo-inositol was reduced by both treatments and sorbitol content  
466 was increased only in WAC5.0-treated plants (Table 4).

467 Finally, the pathway analysis highlighted that treatments carried out on the plants of *E.*  
468 *crus-galli* differentially affected several pathways (Table 5). In particular, both emulsions with  
469 caraway and peppermint EOs affected alanine, aspartate and glutamate metabolism as well as  
470 the galactose metabolism (Table 5). On the other hand, glycine, serine and threonine  
471 metabolism as well as isoquinoline alkaloid biosynthesis was significantly affected only by the  
472 emulsions with peppermint EOs (Table 5).

473 Interestingly, both EOs had a quite completely different effects on *E. crus-galli* seedlings.  
474 In fact, in WAP5.0 treatment a high accumulation of almost all the amino acids and sugars was  
475 observed, whereas in WAC5.0 treated plants an opposite behavior was observed. On the  
476 contrary, organic acid content followed a similar trend in both treatments. As previously  
477 reported amino acid and sugar accumulation plays a pivotal role in protecting plants from  
478 oxidative stress acting as osmoprotectants (Kovács et al., 2012; Good and Zaplachinski, 1994;  
479 Rhodes et al., 1986). Moreover, in plants treated with peppermint EOs (WAP5.0) a higher  
480 increase in sucrose, galactinol and glycerol was observed. These molecules are important plant  
481 protectors during several abiotic stress such as salinity, heat- and cold-shock stress (Nishizawa  
482 et al., 2008; Eastmond, 2004; Taji et al., 2002; Santarius, 1992). Santarius (1992) reported that  
483 sucrose and glycerol, which easily penetrate across chloroplast membranes, strongly protected  
484 isolated thylakoid membranes from cold shock preventing membrane damages and stabilizing  
485 protein complex. In addition, Eastmond (2004) demonstrated that *Arabidopsis* mutants, which  
486 accumulate glycerol, were more resistant to abiotic stresses associated with leaf dehydration.  
487 Concerning galactinol, it has been suggested that this molecule not only acts as osmoprotectant

488 and stabilizer of cellular membranes, but is also a pivotal ROS scavenger playing a novel role  
489 in the protection of cellular metabolism, in particular the photosynthetic apparatus, from  
490 oxidative damages caused by several abiotic stress factors (Nishizawa et al., 2008; Taji et al.,  
491 2002).

492 These results suggest that *E. crus-galli* plants exposed to peppermint EOs were able to cope  
493 with EO-promoting stress by activating some metabolic strategy aiming to enable plant  
494 protection. These results are in agreement with leaf injuries and plant biomass results. In fact,  
495 leaf injuries induced by WAP5.0 treatment were significantly lower compared to those  
496 exhibited by WAC5.0-treated plants. On the other hand, biomass in plants treated with  
497 peppermint EO was significantly lower than that of WAC5.0-treated plants. Probably, as also  
498 suggested by Good and Zaplachinsky (1992) plants underwent a series of reaction which finally  
499 lead to a reduction of protein synthesis in order to increase the amino acid content to cope and  
500 protect themselves from EOs-induced osmotic and oxidative stress.

501

## 502 **5. Conclusions**

503 The dataset presented here offers clear evidence that foliar-applied oil-in-water (o/w)  
504 emulsions containing peppermint EO and fatty acid methyl esters strongly affect both species,  
505 maize and barnyard grass at the growth phases of leaves development, from both a  
506 physiological and biochemical point of view. On the contrary, o/w emulsions containing  
507 caraway EOs (WAC) were more effective on *E. crus-galli* at both concentrations, causing leaf  
508 injuries and reduction in biomass as well as significant alterations on the photosynthetic  
509 apparatus and plant metabolism, whereas biomass as well as photosynthetic apparatus of *Z.*  
510 *mays* seedlings were not affected by WAC2.5, as compared to control plants. Moreover, despite  
511 the presence of some injuries on leaf blades after WAC5.0 treatment, maize seedlings were able  
512 to activate metabolic mechanisms, such as amino acids and sugars accumulation, to protect

513 themselves from EOs-induced stress. Taken together these results suggests that the o/w  
514 emulsion based on caraway EO and fatty acids methyl esters represents a potential candidate  
515 for the development of a commercial botanical herbicide against *E. crus-galli* in maize  
516 cultivation.

517

#### 518 **Conflict of interest statement**

519 The authors declare no conflicts of interest.

520

#### 521 **Acknowledgements**

522 This work was supported by the Polish Ministry of Science and Higher Education as part of the  
523 statutory activities of the University of Agriculture in Krakow [grant number DS 3124].

524

#### 525 **Authors contribution**

526 AS designed and performed the pot experiment, statistically analyzed results and wrote part of  
527 the manuscript; FA designed, performed and statistically analyzed the results of metabolomic  
528 analyses, and wrote part of the manuscript; KM and ML performed and analyzed the  
529 chlorophyll fluorescence analysis and wrote part of the manuscript; AK performed the chemical  
530 analyses of essential oils and wrote methodological part for this analysis.

531

#### 532 **References**

533 Adams, R.P., 2007. Alphabetical listing of compounds with their retention time and arithmetic  
534 retention index on DB-5. Identification of essential oil components by gas  
535 chromatography/mass spectrometry 4, 401.

536 Ahuja, N., Singh, H.P., Batish, D.R., & Kohli, R.K., 2015. Eugenol-inhibited root growth in  
537 *Avena fatua* involves ROS-mediated oxidative damage. Pest. Biochem. Physiol. 118, 64-  
538 70.

539 Araniti, F., Bruno, L., Sunseri, F., Pacenza, M., Forgione, I., Bitonti, M.B., & Abenavoli, M.R.,  
540 2017a. The allelochemical farnesene affects *Arabidopsis thaliana* root meristem altering  
541 auxin distribution. Plant Physiol. Biochem. 121, 14-20.

542 Araniti, F., Graña, E., Krasuska, U., Bogatek, R., Reigosa, M.J., Abenavoli, M.R., & Sánchez-  
543 Moreiras, A.M., 2016. Loss of gravitropism in farnesene-treated arabidopsis is due to  
544 microtubule malformations related to hormonal and ROS unbalance. PloS one 11(8),  
545 e0160202.

546 Araniti, F., Graña, E., Reigosa, M.J., Sánchez-Moreiras, A.M., & Abenavoli, M.R., 2013.  
547 Individual and joint activity of terpenoids, isolated from *Calamintha nepeta* extract, on  
548 *Arabidopsis thaliana*. Nat. Prod. Res. 27(24), 2297-2303.

549 Araniti, F., Landi, M., Lupini, A., Sunseri, F., Guidi, L., & Abenavoli, M.R., 2018. *Origanum*  
550 *vulgare* essential oils inhibit glutamate and aspartate metabolism altering the  
551 photorespiratory pathway in *Arabidopsis thaliana* seedlings. J. Plant Physiol. 231, 297-  
552 309.

553 Araniti, F., Sánchez-Moreiras, A.M., Graña, E., Reigosa, M.J., & Abenavoli, M.R., 2017b.  
554 Terpenoid *trans*-caryophyllene inhibits weed germination and induces plant water status  
555 alteration and oxidative damage in adult Arabidopsis. Plant Biol. 19(1), 79-89.

556 Araniti, F., Lupini, A., Sunseri, F., Abenavoli, M.R., 2017c. Allelopathic Potential of *Dittrichia*  
557 *viscosa* (L.) W. Greuter Mediated by VOCs: A Physiological and Metabolomic  
558 Approach. PloS one. 12(1). e0170161.



559 Bainard, L. D., Isman, M. B., & Upadhyaya, M. K., 2006. Phytotoxicity of clove oil and its  
560 primary constituent eugenol and the role of leaf epicuticular wax in the susceptibility to  
561 these essential oils. *Weed Sci.* 54(5), 833-837.

562 Baj, T., Sieniawska, E., Kowalski, R., Wesołowski, M., & Ulewicz-Magulska, B., 2015.  
563 Effectiveness of the Deryng and Clevenger-type apparatus in isolation of various types  
564 of components of essential oil from the *Mutelina purpurea* Thell. flowers. *Acta Pol.*  
565 *Pharm.* 72, 507-515.

566 Benvenuti, S., Cioni, P. L., Flamini, G., & Pardossi, A., 2017. Weeds for weed control:  
567 Asteraceae essential oils as natural herbicides. *Weed Res.* 57(5), 342-353.

568 Chemat, S., Lagha, A., AitAmar, H., Bartels, P.V., & Chemat, F., 2004. Comparison of  
569 conventional and ultrasound-assisted extraction of carvone and limonene from caraway  
570 seeds. *Flav. Fragr. J.* 19(3), 188-195.

571 Eastmond, P. J., 2004. Glycerol-insensitive Arabidopsis mutants: gli1 seedlings lack glycerol  
572 kinase, accumulate glycerol and are more resistant to abiotic stress. *Plant J.* 37(4), 617-  
573 625.

574 Fejér, J., Gruľová, D., & De Feo, V., 2017. Biomass production and essential oil in a new bred  
575 cultivar of peppermint (*Mentha × piperita* L.). *Ind. Crops and Prod.* 109, 812-817.

576 Ghatak, A., Chaturvedi, P., & Weckwerth, W., 2018. Metabolomics in Plant Stress Physiology:  
577 1-50.

578 Good, A.G., & Zaplachinski, S. T., 1994. The effects of drought stress on free amino acid  
579 accumulation and protein synthesis in *Brassica napus*. *Physiol. Plantarum* 90(1), 9-14.

580 Graña, E., Sotelo, T., Díaz-Tielas, C., Reigosa, M.J., & Sánchez-Moreiras, A.M., 2013. The  
581 phytotoxic potential of the terpenoid citral on seedlings and adult plants. *Weed Sci.* 61(3),  
582 469-481.

583 Guidi, L., & Landi, M., 2014. Aromatic plants: use and nutraceutical properties. *Novel Plant*  
584 *Bioresources: Applications in Food, Medicine and Cosmetics*, 303-345.

585 Hazrati, H., Saharkhiz, M.J., Niakousari, M., & Moein, M., 2017. Natural herbicide activity of  
586 *Satureja hortensis* L. essential oil nanoemulsion on the seed germination and  
587 morphophysiological features of two important weed species. *Ecotox. Environ.*  
588 *Safe*. 142, 423-430.

589 Ibrahim, M.A., Oksanen, E.J., & Holopainen, J.K., 2004. Effects of limonene on the growth  
590 and physiology of cabbage (*Brassica oleracea* L) and carrot (*Daucus carota* L) plants. *J.*  
591 *Sci. Food Agricul.* 84(11), 1319-1326.

592 Kalaji, H.M., Schansker, G., Ladle, R.J., Goltsev, V., Bosa, K., Allakhverdiev, S., I, Brestic,  
593 M., Bussotti, F., Calatayud, A., Dąbrowski, P., Elsheery, N.I., Ferroni, L., Guidi, L.,  
594 Hogewoning, S.W., Jajoo, A., Misra, A.N., Nebauer, S.G., Pancaldi, S., Penella, C., Poli,  
595 D.B., Pollastrini, M., Romanowska-Duda, Z.B., Rutkowska, B., Serôdio, J., Suresh, K.,  
596 Szulc, W., Tambussi, E., Yanniccari, M., Zivcak, M., 2014. Frequently asked questions  
597 about in vivo chlorophyll fluorescence: practical issues. *Photosynth. Res.* 122, 121–158.

598 Kaplan, F., & Guy, C.L., 2005. RNA interference of Arabidopsis beta-amylase8 prevents  
599 maltose accumulation upon cold shock and increases sensitivity of PSII photochemical  
600 efficiency to freezing stress. *Plant J.* 44(5), 730-743.

601 Kaur, S., Singh, H.P., Mittal, S., Batish, D.R., & Kohli, R.K., 2010. Phytotoxic effects of  
602 volatile oil from *Artemisia scoparia* against weeds and its possible use as a bioherbicide.  
603 *Ind. Crops and Prod.* 32(1), 54-61.

604 Kovács, Z., Simon-Sarkadi, L., Vashegyi, I., & Kocsy, G., 2012. Different accumulation of free  
605 amino acids during short-and long-term osmotic stress in wheat. *Sci. World J.* 2012: 1-  
606 10.

607 Lichtenthaler, H.K., Buschmann, C., Knapp, M., 2005. How to correctly determine the different  
608 chlorophyll fluorescence parameters and the chlorophyll fluorescence decrease ratio Rfd  
609 of leaves with the PAM fluorometer. *Photosynthetica* 43, 379–393.

610 Mosa, K.A., Ismail, A., & Helmy, M., 2017. Omics and system biology approaches in plant  
611 stress research. In: *Plant Stress Tolerance* (pp. 21-34). Springer, Cham.

612 Müller, P., Li, X.P., Niyogi, K.K., 2001. Non-photochemical quenching. A response to excess  
613 light energy. *Plant Physiol.* 125, 1558–1566.

614 Nicholson, J. K., & Lindon, J.C., 2008. Systems biology: metabolomics. *Nature*, 455(7216),  
615 1054.

616 Nishizawa, A., Yabuta, Y., & Shigeoka, S., 2008. Galactinol and raffinose constitute a novel  
617 function to protect plants from oxidative damage. *Plant Physiol.* 147(3), 1251-1263.

618 Oroian, C., Covrig, I., Odagiu, A., Mălinaş, C., Moldovan, C., & Fleşeriu, A., 2017. Effects of  
619 cultivation systems and environmental conditions on peppermint (*Mentha× piperita* L.)  
620 biomass yield and oil content. *Notulae botanicae horti agrobotanici cluj-napoca*, 45(2),  
621 576-581.

622 Pisulewska E., Fijołek M., Witkowicz R., 2010. The share of peppermint (*Mentha piperita* L.)  
623 in the cropping structure and its cultivation technology in the Michalow commune.  
624 *Zeszyty Problemowe Postępów Nauk Rolniczych* 549, 149-156.

625 R Core Team, 2014. R: A language and environment for statistical computing. R Foundation  
626 for Statistical Computing, Vienna, Austria. URL <http://www.R-project.org/>.

627 Rhodes, D., Handa, S., & Bressan, R.A., 1986. Metabolic changes associated with adaptation  
628 of plant cells to water stress. *Plant Physiol.* 82(4), 890-903.

629 Rolli, E., Marieschi, M., Maietti, S., Sacchetti, G., Bruni, R., 2014. Comparative phytotoxicity  
630 of 25 essential oils on pre-and post-emergence development of *Solanum lycopersicum* L.:  
631 a multivariate approach. *Ind. Crops and Prod.* 60, 280–290.

632 Santarius, K.A., 1992. Freezing of isolated thylakoid membranes in complex media. VIII.  
633 Differential cryoprotection by sucrose, proline and glycerol. *Physiol. Plantarum*, 84(1),  
634 87-93.

635 Schulz, M., Kussmann, P., Knop, M., Kriegs, B., Gresens, F., Eichert, T., Ulbrich, A., Marx,  
636 F., Fabricius, H., Goldbach, H. & Noga, G., 2007. Allelopathic monoterpenes interfere  
637 with *Arabidopsis thaliana* cuticular waxes and enhance transpiration. *Plant signaling &*  
638 *behavior*, 2(4), 231-239.

639 Seidler-Łożykowska, K., & Bocianowski, J., 2012. Evaluation of variability of morphological  
640 traits of selected caraway (*Carum carvi* L.) genotypes. *Ind. Crops and Prod.* 35(1), 140-  
641 145.

642 Seidler-Łożykowska, K., Kędzia, B., Karpińska, E., & Bocianowski, J., 2013. Microbiological  
643 activity of caraway (*Carum carvi* L.) essential oil obtained from different origin. *Acta*  
644 *Sci. Agr.* 35(4), 495-500.

645 Stokłosa, A., Matraszek, R., Isman, M.B., & Upadhyaya, M.K., 2012. Phytotoxic activity of  
646 clove oil, its constituents, and its modification by light intensity in broccoli and common  
647 lambsquarters (*Chenopodium album*). *Weed Sci.* 60(4), 607-611.

648 Sunohara, Y., Baba, Y., Matsuyama, S., Fujimura, K., & Matsumoto, H., 2015. Screening and  
649 identification of phytotoxic volatile compounds in medicinal plants and characterizations  
650 of a selected compound, eucarvone. *Protoplasma*, 252(4), 1047-1059.

651 Synowiec, A. & Drozdek, E. 2016. Physicochemical and herbicidal properties of emulsions of  
652 essential oils against *Avena fatua* L. and *Chenopodium album* L. *J. Plant Dis. Prot.* 123(2),  
653 65-74.

654 Synowiec, A., Kalemba, D., Drozdek, E., & Bocianowski, J., 2017. Phytotoxic potential of  
655 essential oils from temperate climate plants against the germination of selected weeds and  
656 crops. *J. Pest Sci.* 90(1), 407-419.

657 Synowiec, A., Możdżeń, K., & Skoczowski, A., 2015. Early physiological response of broccoli  
658 leaf to foliar application of clove oil and its main constituents. *Ind. Crops and Prod.* 74,  
659 523-529.

660 Taji, T., Ohsumi, C., Iuchi, S., Seki, M., Kasuga, M., Kobayashi, M., Yamaguchi-Shinozaki,  
661 K. & Shinozaki, K., 2002. Important roles of drought-and cold-inducible genes for  
662 galactinol synthase in stress tolerance in *Arabidopsis thaliana*. *Plant J.* 29(4), 417-426.

663 Vaid, S., Batish, D.R., Singh, H.P., & Kohli, R.K., 2011. Phytotoxicity of limonene against  
664 *Amaranthus viridis* L. *The Bioscan*, 6(1), 163-165.

665 Vaughn, S.F., & Spencer, G.F., 1993. Volatile monoterpenes as potential parent structures for  
666 new herbicides. *Weed Sci.* 41(1), 114-119.

667 Vokou, D., Douvli, P., Blionis, G.J., & Halley, J.M., 2003. Effects of monoterpenoids, acting  
668 alone or in pairs, on seed germination and subsequent seedling growth. *J. Chem. Ecol.*  
669 29(10), 2281-2301.

670 Xia, J., Sinelnikov, I.V., Han, B., & Wishart, D.S., 2015. MetaboAnalyst 3.0—making  
671 metabolomics more meaningful. *Nucl. Acids Res.* 43(W1). W251-W257.

672 **Table 1**

673 Main constituents (%) of caraway seed and peppermint herb essential oils distilled from  
 674 plants grown in the temperate climate.

Caraway oil (WAC5.0)				Peppermint oil (WAP5.0)			
RI	RI <sub>lit</sub>	Constituent	%	RI	RI <sub>lit</sub>	Constituent	%
927	934	$\alpha$ -Pinene	0.1	927	924	$\alpha$ -Thujene	0.3
963	970	Sabinene	0.1	939	934	$\alpha$ -Pinene	t
966	974	$\beta$ -Pinene	0.1	962	970	Sabinene	0.2
982	983	Myrcene	0.1	965	974	$\beta$ -Pinene	0.7
1001	1006	Car-3-ene	t	982	983	Myrcene	0.1
1011	1016	p-Cymene	t	994	998	$\alpha$ -Phellandrene	t
1023	1025	Limonene	32.5	1006	1006	Car-3-ene	0.1
1086	1086	Linalool	t	1009	1015	p-Cymene	0.2
1101		p-Mentha-2,8-					
	1103	dien-1-ol	t	1016	1025	1.8-Cyeneol	5.2
1112		cis-Limonen					
	1116	oxide	0.1	1018	1025	Limonene	2.1
1117		trans-Limoene					
	1121	oxide	t	1026	1025	(Z)- $\beta$ -Ocimene	t
1168		cis-					
	1172	Dihydrocarvone	0.1	1046	1055	$\gamma$ -Terpinene	0.2
1174		trans-					
	1177	Dihydrocarvone	0.1	1049	1053	trans-Sabinene hydrate	0.3
1201		Dihydrocarveol					
		(isomer)	t	1076	1081	Terpinolene	0.1
1206	1210	cis-Carveol	0.1	1080	1082	cis-Sabinene hydrate	t

1224	1218	Carvone	66.4	1084	1087	Linalool	0.1
1413		(E)- $\beta$ -				2-Methylbutyl-2-methyl	
	1421	Caryophyllene	t	1087	1091	isobutyrate	t
1446	1446	(E)- $\beta$ -Farnesene	t	1091	1094	2-Methylbutylisovalerate	t
1565		$\beta$ -Caryophyllene					
	1546	oxide	t	1105	1108	cis-p-Menth-2-en-1-ol	t
				1125	1129	trans-Sabinol	t
				1134	1139	Menthone	25.5
				1140	1146	Isomenthone	4.0
				1144	1150	Menthofuran	2.0
				1148	1056	Neomenthol	3.3
				1163	1163	Menthol	42.7
				1167	1176	Isomenthol	0.4
				1171	1176	$\alpha$ -Terpineol	0.4
				1188	1176	Neoisomenthol	t
				1210	1215	Pulegone	0.8
				1223	1226	Piperiton	0.4
						Isopulegol acetate (Isomer	
				1253	1259	D)	t
				1256	1263	Neomenthyl acetate	0.3
				1276	1280	Menthyl acetate	5.9
				1289	1298	Isomenthyl acetate	0.2
				1372	1380	$\alpha$ -Copaene	t
				1379	1386	$\beta$ -Bourbonene	0.2
				1384	1389	$\beta$ -Elemene	0.1

1414	1421	(E)- $\beta$ -Caryophyllene	2.2
1422	1430	$\beta$ -Copaene	t
1436	1445	Isogermacrene D	t
1444	1446	(E)- $\beta$ -Farnesene	0.1
1446	1455	$\alpha$ -Humulene	0.1
1466	1474	$\gamma$ -Muurolene	t
1471	1479	Germacrene D	0.9
1486	1494	Bicyclogermacrene	0.1
1489	1496	$\alpha$ -Muurolene	t
1491	1497	$\alpha$ -Cuprenene	0.1
1509	1507	$\gamma$ -Cadinene	t
1561	1572	Spathulenol	t
1565	1578	Caryophyllene oxide	0.1
1576	1589	Globulol	0.1

Sum of 99.7  
constituents

Sum of constituents 99.5



676 **Table 2**

677 Effects of water plus adjuvant (WA), caraway (WAC5.0) and peppermint  
 678 (WAP5.0) essential oil on *Zea mays* metabolites content.

Feature	WA	WAC5.0	WAP5.0	Class
Isoleucine	1.32	0.81	0.71	<b>Amino acid</b>
Aspartic acid	8.65 <sup>b</sup>	7.96 <sup>b</sup>	4.88 <sup>a</sup>	
Glutamic acid	15.15 <sup>a</sup>	20.68 <sup>b</sup>	13.78 <sup>a</sup>	
Valine	0.82 <sup>b</sup>	0.88 <sup>b</sup>	0.48 <sup>a</sup>	
Serine	4.07 <sup>a</sup>	9.44 <sup>b</sup>	10.72 <sup>b</sup>	
L-Alanine	9.94 <sup>a</sup>	17.82 <sup>b</sup>	10.00 <sup>a</sup>	
Norvaline	1.22 <sup>b</sup>	0.72 <sup>a</sup>	1.16 <sup>b</sup>	
Threonine	2.32	4.16	2.07	
Glycine	3.76	3.49	3.46	
Pyroglutamic acid	5.24	7.67	6.35	
Aconitic acid	217.97 <sup>c</sup>	170.33 <sup>b</sup>	90.75 <sup>a</sup>	<b>Organic acid</b>
Cinnamic acid	2.51	2.65	2.49	
Carbamate	17.69 <sup>b</sup>	9.82 <sup>a</sup>	15.81 <sup>b</sup>	
Citric acid	16.13 <sup>b</sup>	13.23 <sup>a</sup>	12.26 <sup>a</sup>	
cyclohexanecarboxylicacid	108.71	116.03	71.46	
Itaconic acid	1.68	5.81	2.83	
Malic acid	105.73	110.11	112.04	
Malonic acid	1.09 <sup>b</sup>	0.33 <sup>a</sup>	0.19 <sup>a</sup>	
Oxalic acid	50.38 <sup>a</sup>	66.47 <sup>b</sup>	73.63 <sup>b</sup>	
2-Oxoglutaric acid	0.77 <sup>a</sup>	3.00 <sup>b</sup>	1.94 <sup>b</sup>	
Quinic acid	282.40	249.73	250.44	
Succinic acid	3.41	6.79	5.51	
Threonic acid	18.48	18.13	12.98	
Glycolic acid	2.25 <sup>b</sup>	0.93 <sup>a</sup>	0.93 <sup>a</sup>	
Lactate	15.75 <sup>a</sup>	24.15 <sup>b</sup>	15.32 <sup>a</sup>	
Arabinose	4.35 <sup>a</sup>	8.30 <sup>b</sup>	9.29 <sup>b</sup>	
Fructose	763.56 <sup>c</sup>	223.89 <sup>a</sup>	479.78 <sup>b</sup>	
Glucose	1128.24 <sup>c</sup>	352.92 <sup>a</sup>	907.93 <sup>b</sup>	

Inosose	6.96	7.49	5.40	
Lactose	3.23 <sup>a</sup>	21.00 <sup>c</sup>	5.44 <sup>b</sup>	
Lyxose	1.95 <sup>b</sup>	0.86 <sup>a</sup>	2.52 <sup>b</sup>	
Maltose	10.77 <sup>a</sup>	29.16 <sup>b</sup>	7.54 <sup>a</sup>	
Mannobiose	4.20 <sup>a</sup>	8.43 <sup>b</sup>	3.44 <sup>a</sup>	
Sedoheptulose	3.88 <sup>a</sup>	12.89 <sup>b</sup>	4.09 <sup>a</sup>	
Sucrose	409.95	283.04	363.48	
Tagatose	23.08 <sup>b</sup>	4.97 <sup>a</sup>	3.78 <sup>a</sup>	
Threose	76.50 <sup>c</sup>	41.85 <sup>a</sup>	65.47 <sup>b</sup>	
<hr/>				
Glyceric acid	6.45	11.80	7.86	
Threonic acid	18.48	18.13	12.98	<b>Sugar Acid</b>
Erythronic acid	7.57 <sup>a</sup>	15.37 <sup>b</sup>	8.65 <sup>a</sup>	
<hr/>				
Arabitol	1.58 <sup>a</sup>	1.47 <sup>a</sup>	2.98 <sup>b</sup>	
Dithioerythritol	2.69 <sup>a</sup>	14.42 <sup>c</sup>	6.69 <sup>b</sup>	
Galactinol	2.32 <sup>b</sup>	3.03 <sup>b</sup>	0.93 <sup>a</sup>	<b>Sugar alcohol</b>
Glycerol	2.76	4.86	2.84	
Myoinositol	59.06 <sup>a</sup>	70.17 <sup>b</sup>	53.75 <sup>a</sup>	
<hr/>				
Ethanolamine	6.12	7.00	5.26	
Silanamine	1.82 <sup>a</sup>	6.30 <sup>b</sup>	5.54 <sup>b</sup>	<b>Amine</b>
Hydroxylamine	91.68	110.88	93.80	
<hr/>				
Octadecanoic acid	9.75 <sup>a</sup>	16.62 <sup>b</sup>	14.78 <sup>b</sup>	<b>Fatty acid</b>
Palmitic acid	17.72 <sup>a</sup>	22.34 <sup>b</sup>	20.79 <sup>b</sup>	
<hr/>				
Galacturonic acid	20.35 <sup>b</sup>	16.31 <sup>a</sup>	17.48 <sup>a</sup>	<b>Glycan</b>
<hr/>				
Ribono-1,4-lactone	0.21 <sup>a</sup>	1.08 <sup>b</sup>	0.70 <sup>b</sup>	<b>Lactone</b>
<hr/>				
Glyceryl-glycoside	10.05 <sup>a</sup>	58.02 <sup>c</sup>	34.41 <sup>b</sup>	<b>Glycoside</b>
<hr/>				
Phosphoric acid	21.85	22.32	21.88	<b>Inorganic acid</b>

679 Different letters along the rows indicate statistical differences with  $P \leq 0.05$  (LSD's test). N=3.

680 **Table 3**

681 Pathway analysis: result from “Pathway Analysis” carried on the concentrations of metabolite  
 682 identified in Zea mays treated with water plus adjuvant (WA), caraway (WAC5.0) and  
 683 peppermint (WAP5.0) essential oils. In the table are reported the results obtained through the  
 684 ingenuity pathway analysis carried out with MetPa.

Pathways	T Cmpd	Hits	WA vs		Impact
			WAC5.0 Raw p	WA vs WAP5.0 Raw p	
Alanine, aspartate and glutamate metabolism	21	5	0.014344	0.034342	0.66439
Glycine, serine and threonine metabolism	29	4	0.0034682	0.012128	0.53477
Citrate cycle (TCA cycle)	20	4	0.012643	//	0.24667
Inositol phosphate metabolism	17	1	0.039874	//	0.24503
Glyoxylate and dicarboxylate metabolism	17	4	0.0057461	0.0054114	0.23944
Galactose metabolism	26	6	0.0003485	0.0030181	0.1668
Methane metabolism	11	2	0.0052073	0.0089587	0.16667
Arginine and proline metabolism	37	2	0.053811	0.0056003	0.1268
Starch and sucrose metabolism	25	4	3.59E-01	//	0.11156
Glycerolipid metabolism	14	2	0.053595	//	0.09402
Aminoacyl-tRNA biosynthesis	67	8	0.0052887	0.013395	0.09302

685 T Cmpd: the total number of compounds in the pathway; Hits: is the actually matched number  
 686 from the uploaded data; Raw P: is the original p value calculated from the enrichment analysis;  
 687 Impact: is the pathway impact value calculated from pathway topology analysis.

688 **Table 4**

689 Effects of caraway (WAC5.0) and peppermint (KWAP5.0) essential oil on  
 690 *Echinocloa curs-galli* metabolites content.

Feature	WA	WAC5.0	WAP5.0	Class
Asparagine	1.53 <sup>b</sup>	0.61 <sup>a</sup>	9.66 <sup>c</sup>	
Aspartic acid	18.66	21.13	14.78	
GABA	4.79 <sup>a</sup>	7.18 <sup>a</sup>	16.91 <sup>b</sup>	
Glutamic acid	46.32 <sup>b</sup>	24.34 <sup>a</sup>	25.91 <sup>a</sup>	
Glutamine	3.7 <sup>a</sup>	12.09 <sup>b</sup>	24.30 <sup>c</sup>	
Glycine	2.60 <sup>a</sup>	2.80 <sup>a</sup>	5.90 <sup>b</sup>	
L-Alanine	24.22	20.72	35.24	<b>Amino acid</b>
Leucine	2.50	4.06	6.62	
Proline	19.69 <sup>a</sup>	32.08 <sup>a</sup>	53.85 <sup>b</sup>	
Serine	12.21 <sup>a</sup>	15.94 <sup>a</sup>	34.27 <sup>b</sup>	
Threonine	5.45 <sup>a</sup>	4.59 <sup>a</sup>	7.10 <sup>b</sup>	
Tyrosine	2.37 <sup>a</sup>	1.88 <sup>b</sup>	8.52 <sup>c</sup>	
Valine	19.30	17.49	19.24	
Aconitic acid	380.50 <sup>b</sup>	219.48 <sup>a</sup>	219.13 <sup>a</sup>	
Carbamate	16.72	12.20	13.76	
Citric acid	8.63 <sup>a</sup>	11.19 <sup>b</sup>	16.45 <sup>c</sup>	
Malic acid	37.11 <sup>b</sup>	21.64 <sup>a</sup>	24.30 <sup>a</sup>	
Malonic acid	0.93 <sup>b</sup>	0.39 <sup>a</sup>	0.50 <sup>a</sup>	
Methylmaleic acid	1.01	0.89	1.11	<b>Organic acid</b>
Oxalic acid	105.06 <sup>b</sup>	90.40 <sup>b</sup>	71.51 <sup>a</sup>	
Quinic acid	91.96 <sup>b</sup>	15.23 <sup>a</sup>	124.41 <sup>c</sup>	
Succinic acid	6.68 <sup>a</sup>	4.10 <sup>a</sup>	9.42 <sup>b</sup>	
Threonic acid	8.21	4.13	7.09	
Glycolic acid	0.77 <sup>a</sup>	1.43 <sup>b</sup>	2.12 <sup>c</sup>	
Lactic acid	159.93	27.43	34.14	
Arabinose	10.11 <sup>b</sup>	5.79 <sup>a</sup>	28.41 <sup>c</sup>	
Cellobiose	1.46	0.16	0.85	<b>Sugar</b>
Fructose	150.39 <sup>b</sup>	46.45 <sup>a</sup>	391.27 <sup>c</sup>	

Galactose	0.64 <sup>b</sup>	0.42 <sup>a</sup>	1.05 <sup>c</sup>	
Glucose	120.19 <sup>b</sup>	38.83 <sup>a</sup>	428.70 <sup>c</sup>	
Lactose	5.45 <sup>b</sup>	2.16 <sup>a</sup>	7.26 <sup>c</sup>	
Maltose	1.90 <sup>a</sup>	2.29 <sup>a</sup>	31.11 <sup>b</sup>	
Mannose	66.59 <sup>b</sup>	28.87 <sup>a</sup>	240.91 <sup>c</sup>	
Sucrose	404.63 <sup>b</sup>	243.74 <sup>a</sup>	792.80 <sup>c</sup>	
Tagatose	179.38 <sup>b</sup>	52.78 <sup>a</sup>	410.22 <sup>c</sup>	
Turanose	1.23 <sup>c</sup>	0.27 <sup>b</sup>	0.09 <sup>a</sup>	
Levoglucosan	8.54 <sup>a</sup>	19.95 <sup>c</sup>	11.24 <sup>b</sup>	
Glyceric acid	5.89 <sup>a</sup>	4.15 <sup>a</sup>	9.95 <sup>b</sup>	<b>Sugar acid</b>
Galactinol	7.48 <sup>b</sup>	0.75 <sup>a</sup>	14.71 <sup>c</sup>	
Glycerol	8.38 <sup>a</sup>	9.41 <sup>a</sup>	31.16 <sup>b</sup>	<b>Sugar alcohol</b>
Myoinositol	52.02 <sup>c</sup>	26.33 <sup>a</sup>	45.41 <sup>b</sup>	
Sorbitol	59.40 <sup>a</sup>	137.32 <sup>b</sup>	60.22 <sup>a</sup>	
Urea	5.97	2.61	4.85	
Silamine	0.47 <sup>a</sup>	2.06 <sup>b</sup>	8.94 <sup>c</sup>	<b>Amine</b>
Hydroxylamine	120.87 <sup>a</sup>	181.72 <sup>b</sup>	126.36 <sup>a</sup>	
Urea	5.97	2.61	4.85	
Palmitoleic acid	4.10 <sup>b</sup>	2.64 <sup>a</sup>	6.20 <sup>c</sup>	
Palmitic acid	30.66	32.40	29.09	<b>Fatty acid</b>
Oleic acid	1.72 <sup>a</sup>	1.45 <sup>a</sup>	3.18 <sup>b</sup>	
Stearic acid	18.08 <sup>b</sup>	18.41 <sup>b</sup>	15.47 <sup>a</sup>	
Glucuronic acid $\gamma$ -lactone	10.60 <sup>b</sup>	4.72 <sup>a</sup>	22.54 <sup>c</sup>	<b>Glycan</b>
Glyceryl-glycoside	11.20 <sup>a</sup>	15.50 <sup>b</sup>	44.31 <sup>c</sup>	<b>Glycoside</b>
Phosphoric acid	32.98	48.08	40.04	<b>Inorganic acid</b>


691 Different letters along the rows indicate statistical differences with  $P \leq 0.05$  (LSD's test). N=3.

692 **Table 5**

693 Pathway analysis: result from “Pathway Analysis” carried on the concentrations of metabolite  
 694 identified in *Echinochloa crus-galli* treated with water plus adjuvant (WA), caraway (WAC5.0)  
 695 and peppermint (WAP5.0) essential oils. In the table are reported the results obtained through  
 696 the ingenuity pathway analysis carried out with MetPa.

Pathways	T Cmpd	Hits	WA vs	WA vs	Impact
			WAC5.0	WAP5.0	
			Raw p	Raw p	
Alanine, aspartate and glutamate metabolism	21	7	0.021001	2.72E-01	0.74658
Glycine, serine and threonine metabolism	29	4	//	6.20E-01	0.53477
Galactose metabolism	26	9	1.04E-01	0.0001071	0.51278
Isoquinoline alkaloid biosynthesis	6	1	//	1.65E-01	0.5
Tyrosine metabolism	18	2	//	4.41E-01	0.27273
Inositol phosphate metabolism	17	1	0.0001203	0.048818	0.24503
Glyoxylate and dicarboxylate metabolism	17	4	0.0032152	0.00069149	0.23944
Citrate cycle (TCA cycle)	20	3	0.0093208	8.05E-01	0.17418
Methane metabolism	11	2	//	0.00056205	0.16667
Arginine and proline metabolism	37	6	0.0089378	2.58E-01	0.14946
Starch and sucrose metabolism	25	4	1.51E-02	4.42E-01	0.11156
Glycerolipid metabolism	14	2	//	0.014229	0.09402
Aminoacyl-tRNA biosynthesis	67	12	0.043696	0.00037297	0.09302
Valine, leucine and isoleucine biosynthesis	26	4	//	//	0.03645
Glutathione metabolism	26	2	0.003328	7.52E-01	0.03345

697 T Cmpd: the total number of compounds in the pathway; Hits: is the actually matched number  
 698 from the uploaded data; Raw P: is the original p value calculated from the enrichment analysis;  
 699 Impact: is the pathway impact value calculated from pathway topology analysis.



700

701 **Fig. 1. Leaf injuries and effects of different doses of adjuvant and essential oils on maize**

702 **biomass:** The average leaf injuries (A) and plant biomass (B) of maize sprayed in the stage of


703 4-6 leaves with the oil-in-water emulsions containing caraway or peppermint essential oil and

704 commercial adjuvant in the concentrations of 2.5 % or 5.0 %. Different letters refer to

705 significant differences between means, as separated by post-hoc Tukey HSD test.

706 Abbreviations: W – water; A – adjuvant; C – caraway oil; P – peppermint oil. The bars represent

707 mean value  $\pm$  standard error; N = 10.



708

709 **Fig. 2. Leaf injuries and effects of different doses of adjuvant and essential oils on *E. crus-***

710 ***galli* biomass:** The average leaf injuries (A) and plant biomass (B) of *E. crus-galli* sprayed in

711 the stage of 3-4 leaves with the oil-in-water emulsions containing caraway or peppermint


712 essential oil and commercial adjuvant in the concentrations of 2.5 % or 5.0 %. Different letters

713 refer to significant differences between means, as separated by post-hoc Tukey HSD test.

714 Abbreviations: W – water; A – adjuvant; C – caraway oil; P – peppermint oil. The bars represent


715 mean value  $\pm$  standard error; N = 10.






716

717 **Fig. 3. Pseudo-colour images of PSII parameters after the treatments of *Z. mays* with**  
 718 **essential oils:** The selected parameters of chlorophyll *a* fluorescence of maize second leaf 48  
 719 hours after spraying with water (control) or with the oil-in-water emulsions containing caraway  
 720 essential oil and commercial adjuvant in the concentrations of 2.5 % or 5.0 %. A conventional  
 721 color scale for the comparisons is on the right. Abbreviations: W – water; A – adjuvant; C –  
 722 caraway oil; Fv/Fm - maximum quantum yield of photosystem II photochemistry parameter;  
 723 NPQ – non-photochemical quenching; Rfd –fluorescence decrease ratio. N = 5.



724

725 **Fig. 4. Pseudo-colour images of PSII parameters after the treatments of *Z. mays* with**  
 726 **essential oils:** The selected parameters of chlorophyll *a* fluorescence of maize second leaf 48  
 727 hours after spraying with water (control) or with the oil-in-water emulsions containing  
 728 peppermint essential oil and commercial adjuvant in the concentrations of 2.5 % or 5.0 %. A  
 729 conventional color scale for the comparisons is on the right. Abbreviations: W – water; A –  
 730 adjuvant; P – peppermint oil; Fv/Fm - maximum quantum yield of photosystem II  
 731 photochemistry parameter; NPQ – non-photochemical quenching; Rfd – fluorescence decrease  
 732 ratio. N = 5.



733

734 **Fig. 5. Pseudo-colour images of PSII parameters after the treatments of *E. crusgalli* with**

735 **essential oils:** The selected parameters of chlorophyll *a* fluorescence of *E. crusgalli* second leaf


736 48 hours after spraying with water (control) or with the oil-in-water emulsions containing

737 caraway essential oil and commercial adjuvant in the concentrations of 2.5 % or 5.0 %. A

738 conventional color scale for the comparisons is on the right. Abbreviations: W – water; A –

739 adjuvant; C – caraway oil; Fv/Fm - maximum quantum yield of photosystem II photochemistry

740 parameter; NPQ – non-photochemical quenching; Rfd –fluorescence decrease ratio. N = 5.



741

742 **Fig. 6. Pseudo-colour images of PSII parameters after the treatments of *E. crus-galli* with**

743 **essential oils:** The selected parameters of chlorophyll *a* fluorescence of *E. crus-galli* second

744 leaf 48 hours after spraying with water (control) or with the oil-in-water emulsions containing


745 peppermint essential oil and commercial adjuvant in the concentrations of 2.5 % or 5.0 %. A

746 conventional color scale for the comparisons is on the right. Abbreviations: W – water; A –

747 adjuvant; P – peppermint oil; Fv/Fm - maximum quantum yield of photosystem II


748 photochemistry parameter; NPQ – non-photochemical quenching; Rfd – fluorescence decrease

749 ratio. N = 5.



750


751 **Fig. 7. Effects of essential oils on *Z. mays* photosystem II parameters:** A mean gray values  
 752 for the selected parameters of chlorophyll *a* fluorescence of maize. Different letters refer to  
 753 significant differences between means, as separated by post-hoc Tukey HSD test. Line  
 754 represents a mean value, box – a mean value ± standard error, and whiskers – a mean value ±  
 755 2\*standard deviation. N = 5. Abbreviations: W – water; A – adjuvant; C – caraway oil; P –  
 756 peppermint oil; Fv/Fm - maximum quantum yield of photosystem II photochemistry parameter;  
 757 NPQ – non-photochemical quenching; Rfd – fluorescence decrease ratio.



758 - Mean; □ Mean SE; I Mean 2\*SD

758

759 **Fig. 8. Effects of essential oils on *E. crus-galli* photosystem II parameters:** A mean gray  
 760 values for the selected parameters of chlorophyll *a* fluorescence of *E. crus-galli*. Different  
 761 letters refer to significant differences between means, as separated by post-hoc Tukey HSD test.  
 762 Line represents a mean value, box – a mean value ± standard error, and whiskers – a mean value  
 763 ± 2\*standard deviation. N = 5. Abbreviations: W – water; A – adjuvant; C – caraway oil; P –  
 764 peppermint oil; Fv/Fm - maximum quantum yield of photosystem II photochemistry parameter;  
 765 NPQ – non-photochemical quenching; Rfd – fluorescence decrease ratio.




766

767 **Fig. 9. Effects of essential oils on *Z. mays* metabolome:** Principal component analysis (PCA)  
 768 carried on the metabolite identified in maize plants 48 hours after spraying with water (control)  
 769 or with the oil-in-water emulsions containing caraway (WAC5.0) or peppermint (WAP5.0)  
 770 essential oil and commercial adjuvant in the concentrations of 5.0 %. A) PCA analysis model  
 771 scores A) and loading plot B) of metabolite profile of control plants [red dots (0) - WA; green  
 772 dots (1) – WAC5.0; blue dots (3) – WAP5.0]. Both score and loading plots were generated  
 773 using the first two PCs, PC1 vs PC2, with the explained variances shown in brackets; C)

774 Overlay heat map of metabolite profiles in seedlings exposed to caraway (WAC5.0) and  
775 peppermint (WAP5.0) essential oils. Each square represents the effect of the essential oils on  
776 the amount of every metabolite using a false-color scale. Red and green regions indicate  
777 increase or decrease of metabolite content, respectively. WA (1-3), indicate control replicates;  
778 WAC5.0 (1-3) indicate the replicates of seedlings treated with caraway essential oils; WAP5.0  
779 (1-3) C-24 and C-48 indicate the replicates of seedlings treated with peppermint essential oils.  
780 N=3.  
781





782

783 **Fig. 10. Effects of essential oils on *E.crus-galli* metabolome:** Principal component analysis  
 784 carried (PCA) on the metabolite identified in *E. crus-galli* plants 48 hours after spraying with  
 785 water (control) or with the oil-in-water emulsions containing caraway (WAC5.0) or peppermint  
 786 (WAP5.0) essential oil and commercial adjuvant in the concentrations of 5.0 %. A) PCA model  
 787 scores A) and loading plot B) of metabolite profile of control plants [red dots (0) - WA; green  
 788 dots (1) – WAC5.0; blue dots (3) – WAP5.0]. Both score and loading plots were generated  
 789 using the first two PCs, PC1 vs PC2, with the explained variances shown in brackets; C)

790 Overlay heat map of metabolite profiles in seedlings exposed to caraway (WAC5.0) and  
791 peppermint (WAP5.0) essential oils. Each square represents the effect of the essential oils on  
792 the amount of every metabolite using a false-color scale. Red and green regions indicate  
793 increase or decrease of metabolite content, respectively. WA (1-3), indicate control replicates;  
794 WAC5.0 (1-3) indicate the replicates of seedlings treated with caraway essential oils; WAP5.0  
795 (1-3) C-24 and C-48 indicate the replicates of seedlings treated with peppermint essential oils.  
796 N=3.