

# A high-order nonlinear Schrödinger equation with the weak non-local nonlinearity and its optical solitons

K. Hosseini<sup>a</sup>, K. Sadri<sup>a</sup>, M. Mirzazadeh<sup>b</sup>, Y.M. Chu<sup>c,\*</sup>, A. Ahmadian<sup>d,e</sup>, B.A. Pansera<sup>e</sup>, S. Salahshour<sup>f,\*</sup>

<sup>a</sup> Department of Mathematics, Rasht Branch, Islamic Azad University, Rasht, Iran

<sup>b</sup> Department of Engineering Sciences, Faculty of Technology and Engineering, East of Guilan, University of Guilan, P.C. 44891-63157 Rudsar-Vajargah, Iran

<sup>c</sup> Department of Mathematics, Huzhou University, Huzhou 313000, PR China

<sup>d</sup> Institute of IR 4.0, The National University of Malaysia, Bangi, 43600 UKM, Selangor, Malaysia

<sup>e</sup> Department of Law, Economics and Human Sciences & Decisions Lab, Mediterranean University of Reggio Calabria, Reggio Calabria 89125, Italy

<sup>f</sup> Faculty of Engineering and Natural Sciences, Bahcesehir University, Istanbul, Turkey

## ARTICLE INFO

### Keywords:

High-order nonlinear Schrödinger equation  
Weak non-local nonlinearity  
Modified Kudryashov method  
Optical solitons

## ABSTRACT

The present paper explores a high-order nonlinear Schrödinger equation in a non-Kerr law media with the weak non-local nonlinearity describing solitons' propagation through nonlinear optical fibers. To this end, the real and imaginary parts of the model are firstly extracted using a wave variable transformation. The modified Kudryashov method and symbolic computations are then adopted to successfully retrieve optical solitons of the model. The results presented in the current study demonstrate the great performance of the modified Kudryashov method in handling high-order nonlinear Schrödinger equations.

## Introduction

The extent of published papers in the last two decades with the subject of "Soliton Solutions of Nonlinear PDEs" reveals the importance of such an important area of research. Particularly, the spread of symbolic computation packages led to emerging a group of systematic methods to retrieve soliton solutions of nonlinear PDEs. Among recently introduced methods, the new Kudryashov method [1–4] has gained a significant interest because of its potential in dealing with nonlinear PDEs. The new Kudryashov method considers the solution of a nonlinear ODE like

$$O(U(\epsilon), U'(\epsilon), U''(\epsilon), \dots) = 0, \quad (1)$$

as the following finite series

$$U(\epsilon) = a_0 + \sum_{i=1}^N a_i K^i(\epsilon), \quad a_N \neq 0, \quad (2)$$

where  $a_i$  ( $0 \leq i \leq N$ ) are found later,  $N$  is derived through applying the balance principle, and  $K(\epsilon)$  satisfies the following equation

$$\begin{aligned} (K'(\epsilon))^2 &= K^2(\epsilon)(1 - \eta K^2(\epsilon)), \quad K(\epsilon) \\ &= \frac{4A}{(4A^2 - \eta)\sinh(\epsilon) + (4A^2 + \eta)\cosh(\epsilon)}, \quad \eta = 4AB, \end{aligned}$$

where  $A$  and  $B$  are free parameters. The impressive performance of the new Kudryashov method has persuaded Hosseini and his collaborators to frequently adopt such a distinguished method in their works. More information can be found in [5–8].

There are many nonlinear PDEs to describe the pulse propagation in optical fibers. The Triki–Biswas equation [9–11], the Fokas–Lenells equation [12–14], the Biswas–Arshed equation [15–17], and the Kudryashov equation [18–20] are the most commonly used models that have been studied by a group of academic researchers. In this paper, the following high-order nonlinear Schrödinger equation with the weak non-local nonlinearity [21]

$$\begin{aligned} i \frac{\partial u}{\partial t} + i c_1 \frac{\partial^3 u}{\partial x^3} + c_2 \frac{\partial^4 u}{\partial x^4} + \left( c_3 |u|^2 + c_4 |u|^4 + c_5 \frac{\partial^2 |u|^2}{\partial x^2} \right) u - i \left( c_6 \frac{\partial u}{\partial x} + c_7 \frac{\partial |u|^2 u}{\partial x} + c_8 u \frac{\partial |u|^2}{\partial x} \right) \\ = 0, \end{aligned} \quad (3)$$

\* Corresponding authors.

E-mail addresses: [chuyuming@zjhu.edu](mailto:chuyuming@zjhu.edu) (Y.M. Chu), [soheil.salahshour@eng.bau.edu.tr](mailto:soheil.salahshour@eng.bau.edu.tr) (S. Salahshour).

is considered. In Eq. (3),  $u$  is a complex-valued function, the first term is the linear temporal evolution,  $c_1$  and  $c_2$  are the coefficients of the third order and fourth order dispersions,  $c_3$  and  $c_4$  are the coefficients of the parabolic nonlinearity while  $c_5$  is the coefficient of the non-local nonlinearity,  $c_6$  is the coefficient of the inter-modal dispersion,  $c_7$  is the coefficient of the self-steepening term, and  $c_8$  is the coefficient of the nonlinear dispersion. To inform the readers, new solitons and other solutions of the model (3) have been extracted in [21] using the unified Riccati expansion method. Hosseini et al. [22] obtained soliton solutions of special cases of Eq. (3) using the new Kudryashov method.

The authors' concern in the present paper is employing the modified Kudryashov method to look for soliton solutions of the above high-order nonlinear Schrödinger equation with the weak non-local nonlinearity. The modified Kudryashov method benefits from utilizing the following finite series [23,24]

$$U(\epsilon) = a_0 + \sum_{i=1}^N \left( \frac{K(\epsilon)}{1 + K^2(\epsilon)} \right)^{i-1} \left( a_i \frac{K(\epsilon)}{1 + K^2(\epsilon)} + b_i \frac{1 - K^2(\epsilon)}{1 + K^2(\epsilon)} \right), a_N \text{ or } b_N \neq 0, \tag{4}$$

as the solution of Eq. (1). Such a representation provides a group of soliton solutions that are different from those obtained using its classical version. Recently, Hosseini et al. [24] used successfully the modified Kudryashov method to derive optical solitons of an integrable nonlinear Schrödinger equation in (2 + 1) dimensions. More papers have been listed in [25–37].

The remainder of this paper is as follows: In Section 2, a wave variable transformation is formally adopted to extract the real and imaginary parts of the model. In Section 3, the modified Kudryashov method and symbolic computations are utilized to retrieve optical solitons of the model. The last section presents a summary of the results.

**The governing model and its reduced form**

To obtain the reduced form of the governing model with the weak non-local nonlinearity, a wave variable transformation is considered as

$$u(x, t) = U(\epsilon)e^{i(-\kappa x + \mu t)}, \epsilon = x - vt, \tag{5}$$

where  $v$ ,  $\kappa$ , and  $\mu$  are the soliton velocity, the soliton wave number, and the soliton frequency. Substituting Eq. (5) into the model (3) and separating the real and imaginary parts results in

$$u_1(x, t) = \frac{a_1 A \left( \sinh \left( x + \left( \frac{c_1^3 + 8c_2^2 c_6}{8c_2^2} \right) t \right) + \cosh \left( x + \left( \frac{c_1^3 + 8c_2^2 c_6}{8c_2^2} \right) t \right) \right)}{2A^2 \sinh \left( x + \left( \frac{c_1^3 + 8c_2^2 c_6}{8c_2^2} \right) t \right) \cosh \left( x + \left( \frac{c_1^3 + 8c_2^2 c_6}{8c_2^2} \right) t \right) + 2A^2 \left( \cosh \left( x + \left( \frac{c_1^3 + 8c_2^2 c_6}{8c_2^2} \right) t \right) \right)^2 - A^2 + 1} \times e^{i \left( \frac{-c_1}{4c_2} x + \mu t \right)}, \tag{10}$$

$$c_2 \frac{d^4 U(\epsilon)}{d\epsilon^4} + (3\kappa c_1 - 6\kappa^2 c_2) \frac{d^2 U(\epsilon)}{d\epsilon^2} + (\kappa^4 c_2 - \kappa^3 c_1 - \kappa c_6 - \mu) U(\epsilon) + 2c_5 U(\epsilon) \left( \frac{dU(\epsilon)}{d\epsilon} \right)^2 + 2c_5 \frac{d^2 U(\epsilon)}{d\epsilon^2} U^2(\epsilon) + (c_3 - \kappa c_7) U^3(\epsilon) + c_4 U^5(\epsilon) = 0, \tag{6}$$

$$(c_1 - 4\kappa c_2) \frac{d^3 U(\epsilon)}{d\epsilon^3} + (4\kappa^3 c_2 - 3\kappa^2 c_1 - v - c_6) \frac{dU(\epsilon)}{d\epsilon} - (3c_7 + 2c_8) \frac{dU(\epsilon)}{d\epsilon} U^2(\epsilon) = 0. \tag{7}$$

The linearly independent principle is employed on Eq. (7) to obtain the soliton wave number, the soliton velocity, and the parameter  $c_8$  as

$$\kappa = \frac{c_1}{4c_2}, v = -\frac{c_1^3 + 8c_2^2 c_6}{8c_2^2}, c_8 = -\frac{3c_7}{2}.$$

Now, one can rewrite Eq. (6) as follows

$$c_2 \frac{d^4 U(\epsilon)}{d\epsilon^4} + \frac{3c_1^2}{8c_2} \frac{d^2 U(\epsilon)}{d\epsilon^2} - \left( \frac{3c_1^4}{256c_2^3} + \frac{c_1 c_6}{4c_2} + \mu \right) U(\epsilon) + 2c_5 U(\epsilon) \left( \frac{dU(\epsilon)}{d\epsilon} \right)^2 + 2c_5 \frac{d^2 U(\epsilon)}{d\epsilon^2} U^2(\epsilon) - \left( \frac{c_1 c_7}{4c_2} - c_3 \right) U^3(\epsilon) + c_4 U^5(\epsilon) = 0. \tag{8}$$

**The governing model and its optical solitons**

Due to the terms  $\frac{d^4 U(\epsilon)}{d\epsilon^4}$  and  $U^5(\epsilon)$ , one can find the balance number as  $N = 1$ . Consequently, the solution of Eq. (8) can be written as

$$U(\epsilon) = a_0 + a_1 \frac{K(\epsilon)}{1 + K^2(\epsilon)} + a_2 \frac{1 - K^2(\epsilon)}{1 + K^2(\epsilon)}, a_1 \text{ or } a_2 \neq 0, \tag{9}$$

where  $a_0$ ,  $a_1$ , and  $a_2$  are found later. By substituting Eq. (9) into Eq. (8), we find a system of nonlinear algebraic equations that its solution yields the following cases:

**Case 1..**  $B = 0$ ,

$$a_0 = 0,$$

$$a_2 = 0,$$

$$c_2 = -\frac{1}{384} a_1^4 c_4 + \frac{1}{16} a_1^2 c_5,$$

$$c_3 = -\frac{5a_1^8 c_4^2 - 144a_1^6 c_4 c_5 + 576a_1^4 c_5^2 + 2304a_1^2 c_1 c_7 + 27648c_1^2}{24a_1^4 (a_1^2 c_4 - 24c_5)},$$

$$\mu = -(a_1^{16} c_4^4 - 96a_1^{14} c_4^3 c_5 + 3456a_1^{12} c_4^2 c_5^2 - 55296a_1^{10} c_4 c_5^3 + 55296a_1^8 c_1^2 c_4^2 - 36864a_1^8 c_1 c_4^2 c_6 + 331776a_1^8 c_5^4 - 2654208a_1^6 c_1^2 c_4 c_5 + 1769472a_1^6 c_1 c_4 c_5 c_6 + 31850496a_1^4 c_1^2 c_5^2 - 21233664a_1^4 c_1 c_5^2 c_6 - 254803968c_1^4) / (384a_1^6 (a_1^2 c_4 - 24c_5)^3)$$


Now, based on the Case 1, the following soliton solution to the model (3) with the weak non-local nonlinearity is derived

where

$$c_2 = -\frac{1}{384} a_1^4 c_4 + \frac{1}{16} a_1^2 c_5,$$

$$c_3 = -\frac{5a_1^8 c_4^2 - 144a_1^6 c_4 c_5 + 576a_1^4 c_5^2 + 2304a_1^2 c_1 c_7 + 27648c_1^2}{24a_1^4 (a_1^2 c_4 - 24c_5)},$$

$$c_8 = -\frac{3}{2} c_7,$$



**Fig. 1.** Three-dimensional and density graphs of  $|u_1(x, t)|$  (namely Eq. (10)) when  $A = 1, c_4 = 0.1, c_5 = 1, c_6 = 0.1, a_1 = 1$  and (a)  $c_1 = 0.1$ ; (b)  $c_1 = 0.175$ .

$$\begin{aligned} \mu = & -(a_1^{16}c_4^4 - 96a_1^{14}c_4^3c_5 + 3456a_1^{12}c_4^2c_5^2 - 55296a_1^{10}c_4c_5^3 \\ & + 55296a_1^8c_1^2c_4^2 - 36864a_1^8c_1c_4^2c_6 + 331776a_1^8c_5^4 \\ & - 2654208a_1^6c_1^2c_4c_5 + 1769472a_1^6c_1c_4c_5c_6 + 31850496a_1^4c_1^2c_5^2 \\ & - 21233664a_1^4c_1c_5^2c_6 - 254803968c_1^4)/(384a_1^6(a_1^2c_4 - 24c_5)^3) \end{aligned}$$

$$\begin{aligned} u_2(x, t) = & \frac{a_2 \left( A^4 + 4A^2 \sinh \left( x + \left( \frac{c_1^3 + 8c_2^2c_6}{8c_2^2} \right) t \right) \cosh \left( x + \left( \frac{c_1^3 + 8c_2^2c_6}{8c_2^2} \right) t \right) - 1 \right)}{A^4 + 4A^2 \left( \cosh \left( x + \left( \frac{c_1^3 + 8c_2^2c_6}{8c_2^2} \right) t \right) \right)^2 - 2A^2 + 1} \\ & \times e^{i \left( -\frac{c_1}{4c_2}x + \mu t \right)}, \end{aligned} \tag{11}$$

**Case 2..**  $B = 0,$

$$a_0 = 0,$$

$$a_1 = 0,$$

$$c_2 = -\frac{1}{24}a_2^4c_4 - \frac{1}{4}a_2^2c_5,$$

$$c_3 = -\frac{5a_2^8c_4^2 + 36a_2^6c_4c_5 + 36a_2^4c_5^2 + 18a_2^2c_1c_7 - 54c_1^2}{3a_2^4(a_2^2c_4 + 6c_5)},$$

$$\begin{aligned} \mu = & -2(a_2^{16}c_4^4 + 21a_2^{14}c_4^3c_5 + 162a_2^{12}c_4^2c_5^2 + 540a_2^{10}c_4c_5^3 - 27a_2^8c_1^2c_4^2 \\ & - 9a_2^8c_1c_4^2c_6 + 648a_2^8c_5^4 - 324a_2^6c_1^2c_4c_5 - 108a_2^6c_1c_4c_5c_6 - 972a_2^4c_1^2c_5^2 \\ & - 324a_2^4c_1c_5^2c_6 - 243c_1^4)/(3a_2^6(a_2^2c_4 + 6c_5)^3) \end{aligned}$$

Now, based on the **Case 2**, the following soliton solution to the model (3) with the weak non-local nonlinearity is obtained

where

$$c_2 = -\frac{1}{24}a_2^4c_4 - \frac{1}{4}a_2^2c_5,$$


$$c_3 = -\frac{5a_2^8c_4^2 + 36a_2^6c_4c_5 + 36a_2^4c_5^2 + 18a_2^2c_1c_7 - 54c_1^2}{3a_2^4(a_2^2c_4 + 6c_5)},$$

$$c_8 = -\frac{3}{2}c_7,$$

$$\begin{aligned} \mu = & -2(a_2^{16}c_4^4 + 21a_2^{14}c_4^3c_5 + 162a_2^{12}c_4^2c_5^2 + 540a_2^{10}c_4c_5^3 - 27a_2^8c_1^2c_4^2 \\ & - 9a_2^8c_1c_4^2c_6 + 648a_2^8c_5^4 - 324a_2^6c_1^2c_4c_5 - 108a_2^6c_1c_4c_5c_6 - 972a_2^4c_1^2c_5^2 \\ & - 324a_2^4c_1c_5^2c_6 - 243c_1^4)/(3a_2^6(a_2^2c_4 + 6c_5)^3) \end{aligned}$$

**Case 3..**  $B = 0,$

$$a_0 = 0,$$



**Fig. 2.** Three-dimensional and density graphs of  $|u_1(x, t)|$  (namely Eq. (10)) when  $A = 1, c_1 = 0.1, c_4 = 0.1, c_5 = 1, a_1 = 1$  and (a)  $c_6 = 0.1$ ; (b)  $c_6 = 0.175$ .

$$a_1 = \pm \sqrt{\frac{12c_1^2 - 352c_2^2}{3c_1^2 + 8c_2^2}} a_2,$$

$$c_3 = \frac{72a_2^2 c_1 c_2^2 c_7 - 9c_1^4 - 174c_1^2 c_2^2 - 400c_2^4}{288a_2^2 c_2^3},$$

$$c_4 = \frac{(3c_1^2 + 8c_2^2)^2}{384a_2^4 c_2^3},$$

$$c_5 = -\frac{(3c_1^2 - 40c_2^2)(3c_1^2 + 8c_2^2)}{576a_2^2 c_2^3},$$

$$\mu = -\frac{45c_1^4 + 1104c_1^2 c_2^2 + 576c_1 c_2^2 c_6 + 2816c_2^4}{2304c_2^3}.$$

Now, based on the Case 3, the following exact solutions to the model (3) with the weak non-local nonlinearity are derived

$$\times e^{i \left( \frac{-c_1 x - 45c_1^4 + 1104c_1^2 c_2^2 + 576c_1 c_2^2 c_6 + 2816c_2^4}{2304c_2^3} t \right)},$$

where

$$c_3 = \frac{72a_2^2 c_1 c_2^2 c_7 - 9c_1^4 - 174c_1^2 c_2^2 - 400c_2^4}{288a_2^2 c_2^3},$$

$$c_4 = \frac{(3c_1^2 + 8c_2^2)^2}{384a_2^4 c_2^3},$$

$$c_5 = -\frac{(3c_1^2 - 40c_2^2)(3c_1^2 + 8c_2^2)}{576a_2^2 c_2^3},$$

$$c_8 = -\frac{3}{2}c_7.$$

To illustrate the dynamical behavior of Eqs. (10) and (11), the au-

$$u_{3,4}(x, t) = \frac{a_2 \left( (A^2 + 1) \sinh \left( x + \left( \frac{c_1^3 + 8c_2^2 c_6}{8c_2^2} \right) t \right) + (A^2 - 1) \cosh \left( x + \left( \frac{c_1^3 + 8c_2^2 c_6}{8c_2^2} \right) t \right) \pm 2 \sqrt{\frac{3c_1^2 - 88c_2^2}{3c_1^2 + 8c_2^2}} A \right)}{(A^2 - 1) \sinh \left( x + \left( \frac{c_1^3 + 8c_2^2 c_6}{8c_2^2} \right) t \right) + (A^2 + 1) \cosh \left( x + \left( \frac{c_1^3 + 8c_2^2 c_6}{8c_2^2} \right) t \right)}$$

thors have considered Figs. 1-4. Eq. (10) has been plotted in Fig. 1 for  $A = 1, c_4 = 0.1, c_5 = 1, c_6 = 0.1, a_1 = 1$  and different values of the parameter  $c_1$ , namely (a)  $c_1 = 0.1$ ; (b)  $c_1 = 0.175$ . This equation also is




Fig. 3. Three-dimensional and density graphs of  $|u_2(x,t)|$  (namely Eq. (11)) when  $A = 1, c_4 = 0.1, c_5 = 1, c_6 = 0.1, a_2 = 1$  and (a)  $c_1 = 0.1$ ; (b)  $c_1 = 0.4$ .




Fig. 4. Three-dimensional and density graphs of  $|u_2(x,t)|$  (namely Eq. (11)) when  $A = 1, c_1 = 0.1, c_4 = 0.1, c_5 = 1, a_2 = 1$  and (a)  $c_6 = 0.1$ ; (b)  $c_6 = 0.3$ .

portrayed in Fig. 2 for  $A = 1$ ,  $c_1 = 0.1$ ,  $c_4 = 0.1$ ,  $c_5 = 1$ ,  $a_1 = 1$  and different values of the parameter  $c_6$ , namely (a)  $c_6 = 0.1$ ; (b)  $c_6 = 0.175$ . Fig. 3 shows three-dimensional and density graphs of Eq. (11) for  $A = 1$ ,  $c_4 = 0.1$ ,  $c_5 = 1$ ,  $c_6 = 0.1$ ,  $a_2 = 1$  and different values of the parameter  $c_1$ , namely (a)  $c_1 = 0.1$ ; (b)  $c_1 = 0.4$ . Three-dimensional and density graphs of this equation are also plotted in Fig. 4 for  $A = 1$ ,  $c_1 = 0.1$ ,  $c_4 = 0.1$ ,  $c_5 = 1$ ,  $a_2 = 1$  and different values of the parameter  $c_6$ , namely (a)  $c_6 = 0.1$ ; (b)  $c_6 = 0.3$ . Obviously, the first two figures show bright solitons while the last two figures indicate dark solitons. A bright soliton is featured as a localized intensity peak above a continuous wave background while a dark soliton is characterized as a localized intensity dip below a continuous wave background [38].

Remark: It is noted that soliton solutions given in this study are different from those presented in [21].

## Conclusion

The authors' concern of this paper was to analytically explore a high-order nonlinear Schrödinger equation in a non-Kerr law media with the weak non-local nonlinearity. Such a goal was formally carried out by considering a wave variable transformation and the modified Kudryashov method to successfully retrieve optical solitons of the model. The results presented in the current study demonstrated the great performance of the modified Kudryashov method in handling high-order nonlinear Schrödinger equations.

## Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

## References

- [1] Kudryashov NA. Method for finding highly dispersive optical solitons of nonlinear differential equation. *Optik* 2020;206:163550.
- [2] Kudryashov NA. Highly dispersive solitary wave solutions of perturbed nonlinear Schrödinger equations. *Appl Math Comput* 2020;371:124972.
- [3] Kudryashov NA. Highly dispersive optical solitons of the generalized nonlinear eighth-order Schrödinger equation. *Optik* 2020;206:164335.
- [4] Kudryashov NA, Antonova EV. Solitary waves of equation for propagation pulse with power nonlinearities. *Optik* 2020;217:164881.
- [5] Hosseini K, Mirzazadeh M, Ilie M, Gómez-Aguilar JF. Soliton solutions of the Sasa-Satsuma equation in the monomode optical fibers including the beta-derivatives. *Optik* 2020;224:165425.
- [6] Hosseini K, Mirzazadeh M, Ilie M, Gómez-Aguilar JF. Biswas-Arshed equation with the beta time derivative: Optical solitons and other solutions. *Optik* 2020;217:164801.
- [7] Hosseini K, Salahshour S, Mirzazadeh M. Bright and dark solitons of a weakly nonlocal Schrödinger equation involving the parabolic law nonlinearity. *Optik* 2021;227:166042.
- [8] Matinfar M, Hosseini K. Optical solitons of (2+1)-dimensional nonlinear Schrödinger equation involving linear and nonlinear effects. *Optik* 2021;228:166110.
- [9] Kudryashov NA. First integrals and solutions of the traveling wave reduction for the Triki-Biswas equation. *Optik* 2019;185:275–81. <https://doi.org/10.1016/j.ijleo.2019.03.087>.
- [10] González-Gaxiola O. Numerical solution for Triki-Biswas equation by Adomian decomposition method. *Optik* 2019;194:163014.
- [11] Zhou Q, Ekici M, Sonmezoglu A. Exact chirped singular soliton solutions of Triki-Biswas equation. *Optik* 2019;181:338–42. <https://doi.org/10.1016/j.ijleo.2018.11.054>.
- [12] Biswas A. Chirp-free bright optical soliton perturbation with Fokas-Lenells equation by traveling wave hypothesis and semi-inverse variational principle. *Optik* 2018;170:431–5. <https://doi.org/10.1016/j.ijleo.2018.06.009>.
- [13] Biswas A, Ekici M, Sonmezoglu A, Alqahtani RT. Optical soliton perturbation with full nonlinearity for Fokas-Lenells equation. *Optik* 2018;165:29–34.
- [14] Krishnan EV, Biswas A, Zhou Q, Alifiras M. Optical soliton perturbation with Fokas-Lenells equation by mapping methods. *Optik* 2019;178:104–10. <https://doi.org/10.1016/j.ijleo.2018.10.017>.
- [15] Biswas A, Arshed S. Optical solitons in presence of higher order dispersions and absence of self-phase modulation. *Optik* 2018;174:452–9. <https://doi.org/10.1016/j.ijleo.2018.08.037>.
- [16] N.A. Kudryashov, Periodic and solitary waves of the Biswas-Arshed equation, 200 (2020) 163442.
- [17] Yıldırım Y. Optical solitons with Biswas-Arshed equation by sine-Gordon equation method. *Optik* 2020;223:165622.
- [18] Kudryashov NA. A generalized model for description of propagation pulses in optical fiber. *Optik* 2019;189:42–52. <https://doi.org/10.1016/j.ijleo.2019.05.069>.
- [19] Biswas A, Ekici M, Sonmezoglu A, Alshomrani AS, Belic MR. Optical solitons with Kudryashov's equation by extended trial function. *Optik* 2020;202:163290.
- [20] Biswas A, Asma M, Guggilla P, Mullick L, Moraru L, Ekici M, et al. Optical soliton perturbation with Kudryashov's equation by semi-inverse variational principle. *Phys Lett A* 2020;384:126830.
- [21] Zayed EME, El-Horbaty M, Alngar MEM. Cubic-quartic optical soliton perturbation having four laws non-linearity with a prolific integration algorithm. *Optik* 2020;220:165121.
- [22] Hosseini K, Matinfar M, Mirzazadeh M. Soliton Solutions of High-order Nonlinear Schrödinger Equations with Different Laws of Nonlinearities. *Regul Chaot Dyn* 2021;26(1):105–12. <https://doi.org/10.1134/S1560354721010068>.
- [23] Ma H-C, Zhang Z-P, Deng A-P. A new periodic solution to Jacobi elliptic functions of MKdV equation and BBM equation. *Acta Math Appl Sin Engl Ser* 2012;28(2):409–15. <https://doi.org/10.1007/s10255-012-0153-7>.
- [24] Hosseini K, Sadri K, Mirzazadeh M, Salahshour S. An integrable (2+1)-dimensional nonlinear Schrödinger system and its optical soliton solutions. *Optik* 2021;229:166247.
- [25] Alquran M, Yousef F, Alquran F, Sulaiman TA, Yusuf A. Dual-wave solutions for the quadratic-cubic conformable-Caputo time-fractional Klein-Fock-Gordon equation. *Math Comput Simul* 2021;185:62–76. <https://doi.org/10.1016/j.matcom.2020.12.014>.
- [26] Jaradat I, Alquran M. Construction of solitary two-wave solutions for a new two-mode version of the Zakharov-Kuznetsov equation. *Mathematics* 2020;8:1127.
- [27] Alquran M, Jaradat I, Ali M, Baleanu D. The dynamics of new dual-mode Kawahara equation: interaction of dual-wave solutions and graphical analysis. *Phys Scr* 2020;95:045216.
- [28] Alquran M, Jaradata I, Baleanu D. Shapes and dynamics of dual-mode Hirota-Satsuma coupled KdV equations: Exact traveling wave solutions and analysis. *Chin J Phys* 2019;58:49–56.
- [29] Alquran M, Jaradat I, Yusuf A, Sulaiman TA. Heart-cusp and bell-shaped-cusp optical solitons for an extended two-mode version of the complex Hirota model: application in optics. *Opt Quant Electron* 2021;53(1). <https://doi.org/10.1007/s11082-020-02674-1>.
- [30] Hamed AA, Abdel Kader AH, Abdel Latif MS. Solitons, rogue waves and breather solutions for the (2+1)-dimensional nonlinear Schrödinger equation with variable coefficients. *Optik* 2020;216:164768.
- [31] Abdel Kader AH, Abdel Latif MS, Baleanu D. Modulation instability and some dark and bright optical solitons in weakly nonlocal media with general polynomial law nonlinearity. *Mod Phys Lett B* 2020;34:2050061.
- [32] Abdel Kader AH, Abdel Latif MS, Zhou Q. Exact optical solitons in metamaterials with anti-cubic law of nonlinearity by Lie group method. *Opt Quant Electron* 2019;51:30.
- [33] Elhadedy H, Abdel Latif MS, Elsonbaty A, Abdel Kader AH, Liu W, Zhou Q. Bright and dark solitons in optical materials with polynomial law nonlinearity. *Proceed Roman Acad - Series A* 2019;20:37–43.
- [34] Abdel Kader AH, Abdel Latif MS. New bright and dark soliton solutions for a generalized nonlinear Schrödinger equation. *Optik* 2019;176:699–703. <https://doi.org/10.1016/j.ijleo.2018.09.039>.
- [35] H.M. Srivastava, D. Baleanu, J.A.T. Machado, M.S. Osman, H. Rezazadeh, S. Arshed, H. Günerhan, Traveling wave solutions to nonlinear directional couplers by modified Kudryashov method, *Phys Scrip* 95 (2020) 075217.
- [36] H. Rezazadeh, M. Inc, D. Baleanu, New solitary wave solutions for variants of (3+1)-dimensional Wazwaz-Benjamin-Bona-Mahony equations, *Frontiers in Physics* 8 (2020) 332.
- [37] Rezazadeh H. New solitons solutions of the complex Ginzburg-Landau equation with Kerr law nonlinearity. *Optik* 2018;167:218–27. <https://doi.org/10.1016/j.ijleo.2018.04.026>.
- [38] [https://en.wikipedia.org/wiki/Vector\\_soliton](https://en.wikipedia.org/wiki/Vector_soliton).